Centre No.					Pape	r Refer	ence			Surname	Initial(s)
Candidate			6	6	7	7	/	0	1	Signature	

Paper Reference(s)

6677/01

Edexcel GCE

Mechanics M1

Advanced/Advanced Subsidiary

Wednesday 16 May 2012 – Morning

Time: 1 hour 30 minutes

Materials required for examination	Items included with question paper
Mathematical Formulae (Pink)	Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 7 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates


You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.

©2012 Pearson Education Ltd.

P40689A





Examiner's use only

Turn over

Total



directions along the san before the collision, to	B , of mass $5m$ kg and $2m$ kg respectively, are more straight horizontal line. The particles collide direction speeds of A and B are 3 m s^{-1} and 4 m s^{-1} . A is unchanged by the collision. Immediately after A .	ectly. Immediately respectively. The
(a) Find the speed of .	B immediately after the collision.	(3)
In the collision, the ma	agnitude of the impulse exerted on A by B is 3.3 N	N s.
(b) Find the value of <i>i</i>	m.	(3)



3

2.




Figure 1

A non-uniform rod AB has length 3 m and mass 4.5 kg. The rod rests in equilibrium, in a horizontal position, on two smooth supports at P and at Q, where AP = 0.8 m and QB = 0.6 m, as shown in Figure 1. The centre of mass of the rod is at G. Given that the magnitude of the reaction of the support at P on the rod is twice the magnitude of the reaction of the support at Q on the rod, find

(a)	the magnitude	of the	reaction	of the	support	at Q	on 1	the roo	d,
-----	---------------	--------	----------	--------	---------	------	------	---------	----

(3)

(b)	the	distance	AG
(~)	ULIU	aistailee	110

(4)



Leave blank

3.




Figure 2

A box of mass 5 kg lies on a rough plane inclined at 30° to the horizontal. The box is held in equilibrium by a horizontal force of magnitude 20 N, as shown in Figure 2. The force acts in a vertical plane containing a line of greatest slope of the inclined plane. The box is in equilibrium and on the point of moving down the plane. The box is modelled

as a particle.

Find

(a) the magnitude of the normal reaction of the plane on the box,

(4)

(b) the coefficient of friction between the box and the plane.

(5)





Leave blank

- 4. A car is moving on a straight horizontal road. At time t = 0, the car is moving with speed 20 m s^{-1} and is at the point A. The car maintains the speed of 20 m s^{-1} for 25 s. The car then moves with constant deceleration 0.4 m s^{-2} , reducing its speed from 20 m s^{-1} to 8 m s^{-1} . The car then moves with constant speed 8 m s^{-1} for 60 s. The car then moves with constant acceleration until it is moving with speed 20 m s^{-1} at the point B.
 - (a) Sketch a speed-time graph to represent the motion of the car from A to B.

(3)

(b) Find the time for which the car is decelerating.

(2)

Given that the distance from A to B is 1960 m,

(c) find the time taken for the car to move from A to B.

(8)





Leave
blank

5.	A particle P is projected vertically upwards from a point A with speed u m s ⁻¹ . The point A is 17.5 m above horizontal ground. The particle P moves freely under gravity until it reaches the ground with speed 28 m s ⁻¹ .
	(a) Show that $u = 21$ (3)
	At time t seconds after projection, P is 19 m above A .
	(b) Find the possible values of t. (5)
	The ground is soft and, after P reaches the ground, P sinks vertically downwards into the ground before coming to rest. The mass of P is 4 kg and the ground is assumed to exert a constant resistive force of magnitude 5000 N on P .
	(c) Find the vertical distance that P sinks into the ground before coming to rest. (4)







Leave
blank

6.	[In this question \mathbf{i} and \mathbf{j} are horizontal unit vectors due east and due north respectively and position vectors are given with respect to a fixed origin.]			
	A ship S is moving with constant velocity $(-12\mathbf{i} + 7.5\mathbf{j})$ km h ⁻¹ .			
	(a) Find the direction in which S is moving, giving your answer as a bearing.	(3)		
	At time t hours after noon, the position vector of S is s km. When $t = 0$, $s = 40i - 6j$.			
	(b) Write down \mathbf{s} in terms of t .	(2)		
	A fixed beacon B is at the point with position vector $(7\mathbf{i} + 12.5\mathbf{j})$ km.			
	(c) Find the distance of S from B when $t = 3$	(4)		
	(d) Find the distance of S from B when S is due north of B .	(4)		





(4)

(3)

7.




Figure 3

Two particles P and Q, of mass 0.3 kg and 0.5 kg respectively, are joined by a light horizontal rod. The system of the particles and the rod is at rest on a horizontal plane. At time t=0, a constant force \mathbf{F} of magnitude 4 N is applied to Q in the direction PQ, as shown in Figure 3. The system moves under the action of this force until t=6 s. During the motion, the resistance to the motion of P has constant magnitude 1 N and the resistance to the motion of Q has constant magnitude 2 N.

Find

- (a) the acceleration of the particles as the system moves under the action of **F**, (3)
- (b) the speed of the particles at t = 6 s, (2)
- (c) the tension in the rod as the system moves under the action of **F**. (3)

At t = 6 s, **F** is removed and the system decelerates to rest. The resistances to motion are unchanged. Find

(d) the distance moved by P as the system decelerates,

(e) the thrust in the rod as the system decelerates.





Question 7 continued	Leave



	Leave
Question 7 continued	
	Q7
(Total 15 marks)	
TOTAL FOR PAPER: 75 MARKS	
END	