Centre No.					Pa	iper Re	eferenc	e			Surname	Initial(s)
Candidate No.			6	6	7	9	/	0	1	R	Signature	

Paper Reference(s)

6679/01R

Edexcel GCE

Mechanics M3

Advanced/Advanced Subsidiary

Monday 10 June 2013 – Morning

Time: 1 hour 30 minutes

Materials required for examination	Items included with question paper
Mathematical Formulae (Pink)	Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 7 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates


You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.

©2013 Pearson Education Ltd.

P42829A
W850/R6679/57570 5/5/5/5/






Turn over

Total

PEARSON

1.



Leave blank

Figure 1

A hollow right circular cone, of base radius a and height h, is fixed with its axis vertical and vertex downwards, as shown in Figure 1. A particle moves with constant speed v in a horizontal circle of radius $\frac{1}{3}a$ on the smooth inner surface of the cone.

Show that $v = \sqrt{\left(\frac{1}{3}hg\right)}$.	
((3))	(7)



2. A particle of mass 4 kg is moving along the horizontal *x*-axis under the action of a single force which acts in the positive *x*-direction. At time *t* seconds the force has magnitude

$$\left(1+3t^{\frac{1}{2}}\right)N$$

When t = 0 the particle has speed 2 m s⁻¹ in the positive *x*-direction. Find the work done by the force in the interval $0 \le t \le 4$

(7)



3.	A particle P of mass 0.5 kg is attached to one end of a light elastic spring, of natura 2 m and modulus of elasticity 20 N. The other end of the spring is attached to a fix A . The particle P is held at rest at the point B , which is 1 m vertically below A , a released.	ed point
	(a) Find the acceleration of P immediately after it is released from rest.	(4)
	The particle comes to instantaneous rest for the first time at the point C .	
	(b) Find the distance BC.	(6)



Leave	
blank	

- **4.** A particle *P* is moving along the positive *x*-axis. At time *t* seconds, $t \ge 0$, *P* is *x* metres from the origin *O* and is moving away from *O* with velocity $v \text{ m s}^{-1}$, where $v = \frac{4}{(x+2)}$. When t = 0, *P* is at *O*. Find
 - (a) the distance of *P* from *O* when t = 2

(5)

(b) the magnitude and direction of the acceleration of P when t = 2

(5)



estion 4 continued	



Leave blank

5.




Figure 2

Part of a hollow spherical shell, centre O and radius r, forms a bowl with a plane circular rim. The bowl is fixed to a horizontal surface at A with the rim uppermost and horizontal.

The point A is the lowest point of the bowl. The point B, where $\angle AOB = \alpha$ and $\tan \alpha = \frac{3}{4}$,

is on the rim of the bowl, as shown in Figure 2. A small smooth marble M is placed inside the bowl at A, and given an initial horizontal speed $\sqrt{(gr)}$. The motion of M takes place in the vertical plane OAB.

(a) Show that the speed of
$$M$$
 as it reaches B is $\sqrt{\left(\frac{3}{5}gr\right)}$.

After leaving the surface of the bowl at B, M moves freely under gravity and first strikes the horizontal surface at the point C. Given that r = 0.4 m,

(b)	find the distance AC.	
-----	-----------------------	--

(8)



estion 5 continued	

6. (a) A uniform lamina is in the shape of a quadrant of a circle of radius a. Show, by integration, that the centre of mass of the lamina is at a distance of $\frac{4a}{3\pi}$ from each of its straight edges.

(7)




Figure 3

A second uniform lamina ABCDEFA is shown shaded in Figure 3. The straight sides AC and AE are perpendicular and AC = AE = 2a. In the figure, the midpoint of AC is B, the midpoint of AE is F, and ABDF and DGEF are squares of side a. BCD is a quadrant of a circle with centre B. DGE is a quadrant of a circle with centre G.

(b) Find the distance of the centre of mass of the lamina from the side AE. (5)

The lamina is smoothly hinged to a horizontal axis which passes through E and is perpendicular to the plane of the lamina. The lamina has weight W newtons. The lamina is held in equilibrium in a vertical plane, with A vertically above E, by a horizontal force of magnitude X newtons applied at C.

(c)	Find X in terms of W.	
` /		(3)







Leave
blank

7.	Two points A and B are 4 m apart on a smooth horizontal surface. A light elastic string of natural length 0.8 m and modulus of elasticity 15 N, has one end attached to the point A . A light elastic string, of natural length 0.8 m and modulus of elasticity 10 N, has or end attached to the point B . A particle P of mass 0.2 kg is attached to the free end of each string. The particle rests in equilibrium on the surface at the point C on the straight limbetween A and B .	nt ne ch
	(a) Show that the length of AC is 1.76 m.	4)
	The particle P is now held at the point D on the line AB such that $AD = 2.16$ m. The particle is then released from rest and in the subsequent motion both strings remain taut	
	(b) Show that P moves with simple harmonic motion.	4)
	(c) Find the speed of P as it passes through the point C .	2)
	(d) Find the time from the instant when P is released from D until the instant when P first moving with speed 2 m s ⁻¹ .	is
		4)
		_
		_
		_
		_
		_
		_
		_
		_
		_







(Total 14 marks) TOTAL FOR PAPER: 75 MARKS	Q7
(Total 14 marks)	Q7
	Q7
	blan