Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	7	7	/	0	1	Signature	

Paper Reference(s)

6677/01

Edexcel GCE

Mechanics M1

Advanced/Advanced Subsidiary

Monday 13 May 2013 – Afternoon

Time: 1 hour 30 minutes

Examiner's use only

Team Leader's use only

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
8	
Total	

Materials required for examination
Mathematical Formulae (Pink)

Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature.
Check that you have the correct question paper.
Answer ALL the questions.
You must write your answer to each question in the space following the question.
Whenever a numerical value of g is required, take $g = 9.8$ m s^{-2}.
When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.
Full marks may be obtained for answers to ALL questions.
The marks for individual questions and the parts of questions are shown in round brackets: e.g. **(2)**.
There are 8 questions in this question paper. The total mark for this paper is 75.
There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.
You should show sufficient working to make your methods clear to the Examiner.
Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2013 Pearson Education Ltd.

Printer's Log. No.
P41828A
W850/R6677/57570 5/5/5/5/

P 4 1 8 2 8 A 0 1 2 8

Turn over

PEARSON

https://xtremepape.rs/

Leave blank

1. Particle P has mass 3 kg and particle Q has mass m kg. The particles are moving in opposite directions along a smooth horizontal plane when they collide directly. Immediately before the collision, the speed of P is 4 m s^{-1} and the speed of Q is 3 m s^{-1}. In the collision the direction of motion of P is unchanged and the direction of motion of Q is reversed. Immediately after the collision, the speed of P is 1 m s^{-1} and the speed of Q is 1.5 m s^{-1}.

(a) Find the magnitude of the impulse exerted on P in the collision.

(3)

(b) Find the value of m.

(3)

P 4 1 8 2 8 A 0 2 2 8

https://xtremepape.rs/

Leave blank

Question 1 continued

Q1

(Total 6 marks)

https://xtremepape.rs/

Leave blank

2. A woman travels in a lift. The mass of the woman is 50 kg and the mass of the lift is 950 kg. The lift is being raised vertically by a vertical cable which is attached to the top of the lift. The lift is moving upwards and has constant deceleration of 2 m s^{-2}. By modelling the cable as being light and inextensible, find

(a) the tension in the cable,

(3)

(b) the magnitude of the force exerted on the woman by the floor of the lift.

(3)

P 4 1 8 2 8 A 0 4 2 8

https://xtremepape.rs/

Leave blank

Question 2 continued

Q2

(Total 6 marks)

https://xtremepape.rs/

Leave blank

3.



Figure 1

A box of mass 2 kg is held in equilibrium on a fixed rough inclined plane by a rope. The rope lies in a vertical plane containing a line of greatest slope of the inclined plane. The rope is inclined to the plane at an angle α, where $\tan \alpha = \frac{3}{4}$, and the plane is at an angle of 30° to the horizontal, as shown in Figure 1. The coefficient of friction between the box and the inclined plane is $\frac{1}{3}$ and the box is on the point of slipping up the plane. By modelling the box as a particle and the rope as a light inextensible string, find the tension in the rope.

(8)

P 4 1 8 2 8 A 0 6 2 8

https://xtremepape.rs/

Leave blank

Question 3 continued

https://xtremepape.rs/

Leave blank

Question 3 continued

https://xtremepape.rs/

Leave blank

Question 3 continued

(Total 8 marks)

Q3

https://xtremepape.rs/

Leave blank

4. A lorry is moving along a straight horizontal road with constant acceleration. The lorry passes a point A with speed u m s^{-1}, ($u < 34$), and 10 seconds later passes a point B with speed 34 m s^{-1}. Given that $AB = 240$ m, find

(a) the value of u,

(3)

(b) the time taken for the lorry to move from A to the mid-point of AB.

(6)

https://xtremepape.rs/

Leave blank

Question 4 continued

https://xtremepape.rs/

Leave blank

Question 4 continued

https://xtremepape.rs/

Leave blank

Question 4 continued

Q4

(Total 9 marks)

https://xtremepape.rs/

Leave blank

5. A car is travelling along a straight horizontal road. The car takes 120 s to travel between two sets of traffic lights which are 2145 m apart. The car starts from rest at the first set of traffic lights and moves with constant acceleration for 30 s until its speed is 22 m s^{-1}. The car maintains this speed for T seconds. The car then moves with constant deceleration, coming to rest at the second set of traffic lights.

(a) Sketch, in the space below, a speed-time graph for the motion of the car between the two sets of traffic lights.

(2)

(b) Find the value of T.

(3)

A motorcycle leaves the first set of traffic lights 10 s after the car has left the first set of traffic lights. The motorcycle moves from rest with constant acceleration, a m s^{-2}, and passes the car at the point A which is 990 m from the first set of traffic lights. When the motorcycle passes the car, the car is moving with speed 22 m s^{-1}.

(c) Find the time it takes for the motorcycle to move from the first set of traffic lights to the point A.

(4)

(d) Find the value of a.

(2)

P 4 1 8 2 8 A 0 1 4 2 8

https://xtremepape.rs/

Leave blank

Question 5 continued

https://xtremepape.rs/

Leave blank

Question 5 continued

https://xtremepape.rs/

Leave blank

Question 5 continued

(Total 11 marks)

Q5

https://xtremepape.rs/

Leave blank

6. A beam AB has length 15 m. The beam rests horizontally in equilibrium on two smooth supports at the points P and Q, where $AP = 2$ m and $QB = 3$ m. When a child of mass 50 kg stands on the beam at A, the beam remains in equilibrium and is on the point of tilting about P. When the same child of mass 50 kg stands on the beam at B, the beam remains in equilibrium and is on the point of tilting about Q. The child is modelled as a particle and the beam is modelled as a non-uniform rod.

(a) (i) Find the mass of the beam.

(ii) Find the distance of the centre of mass of the beam from A.

(8)

When the child stands at the point X on the beam, it remains horizontal and in equilibrium. Given that the reactions at the two supports are equal in magnitude,

(b) find AX.

(6)

P 4 1 8 2 8 A 0 1 8 2 8

https://xtremepape.rs/

Leave blank

Question 6 continued

https://xtremepape.rs/

Leave blank

Question 6 continued

https://xtremepape.rs/

Leave blank

Question 6 continued

(Total 14 marks)

Q6

https://xtremepape.rs/

Leave blank

7. *[In this question, the horizontal unit vectors* **i** *and* **j** *are directed due east and due north respectively.]*

The velocity, $\mathbf{v}$ m s^{-1}, of a particle P at time t seconds is given by

$$\mathbf{v} = (1 - 2t)\mathbf{i} + (3t - 3)\mathbf{j}$$

(a) Find the speed of P when $t = 0$

(3)

(b) Find the bearing on which P is moving when $t = 2$

(2)

(c) Find the value of t when P is moving

(i) parallel to $\mathbf{j}$,

(ii) parallel to $(-\mathbf{i} - 3\mathbf{j})$.

(6)

P 4 1 8 2 8 A 0 2 2 2 8

https://xtremepape.rs/

Leave blank

Question 7 continued

https://xtremepape.rs/

Leave blank

Question 7 continued

https://xtremepape.rs/

Leave blank

Question 7 continued

(Total 11 marks)

Q7

https://xtremepape.rs/

Leave blank

8.



Figure 2

Two particles A and B have masses $2m$ and $3m$ respectively. The particles are attached to the ends of a light inextensible string. Particle A is held at rest on a smooth horizontal table. The string passes over a small smooth pulley which is fixed at the edge of the table. Particle B hangs at rest vertically below the pulley with the string taut, as shown in Figure 2. Particle A is released from rest. Assuming that A has not reached the pulley, find

(a) the acceleration of B,

(5)

(b) the tension in the string,

(1)

(c) the magnitude and direction of the force exerted on the pulley by the string.

(4)

P 4 1 8 2 8 A 0 2 6 2 8

https://xtremepape.rs/

Leave blank

Question 8 continued

https://xtremepape.rs/

Leave blank

Question 8 continued

Q8

(Total 10 marks)

TOTAL FOR PAPER: 75 MARKS

END

https://xtremepape.rs/