



Pearson

Mark Scheme (Results)

January 2017

Pearson Edexcel
International Advanced Subsidiary Level
In Core Mathematics C12 (WMA01)
Paper 01

edexcel 

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2017

Publications Code WMA01_01_1701_MS

All the material in this publication is copyright

© Pearson Education Ltd 2017

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

PEARSON EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 125
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \checkmark will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - d... or dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper or ag- answer given
 - \square or d... The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Core Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

Method mark for solving 3 term quadratic:

1. Factorisation

$$(x^2 + bx + c) = (x + p)(x + q), \text{ where } |pq| = |c|, \text{ leading to } x = \dots$$

$$(ax^2 + bx + c) = (mx + p)(nx + q), \text{ where } |pq| = |c| \text{ and } |mn| = |a|, \text{ leading to } x = \dots$$

2. Formula

Attempt to use the correct formula (with values for a , b and c).

3. Completing the square

$$\text{Solving } x^2 + bx + c = 0: \left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0, \quad q \neq 0, \quad \text{leading to } x = \dots$$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \rightarrow x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \rightarrow x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice.

Question Number	Scheme		Marks
1(a)	$\left(\frac{dy}{dx} = \right) \frac{3x^2}{3} - 2 \times 2x + 3$	M1: $x^n \rightarrow x^{n-1}$ or $5 \rightarrow 0$	M1A1
		A1: Any 3 of the 4 terms differentiated correctly - this could be 2 terms correct and $5 \rightarrow 0$ (allow simplified or un-simplified for this mark, including $3x^0$ for 3)	
	$\left(\frac{dy}{dx} = \right) x^2 - 4x + 3$	Cao. All 3 terms correct and simplified and on the same line and no + 0. (<u>Do not</u> allow $1x^2$ for x^2 or x^1 for x or $3x^0$ for 3). Condone poor notation e.g. omission of $dy/dx = \dots$ or if they use $y = \dots$	A1
	<p>Candidates who multiply by 3 before differentiating: e.g. $\left(\frac{x^3}{3} - 2x^2 + 3x + 5\right) \times 3 = x^3 - 6x^2 + 9x + 15 \Rightarrow \frac{dy}{dx} = 3x^2 - 12x + 9$ Scores M1A0A0 but could recover in (a) if they then divide by 3 If they persist with $\frac{dy}{dx} = 3x^2 - 12x + 9$ in (b), allow full recovery in (b)</p>		
			(3)
(b)	$x^2 - 4x + 3 = 0 \Rightarrow x = 1, 3$	M1: Attempt to solve their 3TQ from part (a) as far as $x = \dots$ (see general guidance for solving a 3TQ). If no working is shown and the roots are incorrect for their 3TQ, score M0 here but the second method mark below is still available.	M1A1
		A1: Correct values (may be implied by their inequalities e.g. a correct quadratic followed by just $x > 1$ and $x > 3$ could score M1A1 here)	
	$x < "1", x > "3"$	Chooses outside region ($x <$ their lower limit $x >$ their upper limit). Do not award simply for diagram or table.	M1
	$x < 1, x > 3$ Correct answer. Allow the correct regions separated by a comma or written separately and allow other notation e.g. $(-\infty, 1) \cup (3, \infty)$. Do not allow $1 > x > 3$ or $x < 1$ and $x > 3$ (These score M1A0). ISW if possible e.g. $x > 3, x < 1$ followed by $1 > x > 3$ can score M1A1. $x > 3, x > 1$ followed by $x > 3$ (or) $x < 1$ can score M1A1. Fully correct answer with no working scores both marks. Answers that are otherwise correct but use \leq, \geq lose final mark as would $[-\infty, 1] \cup [3, \infty]$.		A1
			(4)
			(7 marks)

Question Number	Scheme		Marks	
2 (a)	Mark (a) and (b) together			
	$(x \pm 4) \dots (y \pm 2)$	Attempts to complete the square on x and y or sight of $(x \pm 4)$ and $(y \pm 2)$. May be implied by a centre of $(\pm 4, \pm 2)$. Or if considering $x^2 + y^2 + 2gx + 2fy + c = 0$, centre is $(\pm g, \pm f)$.	M1	
	Centre $C = (4, -2)$	Correct centre (allow $x = 4, y = -2$) But not $g = \dots, f = \dots$ or $p = \dots, q = \dots$ etc.	A1	
	Correct answer scores both marks			
				(2)
(b)	$r^2 = 12 + (\pm 4)^2 + (\pm 2)^2$	<p style="text-align: center;">Must reach:</p> $r^2 = 12 + \text{their } (\pm 4)^2 + \text{their } (\pm 2)^2$ or $r = \sqrt{12 + \text{their } (\pm 4)^2 + \text{their } (\pm 2)^2}$ or if considering $x^2 + y^2 + 2gx + 2fy + c = 0,$ $r^2 = g^2 + f^2 - c$ or $r = \sqrt{g^2 + f^2 - c}$ Must clearly be identifying the radius or radius ² May be implied by a correct exact radius or awrt 5.66	M1	
	$r = \sqrt{32}$	$r = \sqrt{32}$. Accept exact equivalents such as $4\sqrt{2}$. $r = \dots$ not needed but must clearly be the radius. Do not allow $\pm\sqrt{32}$ unless minus is rejected	A1	
	Correct answer scores both marks			
				(2)
(c)	$x = 0 \Rightarrow y^2 + 4y - 12 = 0$	Correct quadratic. Allow $16 + (y + 2)^2 = 32$	B1	
	$(y + 6)(y - 2) = 0 \Rightarrow y = \dots$	Attempts to solve a 3TQ that has come from substituting $x = 0$ or $y = 0$ into the given equation or their 'changed' equation. May be implied by correct answers for their quadratic.	M1	
	$y = 2, -6$ or $(0, 2)$ and $(0, -6)$	Correct y values or correct coordinates. Accept sight of these for all 3 marks if no incorrect working seen but must clearly be y values or correct coordinates. This may be implied by the correct roots of a quadratic in y .	A1	
				(3)
			(7 marks)	

Question Number	Scheme		Marks
3(a)	$S = r\theta = 7 \times 0.8 = 5.6(\text{cm})$	M1: Uses $S = r\theta$	M1A1
		A1: 5.6 oe e.g. 28/5	
	<p>Note that if the 0.8 is converted to degrees e.g. $0.8 \times \frac{180}{\pi} = 45.8366\dots$, this angle may be rounded or truncated when attempting $\frac{45.8366\dots}{360} \times 2 \times \pi \times 7$ for the M1 so allow A1 for awrt 5.6</p>		
			(2)
(b)	$\angle POC = \frac{\pi}{2} - 0.8 = \text{awrt } 0.771$	M1: Attempts to find $\frac{\pi}{2} - 0.8$ or $\pi - \frac{\pi}{2} - 0.8$. Allow an attempt to find θ from $\theta + \frac{\pi}{2} + 0.8 = \pi$. Accept as evidence awrt 0.77	M1A1
		A1: awrt 0.771	
		<p>Answers in degrees only can score M1A0 e.g. $180 - 90 - 0.8 \times \frac{180}{\pi} (= 44.163\dots)$</p>	
			(2)
(c)	$4^2 + 5^2 - 2 \times 4 \times 5 \cos'0.771'$ or $\sqrt{4^2 + 5^2 - 2 \times 4 \times 5 \cos'0.771'}$	Correct use of the cosine rule to find CP or CP^2 . NB 0.771 radians is awrt 44 degrees. Ignore lhs for this mark and look for e.g. $4^2 + 5^2 - 2 \times 4 \times 5 \cos'0.771$ or 44'	M1
	$CP^2 = 4^2 + 5^2 - 2 \times 4 \times 5 \cos 0.771$ or $CP = \sqrt{4^2 + 5^2 - 2 \times 4 \times 5 \cos 0.771}$	A correct expression for CP or CP^2 with lhs consistent with rhs . Allow awrt 0.77 radians or awrt 44 degrees. (May be implied if a correct numerical value is used in subsequent work)	A1
	Perimeter = $4 + 5 + 2 \times 7 + '5.6' + '3.5'$	$4 + 5 + 2 \times 7 + \text{their } AQ + \text{their } CP$. Need to see all 6 lengths but may be implied by e.g. $23 + '5.6' + '3.5'$	M1
	= 32.11 (cm)	Awrt 32.11 (ignore units)	A1
			(4)
			(8 marks)

Question Number	Scheme		Marks
4 (a)	$S_9 = 54$ $\Rightarrow 54 = \frac{9}{2}(2a + 8d)$ <p style="text-align: center;">or</p> $\Rightarrow 54 = \frac{9}{2}(a + a + 8d)$	Uses a correct sum formula with $n = 9$ and $S_9 = 54$	M1
	$\Rightarrow a + 4d = 6^*$	cso	A1*
	Listing:		
	$a + a + d + a + 2d + \dots + a + 8d = 54$ $\Rightarrow 9a + 36d = 54$ <p style="text-align: center;">Scores M1 for attempting to sum 9 terms (both lines needed)</p> <p style="text-align: center;">or</p> $a + a + d + a + 2d + a + 3d + a + 4d + a + 5d + a + 6d + a + 7d + a + 8d = 54$ <p style="text-align: center;">Scores M1 on its own and then A1 if they complete correctly.</p>		
(b)	$a + 7d = \frac{1}{2}(a + 6d)$ <p style="text-align: center;">or</p> $\frac{1}{2}(a + 7d) = a + 6d$	Uses $t_8 = \frac{1}{2}t_7$ or $\frac{1}{2}t_8 = t_7$ to produce one of these equations.	M1
	$\Rightarrow 6 - 4d + 7d = \frac{1}{2}(6 - 4d + 6d)$ $\Rightarrow d = \dots$	Uses the given equation from (a) and their second linear equation in a and d and proceeds to find a value for either a or d .	M1
	$\Rightarrow d = -1.5, a = 12$	A1: Either $d = -1.5$ (oe) or $a = 12$	A1A1
		A1: Both $d = -1.5$ (oe) and $a = 12$	
	Note that use of $\frac{1}{2}t_8 = t_7$ in (b) gives $a = 30$ and $d = -6$		
			(6 marks)

Question Number	Scheme		Marks	
5 (a)(i)	$\log_3\left(\frac{x}{9}\right) = \log_3 x - \log_3 9 = y - 2$	M1: $\log_3\left(\frac{x}{9}\right) = \log_3 x - \log_3 9$ or $\log_3\left(\frac{x}{9}\right) = \log_3 x + \log_3 \frac{1}{9}$ Correct use of the subtraction rule or addition rule. Ignore the presence or absence of a base and any spurious “= 0”	M1A1	
		A1: $y - 2$		
		An answer left as $\log_3 3^{y-2}$ scores M1A0		
	Note that $\log_3\left(\frac{x}{9}\right) = \log_3 x - \log_3 9 = y - \log_3 9$ scores M1A0			
(ii)	$\log_3 \sqrt{x} = \log_3 x^{\frac{1}{2}} = \frac{1}{2} \log_3 x = \frac{1}{2} y$	$\frac{1}{2} y$ or equivalent	B1	
	(3)			
(b)	$2 \log_3\left(\frac{x}{9}\right) - \log_3 \sqrt{x} = 2 \Rightarrow 2(y-2) - \frac{1}{2} y = 2$		M1	
	Uses their answers from part (a) to create a linear equation in y (condone poor use of brackets e.g. $2(y-2) = 2y-2$ and also the slip $(y-2) - \frac{1}{2} y = 2$ for this mark)			
	$\Rightarrow y = 4$	Correct value for y.		A1
	Note that arriving at $(y-2)^2 - \frac{1}{2} y = 2$ above scores M0 (not linear) but does have a solution $y = 4$ so look out for $y = 4$ not being derived correctly.			
	$\log_3 x = 4 \Rightarrow x = 3^4$	Correct method for undoing log. Dependent on the first M		dM1
	$\Rightarrow x = 81$	cao		A1
(4)				
(7 marks)				
Alt 1 (b)	$2 \log_3\left(\frac{x}{9}\right) - \log_3 \sqrt{x} = \log_3\left(\frac{(x/9)^2}{\sqrt{x}}\right)$ or $2 \log_3\left(\frac{x}{9}\right) - \log_3 \sqrt{x} = 2 \log_3 x - 2 \log_3 9 - \log_3 \sqrt{x} = \log_3 \frac{x^2}{\sqrt{x}} + \dots$ Combines two log terms in x correctly to obtain a single log term		M1	
	$\log_3\left(\frac{(x/9)^2}{\sqrt{x}}\right) = 2$ or $\log_3\left(\frac{x^2}{\sqrt{x}}\right) = 6$	Correct equation	A1	
	$\left(\frac{(x/9)^2}{\sqrt{x}}\right) = 3^2$ or $\left(\frac{x^2}{\sqrt{x}}\right) = 3^6$	Correct method for undoing log. Dependent on the first M	dM1	
	$\Rightarrow x = 81$	cao	A1	

Alt 2 (b) Uses $x = 3^y$	$2\log_3\left(\frac{x}{9}\right) - \log_3\sqrt{x} = 2\log_3\left(\frac{3^y}{9}\right) - \log_3 3^{\frac{y}{2}} = \log_3\left(\frac{3^{\frac{3y}{2}}}{81}\right)$		M1
	Combines logs correctly		
	$\log_3\left(\frac{3^{\frac{3y}{2}}}{81}\right) = 2 \Rightarrow y = 4$	Correct value for y	A1
	$\log_3 x = 4 \Rightarrow x = 3^4$	Correct method for undoing log. Dependent on the first M	dM1
	$\Rightarrow x = 81$	cao	A1

Question Number	Scheme		Marks
6(a)(i)	$\frac{3}{2}$	Accept exact equivalents	B1
(ii)	$y=0, 3x+5=0 \Rightarrow x=-\frac{5}{3}$	M1: Sets $y=0$ and attempts to find x . Accept as evidence $3x+5=0 \Rightarrow x=..$ or awrt -1.7 A1: $x=-\frac{5}{3}$ or exact equivalent including 1.6 recurring (i.e. a clear dot over the 6)	M1A1
			(3)
(b)	Gradient $l_2 = -\frac{1}{\frac{3}{2}} = -\frac{2}{3}$	Uses $m_2 = -\frac{1}{m_1}$ to find the gradient of l_2 (may be implied by their line equation). Allow an attempt to find m_2 from $m_1 \times m_2 = -1$.	M1
	Point B has y coordinate of 4	This may be embedded within the equation of the line but must be seen in part (b).	B1
	e.g. $y - '4' = '-\frac{2}{3}'(x-1)$ or $\frac{y - '4'}{x-1} = '-\frac{2}{3}'$	A correct straight line method with a changed gradient and their point (1, '4'). There must have been attempt to find the y coordinate of B . If using $y = mx + c$, must reach as far as finding a value for c .	M1
	e.g. $y - 4 = -\frac{2}{3}(x-1)$ or $\frac{y-4}{x-1} = -\frac{2}{3}$	A correct un-simplified equation	A1
	$2x+3y-14=0$	Accept $A(2x+3y-14)=0$ where A is an integer. Terms can be in any order but must have ' $=0$ '.	A1
			(5)
Alt (b)	Gradient $l_2 = -\frac{1}{\frac{3}{2}} = -\frac{2}{3}$	Uses $m_2 = -\frac{1}{m_1}$ to find the gradient of l_2 as before	M1
	$\frac{3}{2}x + \frac{5}{2} = -\frac{2}{3}x + c$	A correct statement for $l_1 = l_2$	B1
	$x=1 \Rightarrow c = \frac{14}{3}$	Substitutes $x=1$ to find a value for c	M1
	$y = -\frac{2}{3}x + \frac{14}{3}$	Correct equation	A1
	$2x+3y-14=0$	Accept $A(2x+3y-14)=0$ where A is an integer.	A1

(c)	$y = 0 \Rightarrow 2x - 14 = 0 \Rightarrow x = 7$	Attempts to find C using $y = 0$ in the equation obtained in part (b)	M1
	Attempts Area of triangle using $\frac{1}{2} \times AC \times (y \text{ coord of } B)$ $= \frac{1}{2} \times \left('7' + ' \frac{5}{3}' \right) \times '4'$ or Attempts Area of triangle using 2 triangles $\frac{1}{2} \times \left(1 + ' \left(\frac{5}{3} \right)' \right) \times (y \text{ coord of } B) + \frac{1}{2} \times ('7' - 1) \times (y \text{ coord of } B)$ If they make a second/different attempt to find the y coordinate of B then still allow this mark.		M1
	$= \frac{52}{3}$	Area = $\frac{52}{3}$ or exact equivalent e.g. $17\frac{1}{3}$ or 17.3 recurring (i.e. a clear dot over the 3)	A1
			(3)
(11 marks)			
Way 2 6(c)	$y = 0 \Rightarrow 2x - 14 = 0 \Rightarrow x = 7$	Attempts to find C using $y = 0$ in the equation obtained in part (b)	M1
	Attempts area of triangle using $\frac{1}{2} AB \times BC = \frac{1}{2} \times \sqrt{\frac{208}{9}} \times \sqrt{52}$ A complete method for the area including correct attempts at finding AB and BC using their values.		M1
	$= \frac{52}{3}$	Area = $\frac{52}{3}$ or exact equivalent e.g. $17\frac{1}{3}$ or 17.3 recurring (i.e. a clear dot over the 3)	A1
			(3)
Way 3 6(c)	$y = 0 \Rightarrow 2x - 14 = 0 \Rightarrow x = 7$	Attempts to find C using $y = 0$ in the equation obtained in part (b)	M1
	$\frac{1}{2} \left \begin{array}{cccc} 1 & 7 & -\frac{5}{3} & 1 \\ 4 & 0 & 0 & 4 \end{array} \right = \frac{1}{2} \left -\frac{20}{3} - 28 \right $	Uses shoelace method. Must see a correct method including $\frac{1}{2}$.	M1
	$= \frac{52}{3}$	Area = $\frac{52}{3}$ or exact equivalent e.g. $17\frac{1}{3}$ or 17.3 recurring (i.e. a clear dot over the 3)	A1
			(3)

Way 4 6(c)	$y = 0 \Rightarrow 2x - 14 = 0 \Rightarrow x = 7$	Attempts to find C using $y = 0$ in the equation obtained in part (b)	M1
	$\int_{-\frac{5}{3}}^1 \left(\frac{3x}{2} + \frac{5}{2} \right) dx + \int_1^7 \left(-\frac{2x}{3} + \frac{14}{3} \right) dx$ $= \left[\frac{3x^2}{4} + \frac{5}{2}x \right]_{-\frac{5}{3}}^1 + \left[-\frac{2x^2}{6} + \frac{14}{3}x \right]_1^7$ $= \left(\frac{3}{4} + \frac{5}{2} \right) - \left(\frac{75}{36} - \frac{25}{6} \right) + \left(-\frac{49}{3} + \frac{98}{3} \right) - \left(-\frac{1}{3} + \frac{14}{3} \right)$ <p>A complete method using their values with correct integration on l_1 and their l_2: Finds the area under the given line between their $-5/3$ and 1 and adds the area under their l_2 between 1 and their 7.</p>		M1
	$= \frac{52}{3}$	Area = $\frac{52}{3}$ or exact equivalent e.g. $17\frac{1}{3}$ or 17.3 recurring (i.e. a clear dot over the 3)	A1
			(3)

Question Number	Scheme		Marks
7 (i)	$\frac{2+4x^3}{x^2} = \frac{2}{x^2} + 4x = 2x^{-2} + 4x$	Attempts to split the fraction. This can be awarded for $\frac{2}{x^2}$ or $\frac{4x^3}{x^2}$ or may be implied by the sight of one correct index e.g. px^{-2} or qx providing one of these terms is obtained correctly. So for example $\frac{2+4x^3}{x^2} = 2+4x^3+x^{-2}$ would be M0 as the x^{-2} has been obtained incorrectly.	M1
	$\int 2x^{-2} + 4x \, dx = 2 \times \frac{x^{-1}}{-1} + 4 \times \frac{x^2}{2} (+c)$	dM1: $x^n \rightarrow x^{n+1}$ on any term. Dependent on the first M. A1: At least one term correct, simplified or un-simplified. Allow powers and coefficients to be un-simplified e.g. $2 \times \frac{x^{-2+1}}{-1}$, $+4 \times \frac{x^{1+1}}{2}$	dM1A1
	$= -\frac{2}{x} + 2x^2 + c$	All correct and simplified including the + c. Accept equivalents such as $-2x^{-1} + 2x^2 + c$	A1
			(4)
There are no marks in (ii) for use of the trapezium rule – must use integration			
(ii)	$\int \left(\frac{4}{\sqrt{x}} + k \right) dx$ $= \int (4x^{-0.5} + k) dx = 4 \frac{x^{0.5}}{0.5} + kx (+c)$	M1: Integrates to obtain either $ax^{0.5}$ or kx A1: Correct integration (simplified or un-simplified). Allow powers and coefficients to be un-simplified e.g. $4 \frac{x^{-0.5+1}}{0.5}$. There is no need for + c	M1A1
	$\left[4 \frac{x^{0.5}}{0.5} + kx \right]_2^4 = 30 \Rightarrow (8\sqrt{4} + 4k) - (8\sqrt{2} + 2k) = 30$ <p>Substitutes both $x = 4$ and $x = 2$ into changed expression involving k, subtracts either way round and sets equal to 30 Condone poor use or omission of brackets when subtracting.</p>		M1
	$2k + 16 - 8\sqrt{2} = 30 \Rightarrow k = 7 + 4\sqrt{2}$	ddM1: Attempts to solve for k from a linear equation in k . Dependent upon both M's and need to have seen $\int k \, dx \rightarrow kx$. A1: $7 + 4\sqrt{2}$ or exact equivalent e.g. $7 + 2^{2.5}$, $7 + 4 \times 2^{0.5}$	ddM1A1
			(5)
			(9 marks)

Question Number	Scheme		Marks	
8(a)	$f(3) = 2(3)^3 - 5(3)^2 - 23(3) - 10$ <p style="text-align: center;">or</p> $\begin{array}{r} 2x^2 + \dots\dots\dots \\ x-3 \overline{) 2x^3 - 5x^2 - 23x - 10} \\ \dots \\ \dots \end{array}$	Attempts to calculate $f(\pm 3)$ or divides by $(x-3)$. For long division need to see minimum as shown with a constant remainder.	M1	
	(Remainder =) -70	-70	A1	
				(2)
Mark (b) and (c) together				
(b)	$f(-2) = 2(-2)^3 - 5(-2)^2 - 23(-2) - 10$ <p style="text-align: center;">Or</p> $\begin{array}{r} 2x^2 + \dots\dots\dots \\ x+2 \overline{) 2x^3 - 5x^2 - 23x - 10} \\ \dots \\ \dots \end{array}$	Attempts $f(\pm 2)$ or divides by $(x+2)$. For long division need to see minimum as shown with a constant remainder.	M1	
	Remainder = 0, hence $x+2$ is a factor	Obtains a remainder zero and makes a conclusion (not just a tick or e.g. QED). Do not need to refer to the remainder in the conclusion but a zero remainder must have been obtained. (May be seen in a preamble)	A1*	
	Note that just $f(-2) = 0$ therefore $(x+2)$ is a factor scores M0A0 as there must be some evidence of a calculation			
				(2)
(c)	$\frac{2x^3 - 5x^2 - 23x - 10}{(x+2)} = ax^2 + bx + c$	Divides $f(x)$ by $(x+2)$ or compares coefficients or uses inspection to obtain a quadratic expression with $2x^2$ as the first term.	M1	
	$2x^2 - 9x - 5$	Correct quadratic seen	A1	
	$f(x) = (x+2)(2x+1)(x-5)$ <p>dM1: Attempt to factorise their 3TQ ($2x^2 \dots$). The usual rules apply here so if $2x^2 - 9x - 5$ is factorised as $(x-5)(x+\frac{1}{2})$, this scores M0 unless the factor of 2 appears later.</p> <p>A1: $(x+2)(2x+1)(x-5)$ oe e.g. $2(x+2)(x+\frac{1}{2})(x-5)$. All factors together on one line. Must appear here and not in (d). Ignore subsequent attempts to solve.</p>		dM1A1	
SC: This is a hence question but we will allow a special case of 1100 for candidates in this part who use their graphical calculators to get roots of -2, -0.5 and 5 and write down the correct factorised form.				

Question Number	Scheme	Marks
	But note that if all that is seen is $(x+2)(x+\frac{1}{2})(x-5)$ this scores 1000	
		(4)

(d)	$3^t = '5' \Rightarrow t \log 3 = \log '5'$	Solves $3^t = k$ where $k > 0$ and follows from their (c) to obtain $t \log 3 = \log k$. Accept sight of $t = \log_3 k$ where $k > 0$ and follows from their (c)	M1
	$\Rightarrow t = \text{awrt } 1.465 \text{ only}$	$t = \text{awrt } 1.465$ and no other solutions	A1
			(2)
			(10 marks)

Question Number	Scheme	Marks	
9(a)	$f(x) = 8x^{-1} + \frac{1}{2}x - 5$ $\Rightarrow f'(x) = -8x^{-2} + \frac{1}{2}$	M1: $-8x^{-2}$ or $\frac{1}{2}$ A1: Fully correct $f'(x) = -8x^{-2} + \frac{1}{2}$ (may be un-simplified)	M1A1
	Sets $-8x^{-2} + \frac{1}{2} = 0 \Rightarrow x = 4$	M1: Sets their $f'(x) = 0$ i.e. a “changed” function (may be implied by their work) and proceeds to find x . A1: $x = 4$ (Allow $x = \pm 4$)	M1A1
	$(4, -1)$	Correct coordinates (allow $x = 4, y = -1$). Ignore their $(-4, \dots)$	A1
			(5)
	(b)(i)	$(x=) 2, 8$	$x = 2$ and $x = 8$ only . Do not accept as coordinates here.
(b)(ii)	$(4, 1)$	$(4, 1)$ or follow through on their solution in (a). Accept $(x, y+2)$ from their (x, y) . With no other points.	B1ft
(b)(iii)	$(x=) 2, \frac{1}{2}$	Both answers are needed and accept $(2, 0), (\frac{1}{2}, 0)$ here. Ignore any reference to the image of the turning point.	B1
			(3)
			(8 marks)

Question Number	Scheme	Marks	
	Mark (a) and (b) together		
10(a)	$(1+ax)^{20} = 1^{20} + {}^{20}C_1 1^{19} (ax)^1 + {}^{20}C_2 1^{18} (ax)^2$. Note that the notation $\binom{20}{1}$ may be seen for ${}^{20}C_1$ etc.		
	${}^{20}C_1 1^{19} (ax)^1 = 4x \Rightarrow 20a = 4 \Rightarrow a = 0.2$	M1: Uses either ${}^{20}C_1 (1^{19})(ax)^1 = 4x^1$ or $20a = 4$ to obtain a value for a . A1: $a = 0.2$ or equivalent	M1A1
			(2)
(b)	${}^{20}C_2 1^{18} (ax)^2 = px^2$ $\Rightarrow \frac{20 \times 19}{2} \times (0.2)^2 = p$ $\Rightarrow p = \dots$	Uses ${}^{20}C_2 (1^{18})(ax)^2 = px^2$ and their value of a to find a value for p . Condone the use of a rather than a^2 in finding p . Maybe implied by an attempt to find a value for $190a^2$ or $190a$. Note: ${}^{20}C_{18}$ can be used for ${}^{20}C_2$	M1
	$p = 7.6$	Accept equivalents such as $\frac{38}{5}, \frac{190}{25}$	A1
			(2)
(c)	Term is ${}^{20}C_4 1^{16} (ax)^4 \Rightarrow q = \dots$	Identifies the correct term and uses their value of a to find a value for q . Condone the use of a rather than a^4 . Must be an attempt to calculate ${}^{20}C_4 a^4$ or ${}^{20}C_4 a$ or ${}^{20}C_{16} a^4$ or ${}^{20}C_{16} a$	M1
	$q = {}^{20}C_4 \times 0.2^4 = \frac{969}{125} = (7.752)$	$q = \frac{969}{125}$ or exact equivalent e.g. $7.752, 7\frac{94}{125}$. $q = \frac{969}{125} x^4$ scores A0 but $qx^4 = \frac{969}{125} x^4$ scores A1.	A1
			(2)
			(6 marks)

Question Number	Scheme		Marks
11(i)	$3\cos^2 x + 1 = 4(1 - \cos^2 x)$ <p style="text-align: center;">or</p> $3(1 - \sin^2 x) + 1 = 4\sin^2 x$ <p style="text-align: center;">or</p> $3 + \tan^2 x + 1 = 4\tan^2 x$ <p style="text-align: center;">or</p> $3\frac{\cos 2x + 1}{2} + 1 = 4\frac{1 - \cos 2x}{2}$	Uses $\sin^2 x = 1 - \cos^2 x$ to produce an equation in $\cos^2 x$ or uses $\cos^2 x = 1 - \sin^2 x$ to produce an equation in $\sin^2 x$ or uses $\cos^2 x + \sin^2 x = 1$ and divides by $\cos^2 x$ to produce an equation in $\tan^2 x$ or uses $\sin^2 x$ and $\cos^2 x$ in terms of $\cos 2x$. Condone missing brackets.	M1
	$\Rightarrow \cos^2 x = \frac{3}{7} \text{ or } \sin^2 x = \frac{4}{7} \text{ or}$ $\tan^2 x = \frac{4}{3} \text{ or } \cos 2x = -\frac{1}{7}$	Correct value for $\cos^2 x$ or $\sin^2 x$ or $\tan^2 x$ or $\cos 2x$. This may be implied by $\cos x = \sqrt{\frac{3}{7}}$ or $\sin x = \sqrt{\frac{4}{7}}$ or $\tan x = \sqrt{\frac{4}{3}}$	A1
	$\Rightarrow \cos x = \pm\sqrt{\frac{3}{7}} \Rightarrow x = \cos^{-1}\left(\sqrt{\frac{3}{7}}\right)$ <p>A correct order of operations to obtain a correct expression for x. E.g.</p> $\cos^2 x = p \Rightarrow \cos x = \sqrt{p} \Rightarrow x = \cos^{-1}\sqrt{p} \text{ or}$ $\sin^2 x = p \Rightarrow \sin x = \sqrt{p} \Rightarrow x = \sin^{-1}\sqrt{p} \text{ or}$ $\tan^2 x = p \Rightarrow \tan x = \sqrt{p} \Rightarrow x = \tan^{-1}\sqrt{p} \text{ or}$ $\cos 2x = p \Rightarrow 2x = \cos^{-1} p \Rightarrow x = \frac{1}{2}\cos^{-1} p$ <p>This may be implied by one correct answer for their values.</p>		M1
	$\Rightarrow x = \text{awrt } 0.86, 2.28, 4.00, 5.43$	A1: Any two of awrt 0.86, 2.28, 4.00, 5.43	A2,1,0
		A1: All four of awrt 0.86, 2.28, 4.00, 5.43 with no additional solutions in the range and ignore solutions outside the range.	
<p>Note that answers in degrees are: 49.11, 130.89, 229.11, 310.89</p> <p>Allow A1 for awrt two of these but deduct the final A mark.</p> <p>For answers given as awrt $0.27\pi, 0.73\pi, 1.27\pi, 1.73\pi$, allow A1 only for any 2 of these but deduct the final A mark.</p>			
		(5)	

(ii)	$5 \sin(\theta + 10^\circ) = \cos(\theta + 10^\circ)$ $\Rightarrow \tan(\theta + 10^\circ) = 0.2$	M1: Reaches $\tan(\dots) = \alpha$ where α is a constant including zero. A1: $\tan(\dots) = 0.2$	M1A1
	$\Rightarrow \theta = \tan^{-1}(0.2) - 10^\circ$	For the correct order of operations to produce one value for θ . Accept $\theta = \tan^{-1}(\alpha) - 10$, $\alpha \neq 0$ or one correct answer as evidence. Dependent on the first M.	dM1
	$\Rightarrow \theta = \text{awrt } 1.3^\circ, 181.3^\circ$	A1: One of awrt $\theta = 1.3, 181.3$	A1A1
		A1: Both awrt $\theta = 1.3, 181.3$ and no other solutions in range and ignore solutions outside the range.	
	Note that final answers in radians in (ii) cannot score the final 2 A marks but the earlier marks are available (maximum 11100)		
			(10 marks)
Alternative 1 for (ii) by squaring:			
	$5 \sin(\dots) = \cos(\dots)$ $\Rightarrow 25 \sin^2(\dots) = \cos^2(\dots)$ $\Rightarrow 25(1 - \cos^2(\dots)) = \cos^2(\dots)$ or $25 \sin^2(\dots) = 1 - \sin^2(\dots)$ Leading to $\sin^2(\dots) = \dots$ or $\cos^2(\dots) = \dots$	Squares both sides, replaces $\sin^2(\dots)$ by $1 - \cos^2(\dots)$ or replaces $\cos^2(\dots)$ by $1 - \sin^2(\dots)$ and reaches $\sin^2(\dots) = \dots$ or $\cos^2(\dots) = \dots$	M1
	$\sin^2(\dots) = \frac{1}{26}$ or $\cos^2(\dots) = \frac{25}{26}$	Correct value for $\sin^2(\dots)$ or $\cos^2(\dots)$. This may be implied by $\sin(\dots) = \frac{1}{\sqrt{26}}$ or $\cos(\dots) = \sqrt{\frac{25}{26}}$	A1
	$\theta = \sin^{-1} \frac{1}{\sqrt{26}} - 10^\circ$ or $\theta = \cos^{-1} \frac{5}{\sqrt{26}} - 10^\circ$	For the correct order of operations to produce one value for θ as shown or accept one correct answer as evidence. Dependent on the first M.	dM1
	$\Rightarrow \theta = 1.3^\circ, 181.3^\circ$	A1: One of awrt $\theta = 1.3, 181.3$	A1A1
		A1: Both awrt $\theta = 1.3, 181.3$ and no other solutions in range and ignore solutions outside the range.	
Note that final answers in radians in (ii) cannot score the final 2 A marks but the earlier marks are available (maximum 11100)			

Alternative 2 for (ii) Using the addition formulae		
Alt (ii)	$5 \sin \theta \cos 10 + 5 \cos \theta \sin 10 = \cos \theta \cos 10 - \sin \theta \sin 10$ Uses the correct addition formulae on both sides and rearranges to $\tan(\dots) =$	M1
	$\tan \theta = \frac{\cos 10 - 5 \sin 10}{5 \cos 10 + \sin 10} = (0.0229)$	Correct value for $\tan \theta$ A1
	$\tan \theta = 0.0229 \Rightarrow \theta = \dots$	Uses arctan to produce one value for θ . Dependent on the first M. dM1
	$\Rightarrow \theta = 1.3^\circ, 181.3^\circ$	A1: One of awrt $\theta = 1.3, 181.3$ A1: Both awrt $\theta = 1.3, 181.3$ and no other solutions in range and ignore solutions outside the range. A1A1
	Note that final answers in radians in (ii) cannot score the final 2 A marks but the earlier marks are available (maximum 11100)	
		(5)


Question Number	Scheme		Marks
12(a)	$y = \frac{3}{4}x^2 - 4\sqrt{x} + 7 \Rightarrow \frac{dy}{dx} = \frac{3}{2}x - 2x^{-0.5}$	M1: Differentiates to obtain at least one correct power for one of the terms in x . (may be un-simplified) e.g. $x^2 \rightarrow x^{2-1}$ or $\sqrt{x} \rightarrow x^{\frac{1}{2}-1}$	M1A1
		A1: Correct derivative. Allow un-simplified e.g. $2 \times \frac{3}{4}x^{2-1}$ or $-4 \times \frac{1}{2}x^{\frac{1}{2}-1}$	
	At $x = 4$ $\frac{dy}{dx} = \frac{3}{2}(4) - 2(4)^{-0.5} = \dots$	Substitutes $x = 4$ into a changed function in an attempt to find the gradient.	M1
	$y - 11 = "5"(x - 4)$ or $y = mx + c \Rightarrow 11 = "5" \times 4 + c \Rightarrow c = \dots$	Correct straight line method using (4, 11) correctly placed and their dy/dx at $x = 4$ for the tangent not the normal . If using $y = mx + c$, must reach as far as finding a value for c . Dependent on the previous M.	dM1
	$y = 5x - 9$	Correct printed equation with no errors seen. Beware of the "5" appearing from wrong working.	A1*
<u>Important Note:</u> Following a correct derivative, if candidate states $x = 4$ so $dy/dx = 5$, this is fine if they then complete correctly – allow full marks. However, following a correct derivative, if the candidate <u>just</u> states $dy/dx = 5$ and then proceeds to obtain the correct straight line equation, the final mark can be withheld. Some evidence is needed that the candidate is considering the gradient at $x = 4$.			
			(5)





For part (b), in all cases, look to apply the appropriate scheme that gives the candidate the best mark

	Finds area under curve between 1 and 4 and subtracts triangle C (see diagram at end)		
(b) Way 1	$\int \frac{3}{4}x^2 - 4\sqrt{x} + 7 \, dx = \frac{1}{4}x^3 - \frac{8}{3}x^{1.5} + 7x (+c)$	M1: $x^n \rightarrow x^{n+1}$ on any term. May be un-simplified e.g. $x^2 \rightarrow x^{2+1}$, $x^{0.5} \rightarrow x^{0.5+1}$, $7 \rightarrow 7x^1$	M1A1
		A1: Correct integration. May be un-simplified e.g. terms such as $\frac{1}{3} \times \frac{3}{4}x^{2+1}$, $-\frac{2}{3} \times 4x^{0.5+1}$, $7x^1$ and $+c$ is not required.	
	Tangent meets x axis at $x = 1.8$	This may be embedded within a triangle area below or may be seen on a diagram.	B1
	<p style="text-align: center;">Area of triangle = $\frac{1}{2} \times (4 - '1.8') \times 11 = (12.1)$</p> <p style="text-align: center;">Correct method for the area of a triangle - look for $\frac{1}{2} \times (4 - '1.8') \times 11$</p> <p style="text-align: center;">This may be implied by the evaluation of $\int_{'1.8'}^4 5x - 9 \, dx = \left[5\frac{x^2}{2} - 9x \right]_{'1.8'}^4$</p>		M1
	<p style="text-align: center;">Correct method for area = Area A + Area B + Area C – Area C</p> $\left(\frac{1}{4}4^3 - \frac{8}{3} \times 4^{1.5} + 7 \times 4 \right) - \left(\frac{1}{4}1^3 - \frac{8}{3} \times 1^{1.5} + 7 \times 1 \right) - '12.1'$ <p style="text-align: center;">Correct combination of areas. Dependent on both previous method marks.</p>		ddM1
	= awrt 5.98	Area of R = awrt 5.98 or allow the exact answer of $\frac{359}{60}$ or equivalent.	A1
		(6)	
		(11 marks)	

	Finds area under curve between 1 and "1.8" and adds "line – curve" or "curve – line" between "1.8" and 4		
(b) Way 2	$\int \frac{3}{4}x^2 - 4\sqrt{x} + 7 \, dx = \frac{1}{4}x^3 - \frac{8}{3}x^{1.5} + 7x (+c)$	M1: $x^n \rightarrow x^{n+1}$ on any term. May be un-simplified e.g. $x^2 \rightarrow x^{2+1}$, $x^{0.5} \rightarrow x^{0.5+1}$, $7 \rightarrow 7x^1$	M1A1
		A1: Correct integration. May be un-simplified e.g. terms such as $\frac{1}{3} \times \frac{3}{4}x^{2+1}$, $-\frac{2}{3} \times 4x^{0.5+1}$, $7x^1$ and $+c$ is not required.	
	Tangent meets x axis at $x = 1.8$	This may be seen on a diagram.	B1
	<p style="text-align: center;">Area between "1.8" and 4 =</p> $\pm \int_{1.8}^4 \left(\frac{3}{4}x^2 - 4\sqrt{x} + 7 \right) - (5x - 9) \, dx = \pm \left[\frac{1}{4}x^3 - \frac{8}{3}x^{1.5} - \frac{5x^2}{2} + 16x \right]_{1.8}^4$ $= \frac{56}{3} - 15.7182... (= 2.9485...)$ <p style="text-align: center;">Attempts to integrate "curve – line" or "line – curve", substitute the limits "1.8" and 4 and subtracts.</p>		M1
	<p style="text-align: center;">Correct method for area = Area A + Area B</p> $\left(\left(\frac{1}{4} "1.8^3 - \frac{8}{3} "1.8^{1.5} + 7 \times "1.8 \right) - \left(\frac{1}{4} 1^3 - \frac{8}{3} 1^{1.5} + 7 \times 1 \right) + '2.9485...' \right)$ <p style="text-align: center;">Correct combination of areas. Dependent on both previous method marks.</p>		ddM1
	= awrt 5.98	Area of R = awrt 5.98 or allow the exact answer of $\frac{359}{60}$ or equivalent.	A1
			(6)

	Uses “line – curve” or “curve – line” between 1 and 4 and subtracts triangle below x axis		
(b) Way 3	$\pm \left(\frac{3}{4}x^2 - 4\sqrt{x} + 7 - 5x + 9 \right) = \pm \left(\frac{3}{4}x^2 - 4\sqrt{x} - 5x + 16 \right)$ $\pm \int \frac{3}{4}x^2 - 4\sqrt{x} - 5x + 16 \, dx = \pm \left(\frac{1}{4}x^3 - \frac{8}{3}x^{1.5} - \frac{5x^2}{2} \right) + kx(+c)$	M1A1	
	M1: $x^n \rightarrow x^{n+1}$ on any term. May be un-simplified e.g. $x^2 \rightarrow x^{2+1}$, $x^{0.5} \rightarrow x^{0.5+1}$, $x \rightarrow x^{1+1}$, $16 \rightarrow 16x^1$. If terms are not collected when subtracting then the same condition applies. A1: Correct integration as shown. May be un-simplified for coefficients and powers and $+c$ is not required.		
	Tangent meets x axis at $x = 1.8$	This may be embedded within a triangle area below or may be seen on a diagram.	B1
	Area of triangle = $\frac{1}{2} \times (1.8 - 1) \times 5 \times 1 - 9 = (1.6)$ Correct method for the area of a triangle - look for $\frac{1}{2} \times (1.8 - 1) \times 5 \times 1 - 9 $		M1
	Correct method for area = Area A + Area B + Area D – Area D $\left(\left(\frac{1}{4}4^3 - \frac{8}{3}4^{1.5} - \frac{5 \times 4^2}{2} + 16 \times 4 \right) - \left(\frac{1}{4}1^3 - \frac{8}{3}1^{1.5} - \frac{5 \times 1^2}{2} + 16 \times 1 \right) - '1.6' \right)$		ddM1
= awrt 5.98	Area of R = awrt 5.98 or allow the exact answer of $\frac{359}{60}$ or equivalent.	A1	
		(6)	



Question Number	Scheme	Marks
13(a)(i)		
	 <p>The maximum must be smooth and not form a point and the branches must not clearly turn back in on themselves.</p> <p style="text-align: center;">or</p> <p>A continuous graph passing through or touching at the points $(-c, 0)$, $(c, 0)$ and $(0, c^2)$. They can appear on their sketch or within the body of the script but there must be a sketch. Allow these marked as $-c$, c and c^2 in the correct places. Allow $(0, -c)$, $(0, c)$ and $(c^2, 0)$ as long as they are marked in the correct places. If there is any ambiguity, the sketch takes precedence.</p>	B1
	A fully correct diagram with the curve in the correct position and the intercepts and shape as described above. The maximum must be on the y-axis and the branches must extend below the x-axis.	B1
(a)(ii)	There must be a sketch to score any marks in (a)	
	 <p> Shape. A positive cubic with only one maximum and one minimum. The curve must be smooth at the maximum and at the minimum (not pointed).</p> <p>A smooth curve that touches or meets the x-axis at the origin and $(3c, 0)$ in the correct place and no other intersections. The origin does not need to be marked but the $(3c, 0)$ does. Allow $3c$ or $(0, 3c)$ to be marked in the correct place. May appear on their sketch or within the body of the script. If there is any ambiguity, the sketch takes precedence.</p> <p>Maximum at the origin (allow the maximum to form a point or cusp)</p>	B1 B1 B1
	There must be a sketch to score any marks in (a)	(5)
(b)	Intersect when $x^2(x-3c) = c^2 - x^2 \Rightarrow x^3 - 3cx^2 = c^2 - x^2$ Sets equations equal to each other and attempts to multiply out the bracket or vice versa	M1
	$x^3 + x^2 - 3cx^2 - c^2 = 0$ $\Rightarrow x^3 + (1-3c)x^2 - c^2 = 0^*$	Collects to one side (may be implied), factorises the x^2 terms and obtains printed answer with no errors. There must be an intermediate line of working. Allow $x^3 + x^2(1-3c) - c^2 = 0$ or $0 = x^3 + (1-3c)x^2 - c^2$ or $0 = x^3 + x^2(1-3c) - c^2$
		(2)

(c)	$8 + 4(1 - 3c) - c^2 = 0$	Substitutes $x = 2$ to give a correct un-simplified form of the equation.	M1
	$c^2 + 12c - 12 = 0$	Correct 3 term quadratic. Allow any equivalent form with the terms collected (may be implied)	A1
	$(c + 6)^2 - 36 - 12 = 0 \Rightarrow c = \dots$ or $c = \frac{-12 \pm \sqrt{12^2 - 4 \times 1 \times (-12)}}{2}$	Solves their 3TQ by using the formula or completing the square only . This may be implied by a correct exact answer for their 3TQ. (May need to check)	M1
	$4\sqrt{3} - 6$	$c = 4\sqrt{3} - 6$ or $c = -6 + 4\sqrt{3}$ only	A1
			(4)
			(11 marks)

Question Number	Scheme		Marks
14 (a)	Allow the use of S or S_n throughout without penalty. $S = a + ar + ar^2 + \dots + ar^{n-1}$ and $rS = ar + ar^2 + ar^3 + \dots + ar^n$ There must be a minimum of '3' terms and must include the first and the n th term. Condone for this mark only $S = a + ar + ar^2 + \dots + ar^n$ and $rS = ar + ar^2 + ar^3 + \dots + ar^{n+1}$ and allow commas instead of '+'s but see note below.		M1
	$S - rS = a - ar^n$	Subtracts either way around. As a special case allow $S - rS = a + ar^n$. For this mark, their S and their rS must be different but it must be S and rS they are considering with possible missing terms or slips.	M1
	$\Rightarrow S(1-r) = a(1-r^n) \Rightarrow S = \frac{a(1-r^n)}{(1-r)}$	dM1: Dependent upon both previous M's. It is for taking out a common factor of S and achieving $S = \dots$	dM1A1*
		A1*: Fully correct proof with no errors or omissions. The use of commas instead of '+'s is an error. $S = \frac{a(r^n - 1)}{(r - 1)}$ without reaching the printed answer is A0	
(4)			
(a) Way 2	$S = \frac{(a + ar + ar^2 + \dots + ar^{n-1})(1-r)}{1-r}$	Gives a minimum of '3' terms and must include the first and the n th and multiplies top and bottom by $1-r$	M1
	$S = \frac{a + ar + ar^2 + \dots + ar^{n-1} - ar - ar^2 - \dots - ar^n}{1-r}$	Expands the top with a minimum of '3' terms in each and must include the first and the n th term	M1
	$S = \frac{a(1-r^n)}{(1-r)}$	dM1: Dependent upon both previous M's. It is for taking out a common factor of a on top and achieving $S = \dots$	dM1A1
		A1*: Fully correct proof with no errors or omissions. The use of commas instead of '+'s is an error. $S = \frac{a(r^n - 1)}{(r - 1)}$ without reaching the printed answer is A0	

(b)	$U = 180 \times 0.93^n$ with $n = 4$ or 5	Attempts $U = 180 \times 0.93^n$ with $n = 4$ or 5 . Accept $U = 167.4 \times 0.93^n$ with $n = 3$ or 4 Allow 93% for 0.93	M1
	$U_5 = 180 \times (0.93)^5 = 125.2$ (litres)	Cso. Awrt 125.2	A1*
	Allow 93% or 1 – 7% for 0.93		
(c)	Attempts $S_n = \frac{a(1-r^n)}{(1-r)}$ with any combination of: $n = 20/21$ $a = 180/167.4$ and $r = 0.93$ Allow 93% for 0.93		M1
	$S = \frac{167.4(1-0.93^{20})}{(1-0.93)}$ or $S = 180 \times \frac{0.93(1-0.93^{20})}{(1-0.93)}$ or $S = \frac{180(1-0.93^{21})}{(1-0.93)} - 180$		A1
	A correct numerical expression for the sum (may be implied by awrt 1831) Allow 93% or 1 – 7% for 0.93		
	1831 (litres)	1831 only (Ignore units). Do not isw here, so 1831 followed by $1831 \times 20 = \dots$ scores A0.	A1
			(3)
			(9 marks)

Listing:

(b)	Sight of awrt 180, 167, 156, 145, 135, 125	Starts with 180 and multiplies by 0.93 either 4 or 5 times showing each result at least to the nearest litre and chooses the 5 th or 6 th term	M1
	$U_5 = 125.2$ (litres)	Must see all values accurate to 1dp: e.g. awrt 180, 167.4, 155.7, 144.8, (134.6 or 134.7), 125.2	A1*
			(2)
(c)	Total = $180 \times 0.93 + 180 \times 0.93^2 + \dots + 180 \times 0.93^{19} + 180 \times 0.93^{20} = \dots$ Finds an expression for the sum of 20 or 21 terms		M1
	All sums accurate to awrt 1dp 167.4+155.7+144.8+134.6+125.2+.....42.2 A correct numerical expression for the sum (may be implied by awrt 1831)		A1
	1831 (litres)	1831 only (Ignore units). Do not isw here, so 1831 followed by $1831 \times 20 = \dots$ scores A0.	A1
			(3)

Question Number	Scheme	Marks	
15	Area of triangle = $\frac{1}{2} \times (2r)^2 \sin\left(\frac{\pi}{3} \text{ or } 60\right)$ or $\frac{1}{2} \times (r)^2 \sin\left(\frac{\pi}{3} \text{ or } 60\right)$ Correct method for the area of either triangle. Ignore any reference to which triangle they are finding the area of.	M1	
	Area of sector = $\frac{1}{2} \times r^2 \times \frac{\pi}{3}$	Use of the sector formula $\frac{1}{2} r^2 \theta$ with $\theta = \frac{\pi}{3}$ which may be embedded within a segment	M1
	$\text{Area } R = \text{Sector} + 2 \text{ Segments} = \frac{1}{2} r^2 \times \frac{\pi}{3} + 2 \times \left(\frac{1}{2} r^2 \times \frac{\pi}{3} - \frac{1}{2} r^2 \times \frac{\sqrt{3}}{2} \right)$ $\text{Area } R = \text{Triangle} + 3 \text{ Segments} = \frac{1}{2} r^2 \times \frac{\sqrt{3}}{2} + 3 \times \left(\frac{1}{2} r^2 \times \frac{\pi}{3} - \frac{1}{2} r^2 \times \frac{\sqrt{3}}{2} \right)$ $\text{Area } R = 3 \text{ Sectors} - 2 \text{ Triangles} = 3 \times \frac{1}{2} r^2 \times \frac{\pi}{3} - 2 \times \left(\frac{1}{2} r^2 \times \frac{\sqrt{3}}{2} \right)$ $\text{Area } R = \text{Big triangle} - 3 \text{ White bits}$ $= \frac{1}{2} \times (2r)^2 \frac{\sqrt{3}}{2} - 3 \times \left(\frac{1}{2} r^2 \times \frac{\sqrt{3}}{2} - \left(\frac{1}{2} r^2 \times \frac{\pi}{3} - \frac{1}{2} r^2 \times \frac{\sqrt{3}}{2} \right) \right)$ <p>M1: A fully correct method (may be implied by a final answer of awrt $0.705r^2$)</p> <p>A1: Correct exact expression - for this to be scored $\sin \frac{\pi}{3} = \frac{\sqrt{3}}{2}$ must be seen</p>		M1A1
	$= \frac{1}{2} \pi r^2 - \frac{\sqrt{3}}{2} r^2 = r^2 \left(\frac{1}{2} \pi - \frac{\sqrt{3}}{2} \right)$	Cso (Allow $\frac{r^2}{2} (\pi - \sqrt{3})$ or any exact equivalent with r^2 taken out as a common factor)	A1
			(5 marks)

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom