Surname	Other nar	nes
Pearson Edexcel International Advanced Level	Centre Number	Candidate Number
Core Math		s C12
Tuesday 9 January 2018 – N Time: 2 hours 30 minutes	•	Paper Reference WMA01/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 125.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1. Given that

$$y = \frac{2x^{\frac{2}{3}} + 3}{6}, \quad x > 0$$

find, in the simplest form,

(a) $\frac{\mathrm{d}y}{\mathrm{d}x}$

(2)

(b) $\int y \, dx$

(3)

P 5 0 7 1 3 A 0 2 5 2

		Leave blank
Question 1 continued		Junk
		Q1
	(Total 5 marks)	
	. ,	



2. A sequence is defined by

$$u_1 = 1$$

 $u_{n+1} = 2 - 3u_n \quad n \geqslant 1$

(a) Find the value of u_2 and the value of u_3

(2)

(b) Calculate the value of $\sum_{r=1}^{4} (r - u_r)$

(3)

P 5 0 7 1 3 A 0 4 5 2

Question 2 continued		Leave
		Q2
	(Total 5 marks)	
	(10tal 3 mal Ks)	



3. Simplify fully



(2)

(b)
$$\frac{2y^7 \times (4y)^{-2}}{3y}$$

(2)

	_

	Leave blank
Question 3 continued	
	Q3
(Total 4 marks)	
	$\overline{}$



4. The equation

$$(p-2)x^2 + 8x + (p+4) = 0$$
, where p is a constant

has no real roots.

(a) Show that p satisfies $p^2 + 2p - 24 > 0$

(3)

(b) Hence find the set of possible values of p.

(4)

Question 4 continued		Lea bla
		Q4
	(Total 7 marks)	



- **5.** (In this question, solutions based entirely on graphical or numerical methods are not acceptable.)
 - (i) Solve, for $0 < \theta < \frac{\pi}{2}$

$$5\sin 3\theta - 7\cos 3\theta = 0$$

Give each solution, in radians, to 3 significant figures.

(5)

(ii) Solve, for $0 < x < 360^{\circ}$

$$9\cos^2 x + 5\cos x = 3\sin^2 x$$

Give each solution, in degrees, to one decimal place.

(6)

	Leave
	blank
Question 5 continued	

Question 5 continued	

Question 5 continued	Leave
Question 5 continued	
	Q5
(Total 11 marks)	



6.

$$f(x) = ax^3 - 8x^2 + bx + 6$$

where a and b are constants.

When f(x) is divided by (x + 1) there is no remainder.

When f(x) is divided by (x - 2) the remainder is -12

(a) Find the value of a and the value of b.

(5)

(b) Factorise f(x) completely.

(4)

	Т,
	Leave
	blank
Question 6 continued	
Question o continued	



nestion 6 continued	

	Leave
Question 6 continued	blank
Question 6 continued	
	Q6
(Total 9 marks)	



7.




Figure 2

Figure 1 shows a rectangular sheet of metal of negligible thickness, which measures 25 cm by 15 cm. Squares of side x cm are cut from each corner of the sheet and the remainder is folded along the dotted lines to make an open cuboid box, as shown in Figure 2.

(a) Show that the volume, $V \text{cm}^3$, of the box is given by

$$V = 4x^3 - 80x^2 + 375x \tag{2}$$

(b) Use calculus to find the value of x, to 3 significant figures, for which the volume of the box is a maximum.

(4)

(c) Justify that this value of x gives a maximum value for V.

(2)

(d) Find, to 3 significant figures, the maximum volume of the box.

(2)

	Leave
	blank
Question 7 continued	



estion 7 continued	

	Leave blank
Question 7 continued	
	Q7
(Total 10 marks)	
(



8.




Figure 3

Figure 3 shows a sketch of the curve with equation $y = f(x), x \in \mathbb{R}$.

The curve crosses the y-axis at the point (0, 5) and crosses the x-axis at the point (6, 0).

The curve has a minimum point at (1, 3) and a maximum point at (4, 7).

On separate diagrams, sketch the curve with equation

(a)
$$y = f(-x)$$
 (3)

(b)
$$y = f(2x)$$
 (3)

On each diagram, show clearly the coordinates of any points of intersection of the curve with the two coordinate axes and the coordinates of the stationary points.

Question 8 continued	blank
	23

Question 8 continued

Leave blank

blank **Question 8 continued** Q8 (Total 6 marks) 25



Leave

9.	The first term of a geometric series is 20 and the common ratio is 0.9	
	(a) Find the value of the fifth towns	
	(a) Find the value of the fifth term.	(2)
		()
	(b) Find the sum of the first 8 terms, giving your answer to one decimal place.	(2)
		(2)
	Given that $S_{\infty} - S_N < 0.04$, where S_N is the sum of the first N terms of this series,	
	(c) show that $0.9^N < 0.0002$	
		(4)
	(d) Hence find the smallest possible value of <i>N</i> .	
	(a) Thence find the smallest possible value of 7.	(2)

	Leave
	blank
Question 9 continued	



	Leave blank
Question 9 continued	

Question 9 continued		Leave blank
Question > continued		
		Q9
		Ť
(Total 10 m	arks)	



10. (i) Use the laws of logarithms to solve the equation

$$3\log_8 2 + \log_8 (7 - x) = 2 + \log_8 x$$

(5)

Leave blank

(ii) Using algebra, find, in terms of logarithms, the exact value of y for which

$$3^{2y} + 3^{y+1} = 10$$

(5)

	Leave
	blank
Question 10 continued	



Question 10 continued	

	Leave
Question 10 continued	blank
Question to continued	
	Q10
	VIO
(Total 10 marks)	
	\rightarrow



11. The circle C has equation

$$x^2 + y^2 - 8x - 10y + 16 = 0$$

The centre of C is at the point T.

- (a) Find
 - (i) the coordinates of the point T,
 - (ii) the radius of the circle *C*.

(4)

The point M has coordinates (20, 12).

(b) Find the exact length of the line MT.

(2)

Point P lies on the circle C such that the tangent at P passes through the point M.

- (c) Find the exact area of triangle MTP, giving your answer as a simplified surd.
- (3)

34

estion 11 continued	



Question 11 continued	

	Leave blank
Question 11 continued	
	Q11
(Total 9 marks)	



12. The line l_1 has equation x + 3y - 11 = 0

The point A and the point B lie on l_1

Given that A has coordinates (-1, p) and B has coordinates (q, 2), where p and q are integers,

(a) find the value of p and the value of q,

(2)

(b) find the length of AB, giving your answer as a simplified surd.

(2)

The line l_2 is perpendicular to l_1 and passes through the midpoint of AB.

(c) Find an equation for l_2 giving your answer in the form y = mx + c, where m and c are constants to be found.

(5)

	Lea
	bla
Question 12 continued	l Ola
Question 12 continued	



Leave blank

Question 12 continued

40

Question 12 continued	blank
	Q12
(Total 9 marks)	



DO NOT WRITE IN THIS AREA




Figure 4 shows the position of two stationary boats, A and B, and a port P which are assumed to be in the same horizontal plane.

Boat A is $8.7 \,\mathrm{km}$ on a bearing of 314° from port P.

Boat *B* is $3.5 \,\mathrm{km}$ on a bearing of 052° from port *P*.

(a) Show that angle APB is 98°

(1)

- (b) Find the distance of boat B from boat A, giving your answer to one decimal place. (2)
- (c) Find the bearing of boat B from boat A, giving your answer to the nearest degree. (4)

	Leave
	blank
Question 13 continued	



Leave blank

Question 13 continued		

Question 13 continued		Leave
Q13		blank
Q13	Question 13 continued	
		Q13
(Total 7 marks)		
(Total / marks)	(Total 7 marks)	
	(Total / marks)	






Figure 5 shows a sketch of part of the line l with equation y = 8 - x and part of the curve C with equation $y = 14 + 3x - 2x^2$

The line l and the curve C intersect at the point A and the point B as shown.

(a) Use algebra to find the coordinates of A and the coordinates of B.

(5)

The region R, shown shaded in Figure 5, is bounded by the coordinate axes, the line l, and the curve C.

(b) Use algebraic integration to calculate the exact area of R.

(8)

	Leave
	blank
Question 14 continued	Chaine
Question 14 continued	



stion 14 continued	

https://xtremepape.rs/

	Leave
Question 14 continued	blank
Question 11 continued	
	014
	Q14
(Total 13 marks)	
(10th 15 marks)	



15. The binomial expansion, in ascending powers of x, of $(1 + kx)^n$ is

$$1 + 36x + 126kx^2 + \dots$$

where k is a non-zero constant and n is a positive integer.

(a) Show that nk(n - 1) = 252

(2)

(b) Find the value of k and the value of n.

- **(5)**
- (c) Using the values of k and n, find the coefficient of x^3 in the binomial expansion of $(1 + kx)^n$
- **(3)**

uestion 15 continued	



Question 15 continued		blank
		Q15
	(Total 10 marks)	
TO	OTAL FOR PAPER: 125 MARKS	
END		