



Pearson
Edexcel

Mark Scheme (Results)

June 2018

Pearson Edexcel
International Advanced Subsidiary Level
In Further Pure Mathematics F1 (WFM01)
Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

June 2018

Publications Code

All the material in this publication is copyright

© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

PEARSON EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
- ft – follow through
- the symbol \surd will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- o.e. – or equivalent (and appropriate)
- d... or dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper or ag- answer given
- \square or d... The second mark is dependent on gaining the first mark

4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Further Pure Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$, leading to $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$, where $|pq| = |c|$ and $|mn| = |a|$, leading to $x = \dots$

2. Formula

Attempt to use the correct formula (with values for a, b and c).

3. Completing the square

Solving $x^2 + bx + c = 0$: $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$, $q \neq 0$, leading to $x = \dots$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \rightarrow x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \rightarrow x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

June 2018
WFM01 Further Pure Mathematics F1
Mark Scheme

Question Number	Scheme	Notes	Marks
1.	$\sum_{r=1}^n r(r+3) = \sum_{r=1}^n r^2 + 3 \sum_{r=1}^n r$		
	$= \frac{1}{6}n(n+1)(2n+1) + 3\left(\frac{1}{2}n(n+1)\right)$	Attempts to expand $r(r+3)$ and attempts to substitute at least one correct standard formula into their resulting expression.	M1
		Correct expression (or equivalent)	A1
	$= \frac{1}{6}n(n+1)[(2n+1) + 9]$	dependent on the previous M mark Attempt to factorise at least $n(n+1)$ having attempted to substitute both correct standard formulae.	dM1
	$= \frac{1}{6}n(n+1)(2n+10)$	{this step does not have to be written}	
	$= \frac{n}{3}(n+1)(n+5) \text{ or } \frac{1}{3}n(n+1)(n+5)$	Correct completion with no errors. Note: $a=3, b=5$	A1
			(4)
			4
Question 1 Notes			
1.	Note	Applying e.g. $n=1, n=2$ to the printed equation without applying the standard formulae to give $a=3, b=5$ is M0A0M0A0	
	Alt 1	Alt Method 1 (Award the first two marks using the main scheme) Using $\frac{1}{3}n^3 + 2n^2 + \frac{5}{3}n \equiv \frac{1}{a}n^3 + \left(\frac{b+1}{a}\right)n^2 + \frac{b}{a}n$ o.e.	
	dM1	Equating coefficients to find both $a = \dots$ and $b = \dots$ and at least one correct of $a=3$ or $b=5$	
	A1	Finds $a=3$ and $b=5$	
	Alt 2	Alt Method 2: (Award the first two marks using the main scheme) $\frac{1}{6}n(n+1)(2n+1) + \frac{3}{2}n(n+1) \equiv \frac{n}{a}(n+1)(n+b)$	
	dM1	Substitutes $n=1, n=2$, into this identity o.e. and solves to find both $a = \dots$ and $b = \dots$ and at least one correct of $a=3, b=5$	
	A1	Note: $n=1$ gives $4 = \frac{2(1+b)}{a}$ or $2a-b=1$ and $n=2$ gives $14 = \frac{6(2+b)}{a}$ or $7a-3b=6$ Finds $a=3$ and $b=5$	
	Note	Allow final dM1A1 for $\frac{1}{3}n^3 + 2n^2 + \frac{5}{3}n$ or $\frac{1}{3}(n^3 + 6n^2 + 5n) \rightarrow \frac{n}{3}(n+1)(n+5)$ with no incorrect working.	
	Note	A correct proof $\sum_{r=1}^n r(r+3) = \frac{n}{3}(n+1)(n+5)$ followed by stating an incorrect e.g. $a=5, b=3$ is M1A1dM1A1 (ignore subsequent working)	
	Note	Give A0 for $\frac{2}{6}n(n+1)(n+5)$ without reference to $a=3$ or $\frac{n}{3}(n+1)(n+5)$ or $\frac{1}{3}n(n+1)(n+5)$	

Question Number	Scheme	Notes	Marks
2.	P represents an anti-clockwise rotation about the origin through 45 degrees		
(a)	$\{\mathbf{P} = \left\{ \begin{array}{cc} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{array} \right\}$ or $\left\{ \begin{array}{cc} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{array} \right\}$ or e.g. $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$	Correct matrix which is expressed in exact surds	B1
			(1)
(b)	Enlargement	Enlargement or enlarge	M1
	Centre (0, 0) with scale factor $k\sqrt{2}$	About (0, 0) or about <i>O</i> or about the origin and scale or factor or times and $k\sqrt{2}$ Note: Allow $\sqrt{2k^2}$ in place of $k\sqrt{2}$	A1
	Note: Give M0A0 for combinations of transformations		(2)
(c) Way 1	$\{\mathbf{PQ} = \left\{ \begin{array}{cc} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{array} \right\} \begin{pmatrix} k\sqrt{2} & 0 \\ 0 & k\sqrt{2} \end{pmatrix} = \begin{pmatrix} k & -k \\ k & k \end{pmatrix}$ $\{\det \mathbf{PQ} = \} (k)(k) - (-k)(k) = 2k^2$	Multiplies their matrix from part (a) by Q [either way round] and applies " <i>ad - bc</i> " to the resulting matrix to give $2k^2$ or states $ \text{their det } \mathbf{PQ} = 2k^2$ Condone $\{\det \mathbf{PQ} = \} k^2 + k^2$	M1
	$6(2k^2) = 147$ or $2k^2 = \frac{147}{6}$	6(their determinant) = 147 or puts their determinant equal to $\frac{147}{6}$	A1
	$\left\{ \Rightarrow k^2 = \frac{49}{4} \Rightarrow \right\} k = \frac{7}{2}$	Obtains $k = 3.5$, o.e.	A1
			(4)
(c) Way 2	$\det \mathbf{Q} = (k\sqrt{2})(k\sqrt{2}) - (0)(0)$ or $\det \mathbf{Q} = (k\sqrt{2})(k\sqrt{2})$	applies " <i>ad - bc</i> " to Q or applies $(k\sqrt{2})^2$	M1
	$\{\det \mathbf{P} = 1 \Rightarrow \} \det \mathbf{PQ} = (1)(2k^2) = 2k^2$ or $\det \mathbf{Q} = 2k^2$	and deduces that $\det \mathbf{PQ} = 2k^2$ or states $ \text{their det } \mathbf{PQ} = 2k^2$ or $\det \mathbf{Q} = 2k^2$	A1
	$6(2k^2) = 147$ or $2k^2 = \frac{147}{6}$	6(their det(PQ)) = 147 or (their det(PQ)) = $\frac{147}{6}$ or 6(their det(Q)) = 147 or (their det(PQ)) = $\frac{147}{6}$	M1
	$\left\{ \Rightarrow k^2 = \frac{49}{4} \Rightarrow \right\} k = \frac{7}{2}$	Obtains $k = 3.5$, o.e.	A1
			(4)

Question 2 Notes		
2. (b)	Note	“original point” is not acceptable in place of the word “origin”.
	Note	“expand” is not acceptable for M1
	Note	“enlarge x by $k\sqrt{2}$ and no change in y ” is M0A0
(c)	Note	Obtaining $k = \pm 3.5$ with no evidence of $k = 3.5$ {only} is A0
	Way 2 Note 1	Give M1A1M0A0 for writing down $147(2k^2) = 6$ or $\frac{1}{2k^2} = \frac{147}{6}$ or $6\left(\frac{1}{2k^2}\right) = 147$, o.e. with no other supporting working.
	Way 2 Note 2	Give M0A0M1A0 for writing $\det \mathbf{Q} = \frac{1}{k^2 - (-k^2)}$ or $\frac{1}{2k^2}$, followed by $6\left(\frac{1}{2k^2}\right) = 147$
	Note	Allow M1A1 for an incorrect rotation matrix \mathbf{P} , leading to $\det \mathbf{PQ} = 2k^2$
	Note	Allow M1A1M1A1 for an incorrect rotation matrix \mathbf{P} , leading to $\det \mathbf{PQ} = 2k^2$ and $k = 3.5$, o.e.
	Note	Using the scale factor of enlargement to write down $k\sqrt{2} = \sqrt{\frac{147}{6}} \Rightarrow k = 3.5$ is M1A1dM1A1
	Note	Using the scale factor of enlargement to write down $k\sqrt{2} = \sqrt{\frac{6}{147}}$ is M1A1dM0

Question Number	Scheme	Notes	Marks
3.	$C: y^2 = 6x$; S is the focus of C ; $y^2 = 4ax$; $P(at^2, 2at)$; Q lies on the directrix of C . $PQ = 14$		
(a)	$\{a = 1.5 \Rightarrow\}$ S has coordinates $(1.5, 0)$	$(1.5, 0)$ or $(\frac{3}{2}, 0)$ or $(\frac{6}{4}, 0)$	B1 cao
	Note: You can recover this mark for $S(1.5, 0)$ stated either parts (b) or part (c)		(1)
(b)	$\{PQ$ is parallel to the x -axis $\Rightarrow\}$ Focus-directrix Property $\Rightarrow SP \{= PQ\} = 14$	$SP = 14$ or 14 stated by itself in (b)	B1 cao
	Note: $PQ = 14$ stated by itself without reference to $SP = 14$ is B0		(1)
(c) Way 1	$\left\{ \text{directrix } x = -\frac{3}{2} \text{ \& } PQ = 14 \Rightarrow \right\} x_p = 14 - \frac{3}{2} \{= 12.5\}$	$x = 14 - \text{their "a"}$	M1
	$y_p^2 = 6(12.5) \Rightarrow y_p = \dots$	dependent on the previous M mark Substitutes their x into $y^2 = 6x$ and finds $y = \dots$	dM1
	Either $x = 12.5, y = 5\sqrt{3}$ or $(12.5, 5\sqrt{3})$	Correct and paired. Accept $(12.5, \sqrt{75})$	A1
			(3)
(c) Way 2	$(x - 1.5)^2 + (6x) = 14^2$ $\Rightarrow x^2 + 3x - 193.75 = 0 \Rightarrow x = \dots$	Applies Pythagoras to $x - "a"$, $\sqrt{6x}$ and 14, then forms and solves quadratic equation in x to give $x = \dots$	M1
		<i>As in Way 1</i>	dM1 A1
			(3)
(c) Way 3	$11^2 + y^2 = 14^2 \Rightarrow y = \dots$	Applies Pythagoras to $14 - "2a"$, y and 14, and solves to give $y = \dots$	M1
	$(\sqrt{75})^2 = 6x \Rightarrow x = \dots$	dependent on the previous M mark Substitutes their y into $y^2 = 6x$ and finds $x = \dots$	dM1
	Either $x = 12.5, y = 5\sqrt{3}$ or $(12.5, 5\sqrt{3})$	Correct and paired. Accept $(12.5, \sqrt{75})$	A1
			(3)
(c) Way 4	$(1.5t^2 - 1.5)^2 + (3t)^2 = 14^2$ $\Rightarrow 2.25t^4 + 4.5t^2 - 193.75 = 0$ {or $9t^4 + 18t^2 - 775 = 0$ } $\Rightarrow t^2 = \frac{25}{3} \Rightarrow t = \frac{5\sqrt{3}}{3}$ $\Rightarrow x = 1.5 \left(\frac{5\sqrt{3}}{3} \right)^2, y = 3 \left(\frac{5\sqrt{3}}{3} \right)$	Applies Pythagoras to $"1.5t^2 - 1.5"$, $2("1.5")t$ and 14, forms and solves a quadratic equation in t^2 to give $t^2 = \dots$ or $t = \dots$, and finds at least one of $x = \dots$ or $y = \dots$ by using $x = "1.5t^2"$ or $y = 2("1.5")t$	M1
		dependent on the previous M mark Finds both $x = \dots$ and $y = \dots$ by using $x = "1.5t^2"$ and $y = 2("1.5")t$	dM1
	Either $x = 12.5, y = 5\sqrt{3}$ or $(12.5, 5\sqrt{3})$	Correct and paired. Accept $(12.5, \sqrt{75})$	A1
			(3)
			5

Question Number	Scheme	Notes	Marks
3.	$C: y^2 = 6x; S$ is the focus of $C; y^2 = 4ax; P(at^2, 2at); Q$ lies on the directrix of $C. PQ = 14$		
(c) Way 5	$\left\{ x_p = \frac{3}{2}t^2, x_q = -\frac{3}{2}, PQ = 14 \Rightarrow \right\}$ $(1.5t^2 - -1.5) = 14 \Rightarrow 1.5t^2 = 12.5$ $\Rightarrow t^2 = \frac{25}{3} \Rightarrow t = \frac{5\sqrt{3}}{3}$ $\Rightarrow x = 1.5\left(\frac{5\sqrt{3}}{3}\right)^2, y = 2\left(\frac{5\sqrt{3}}{3}\right)t$	Uses horizontal distance $PQ = 14$ to form and solve the equation " $1.5t^2 - -1.5 = 14$ " to give $t^2 = \dots$ or $t = \dots$, and finds at least one of $x = \dots$ or $y = \dots$ by using $x = "1.5"t^2$ or $y = 2("1.5")t$	M1
		dependent on the previous M mark Finds both $x = \dots$ and $y = \dots$ by using $x = "1.5"t^2$ and $y = 2("1.5")t$	dM1
	Either $x = 12.5, y = 5\sqrt{3}$ or $(12.5, 5\sqrt{3})$	Correct and paired. Accept $(12.5, \sqrt{75})$	A1
			(3)
(c) Way 6	$\left\{ S(1.5, 0), P\left(\frac{y^2}{6}, y\right), SP = 14 \Rightarrow \right\}$ $\left(\frac{1}{6}y^2 - \frac{3}{2}\right)^2 + y^2 = 14^2 \Rightarrow y = \dots$ $\{y^4 + 18y^2 - 6975 = 0\}$	Applies Pythagoras to $\frac{y^2}{6} - "1.5", y$ and 14 , and solves to give $y = \dots$	M1
	$(\sqrt{75})^2 = 6x \Rightarrow x = \dots$	dependent on the previous M mark Substitutes their y into $y^2 = 6x$ and finds $x = \dots$	dM1
	Either $x = 12.5, y = 5\sqrt{3}$ or $(12.5, 5\sqrt{3})$	Correct and paired. Accept $(12.5, \sqrt{75})$	A1
Question 3 Notes			
3. (c)	Note	<u>Writing coordinates the wrong way round</u> E.g. writing $x = 12.5, y = 5\sqrt{3}$ followed by $(5\sqrt{3}, 12.5)$ is final A0	
	Note	Obtaining both $(12.5, 5\sqrt{3})$ and $(12.5, -5\sqrt{3})$ with no evidence of only $(12.5, 5\sqrt{3})$ is A0	
	Note	Give final A1 for $(12.5, \text{awrt } 8.66)$, with either $y = \sqrt{75}$ or $y = 5\sqrt{3}$ seen in their working	
	Note	You can mark part (b) and part (c) together	

Question Number	Scheme	Notes	Marks
4.	$\mathbf{A} = \begin{pmatrix} 2p & 3q \\ 3p & 5q \end{pmatrix}; \mathbf{XA} = \mathbf{B}; \mathbf{B} = \begin{pmatrix} p & q \\ 6p & 11q \\ 5p & 8q \end{pmatrix}$		
(a)	$\{\det(\mathbf{A}) =\} 2p(5q) - (3p)(3q) \{= pq\}$	$2p(5q) - (3p)(3q)$ which can be un-simplified or simplified	B1
	$\{\mathbf{A}^{-1} =\} \frac{1}{pq} \begin{pmatrix} 5q & -3q \\ -3p & 2p \end{pmatrix}$ or $\begin{pmatrix} \frac{5}{p} & -\frac{3}{p} \\ -\frac{3}{q} & \frac{2}{q} \end{pmatrix}$	$\begin{pmatrix} 5q & -3q \\ -3p & 2p \end{pmatrix}$	M1
		Correct \mathbf{A}^{-1}	A1
			(3)
(b) Way 1	$\{\mathbf{X} = \mathbf{BA}^{-1} =\}$ $\begin{pmatrix} p & q \\ 6p & 11q \\ 5p & 8q \end{pmatrix} \frac{1}{pq} \begin{pmatrix} 5q & -3q \\ -3p & 2p \end{pmatrix} = \dots$	Attempts \mathbf{BA}^{-1} and finds at least one element (or at least one element calculation) of their matrix \mathbf{X} Note: Allow one slip in copying down \mathbf{B} Note: Allow one slip in copying down \mathbf{A}^{-1}	M1
	$= \frac{1}{pq} \begin{pmatrix} 2pq & -pq \\ -3pq & 4pq \\ pq & pq \end{pmatrix}$	At least 4 correct elements (need not be in a matrix)	A1
		dependent on the first M mark Finds a 3×2 matrix of 6 elements	dM1
	$= \begin{pmatrix} 2 & -1 \\ -3 & 4 \\ 1 & 1 \end{pmatrix}$	Correct simplified matrix for \mathbf{X}	A1
			(4)
(b) Way 2	$\{\mathbf{XA} = \mathbf{B} \Rightarrow\} \begin{pmatrix} a & b \\ c & d \\ e & f \end{pmatrix} \begin{pmatrix} 2p & 3q \\ 3p & 5q \end{pmatrix} = \begin{pmatrix} p & q \\ 6p & 11q \\ 5p & 8q \end{pmatrix}$ $2pa + 3pb = p, \quad 3qa + 5qb = q$ or $2pc + 3pd = 6p, \quad 3qc + 5qd = 11q$ or $2pe + 3pf = 5p, \quad 3qe + 5qf = 8q$ and finds at least one of a, b, c, d, e or f	Applies $\mathbf{XA} = \mathbf{B}$ for a 3×2 matrix \mathbf{X} and attempts simultaneous equations in a and b or c and d or e and f to find at least one of a, b, c, d, e or f Note: Allow one slip in copying down \mathbf{A} Note: Allow one slip in copying down \mathbf{B}	M1
	$\left\{ \begin{array}{l} 2a + 3b = 1, \quad 3a + 5b = 1 \\ 2c + 3d = 6, \quad 3c + 5d = 11 \\ 2e + 3f = 5, \quad 3e + 5f = 8 \end{array} \right\} \Rightarrow \begin{array}{l} a = 2, b = -1 \\ c = -3, d = 4 \\ e = 1, f = 1 \end{array}$	At least 4 correct elements	A1
		dependent on the first M mark Finds all 6 elements for the 3×2 matrix \mathbf{X}	dM1
	$\Rightarrow \mathbf{X} = \begin{pmatrix} 2 & -1 \\ -3 & 4 \\ 1 & 1 \end{pmatrix}$	Correct simplified matrix for \mathbf{X}	A1
			(4)
			7

Question 4 Notes		
4. (a)	Note	Condone $\frac{1}{10pq-9pq} \begin{pmatrix} 5q & -3q \\ -3p & 2p \end{pmatrix}$ or $\frac{1}{2p(5q)-(3p)(3q)} \begin{pmatrix} 5q & -3q \\ -3p & 2p \end{pmatrix}$ for A1
	Note	Condone $\begin{pmatrix} 5q & -3q \\ -3p & 2p \end{pmatrix} \frac{1}{pq}$ or $\begin{pmatrix} 5q & -3q \\ -3p & 2p \end{pmatrix} \frac{1}{2p(5q)-(3p)(3q)}$ for A1
	Note	Condone $\begin{pmatrix} \frac{5q}{pq} & -\frac{3q}{pq} \\ -\frac{3p}{pq} & \frac{2p}{pq} \end{pmatrix}$ for A1
(b)	Note	<p>Way 1: Allow SC 1st A1 for at least 4 correct elements in</p> $\left(\begin{array}{cc} \frac{2pq}{\text{their det A}} & \frac{-pq}{\text{their det A}} \\ \frac{-3pq}{\text{their det A}} & \frac{4pq}{\text{their det A}} \\ \frac{pq}{\text{their det A}} & \frac{pq}{\text{their det A}} \end{array} \right)$ <p>or for at least 4 of these elements seen in their calculations</p>

Question Number	Scheme	Notes	Marks
5.	$z^4 - 6z^3 + 34z^2 - 54z + 225 \equiv (z^2 + 9)(z^2 + az + b)$; a, b are real numbers		
(a)	$a = -6, b = 25$	At least one of $a = -6$ or $b = 25$	B1
		Both $a = -6$ and $b = 25$	B1
			(2)
(b)	$\{z^2 + 9 = 0 \Rightarrow\} z = 3i, -3i$	At least one of $3i, -3i, \sqrt{9}i$ or $-\sqrt{9}i$	M1
		Both $3i$ and $-3i$	A1
	$\{z^2 - 6z + 25 = 0 \Rightarrow\}$ <ul style="list-style-type: none"> $z = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(1)(25)}}{2(1)}$ or $(z - 3)^2 - 9 + 25 = 0 \Rightarrow z = \dots$ 	Correct method of applying the quadratic formula or completing the square for solving their $z^2 + az + b = 0$; $a, b \neq 0$	M1
			(4)
(c)		Criteria <ul style="list-style-type: none"> $\pm 3i$ or \pm (their k)i plotted correctly on the imaginary axis, where $k \in \mathbb{R}, k > 0$ dependent on the final M mark being awarded in part (b) Their final two roots of the form $\lambda \pm \mu i, \lambda, \mu \neq 0$, are plotted correctly 	
		Satisfies at least one of the criteria	B1ft
		Satisfies both criteria with some indication of scale or coordinates stated with at least one pair of roots symmetrical about the real axis	B1ft
			(2)
			8
Question 5 Notes			
5. (a)	Note	Give B1B0 for writing down a correct $(z^2 - 6z + 25)$, followed by $a = 25, b = -6$	
	Note	If the values of a and b are not stated , then <ul style="list-style-type: none"> give B1B1 for writing down a correct $(z^2 - 6z + 25)$, give B1B0 for writing down $(z^2 + \text{their "a"}z + \text{their "b"})$, with exactly one of their a or their b correct 	
(b)	Note	No working leading to $z = 3i, -3i$ is 1 st M1 1 st A1	
	Note	$z = \pm \sqrt{9}i$ unless recovered is 1 st M0 1 st A0	
	Note	You can assume $x \equiv z$ for solutions in this question	
	Note	<ul style="list-style-type: none"> Give 2nd M1 2nd A1 for $z^2 - 6z + 25 = 0 \Rightarrow z = 3 + 4i, 3 - 4i$ with no intermediate working. Give 2nd M1 2nd A1 for $z = 3 + 4i, 3 - 4i$ with no intermediate working having stated $a = -6, b = 25$ in part (a) or part (b). Otherwise, give 2nd M0 2nd A0 for $z = 3 + 4i, 3 - 4i$ with no intermediate working. 	

Question 5 Notes Continued		
5. (b)	Note	Special Case: If their <i>3-term quadratic</i> factor $z^2 + "a"z + "b"$ can be factorised then give Special Case 2 nd M1 for correct factorisation leading to $z = \dots$
	Note	Otherwise, give 2 nd M0 for applying a method of factorisation to solve their 3TQ.
	Note	Reminder: Method Mark for solving a 3TQ, " $az^2 + bz + c = 0$ " Formula: Attempt to use the correct formula (with values for a, b and c) Completing the square: $\left(z \pm \frac{b}{2}\right)^2 \pm q \pm c = 0, q \neq 0$, leading to $z = \dots$
5. (b)(c)	Note	You can mark part (b) and part (c) together

Question Number	Scheme	Notes	Marks	
6.	Given $f(x) = \frac{2(x^3 + 3)}{\sqrt{x}} - 9, x > 0$; Roots $\alpha, \beta: 0.4 < \alpha < 0.5$ and $1.2 < \beta < 1.3$			
(a)	$\left\{ \begin{aligned} f(x) &= 2x^{\frac{5}{2}} + 6x^{-\frac{1}{2}} - 9 \Rightarrow \\ f'(x) &= 5x^{\frac{3}{2}} - 3x^{-\frac{3}{2}} \end{aligned} \right\}$	Some evidence of $\pm \lambda x^n \rightarrow \pm \mu x^{n-1}; \lambda, \mu \neq 0$	M1	
		Differentiates to give $\pm Ax^{\frac{3}{2}} \pm Bx^{-\frac{3}{2}}; A, B \neq 0$	M1	
	Correct differentiation which can be simplified or un-simplified		A1	
	$\left\{ \alpha \approx 0.45 - \frac{f(0.45)}{f'(0.45)} \right\} \Rightarrow \alpha \approx 0.45 - \frac{0.2159541693...}{-8.428734015...}$	Valid attempt at Newton-Raphson using their values of $f(0.45)$ and $f'(0.45)$	M1	
	$\{\alpha = 0.4756211869...\} \Rightarrow \alpha = 0.476$ (3 dp)	dependent on all 4 previous marks 0.476 on their first iteration (Ignore any subsequent iterations)	A1 cso	
	Correct differentiation followed by a correct answer of 0.476 scores full marks in part (a) Correct answer with no working scores no marks in part (a)		(5)	
(a) Alt 1	Alternative method 1 for the first 3 marks			
	$\left\{ \begin{aligned} u &= 2x^3 + 6 & v &= \sqrt{x} \\ u' &= 6x^2 & v' &= \frac{1}{2}x^{-\frac{1}{2}} \end{aligned} \right\} \Rightarrow$	Some evidence of $\pm \lambda x^n \rightarrow \pm \mu x^{n-1}; \lambda, \mu \neq 0$	M1	
		Differentiates to give $\frac{\pm Ax^2(\sqrt{x}) \pm Bx^{-\frac{1}{2}}(2x^3 + 6)}{x}; A, B \neq 0$	M1	
	$f'(x) = \frac{6x^2(\sqrt{x}) - \frac{1}{2}x^{-\frac{1}{2}}(2x^3 + 6)}{x}$	Correct differentiation which can be simplified or un-simplified	A1	
(b)	Either <ul style="list-style-type: none"> $\frac{\beta - 1.2}{"0.3678924937..."} = \frac{1.3 - \beta}{"0.1161410527..."}$ $\frac{\beta - 1.2}{1.3 - \beta} = \frac{"0.3678924937..."}{"0.1161410527..."}$ $\frac{\beta - 1.2}{"0.3678924937..."} = \frac{1.3 - 1.2}{"0.1161410527..." + "0.3678924937..."}$ 		At least one of either \pm (awrt 0.37, trunc. 0.36, awrt 0.12, or trunc. 0.11) This mark may be implied.	B1
			A correct linear interpolation method. Do not allow this mark if a total of one or three negative lengths are used or if either fraction is the wrong way up. This mark may be implied.	M1
	<ul style="list-style-type: none"> $\beta = \left(\frac{(1.3)("0.3678924937...") + (1.2)("0.1161410527...")}{"0.1161410527..." + "0.3678924937..."} \right)$ $= \left(\frac{0.4782602418... + 0.1393692632...}{0.4840335464...} \right) = \left(\frac{0.617629505...}{0.484033546...} \right)$ $\beta = 1.2 + \left(\frac{"0.3678924937..."}{"0.1161410527..." + "0.3678924937..."} \right) (0.1)$ $\beta = 1.2 + \left(\frac{"-0.3678924937..."}{"-0.1161410527..." + "-0.3678924937..."} \right) (0.1)$ 		dependent on the previous M mark Rearranges to give $\beta = \dots$	dM1
	$\{\beta = 1.276005578...\} \Rightarrow \beta = 1.276$ (3 dp)	1.276 (Ignore any subsequent iterations)	A1 cao	
			(4)	
			9	

Question Number	Scheme		Notes	Marks
6. (b) Way 2	$\frac{x}{\text{"0.3678924937..."}} = \frac{0.1 - x}{\text{"0.1161410527..."/>$		At least one of either \pm (awrt 0.37, trunc. 0.36, awrt 0.12, or trunc. 0.11) This mark may be implied.	B1
	$x = \frac{(0.1)(\text{"0.3678924937..."})}{0.4840335464...} = 0.0760055778...$			
	$\Rightarrow \beta = 1.2 + 0.0760055778...$		Finds x using a correct method of similar triangles and applies "1.5 + their x "	M1 dM1
	$\{\beta = 1.276005578...\} \Rightarrow \beta = 1.276 \text{ (3 dp)}$		1.276	A1 cao
(b) Way 3	$\frac{0.1 - x}{\text{"0.3678924937..."}} = \frac{x}{\text{"0.1161410527..."/>$		At least one of either \pm (awrt 0.37, trunc. 0.36, awrt 0.12, or trunc. 0.11) This mark may be implied.	B1
	$x = \frac{(0.1)(\text{"0.1161410527..."})}{0.4840335464...} = 0.0239944222...$			
	$\Rightarrow \beta = 1.3 - 0.0239944222...$		Finds x using a correct method of similar triangles and applies "1.6 - their x "	M1 dM1
	$\{\beta = 1.276005578...\} \Rightarrow \beta = 1.276 \text{ (3 dp)}$		1.276	A1 cao
Question 6 Notes				
6. (a)	Note	Incorrect differentiation followed by their estimate of α with no evidence of applying the NR formula is final dM0A0.		
	M1	This mark can be implied by applying at least one correct <i>value</i> of either $f(0.45)$ or $f'(0.45)$ to 1 significant figure in $0.45 - \frac{f(0.45)}{f'(0.45)}$. So just $0.45 - \frac{f(0.45)}{f'(0.45)}$ with an incorrect answer and no other evidence scores final dM0A0.		
	Note	You can imply the M1A1A1 marks for algebraic differentiation for either <ul style="list-style-type: none"> $f'(0.45) = 5(0.45)^{\frac{3}{2}} - 3(0.45)^{-\frac{3}{2}}$ $f'(1.5)$ applied correctly in $\alpha \approx 0.45 - \frac{2((0.45)^3 + 3) - 9}{\sqrt{0.45} \cdot (5(0.45)^{\frac{3}{2}} - 3(0.45)^{-\frac{3}{2}})}$ 		
(a) Alt 2	Alternative method 2 for the first 3 marks		Some evidence of $\pm \lambda x^n \rightarrow \pm \mu x^{n-1}$; $\lambda, \mu \neq 0$ Note: Allow M1 for either $\pm Ax^2(x^{-\frac{1}{2}})$ or $\pm Bx^{-\frac{3}{2}}(2x^3 + 6)$ or $\pm Bx^{-\frac{3}{2}}(x^3 + 3)$; $A, B \neq 0$	M1
	$\left\{ \begin{array}{l} u = 2x^3 + 6 \quad v = x^{-\frac{1}{2}} \\ u' = 6x^2 \quad v' = -\frac{1}{2}x^{-\frac{3}{2}} \end{array} \right\} \Rightarrow$			
	$f'(x) = 6x^2(x^{-\frac{1}{2}}) - \frac{1}{2}x^{-\frac{3}{2}}(2x^3 + 6)$			
		Differentiates to give $\pm Ax^2(x^{-\frac{1}{2}}) \pm Bx^{-\frac{3}{2}}(2x^3 + 6)$; $A, B \neq 0$	M1	
		Correct differentiation which can be simplified or un-simplified	A1	

Question 6 Notes Continued	
6. (b)	Note Condone writing the symbol α in place of β in part (b)
	Note $\frac{\beta - 1.2}{1.3 - \beta} = \left \frac{-0.3678924937\dots}{0.1161410527\dots} \right $ is a valid method for the first M mark
	Note Give 1 st M1 for either $\frac{-f(1.2)}{f(1.3)} = \frac{\beta - 1.2}{1.3 - \beta}$ or $\frac{ f(1.2) }{f(1.3)} = \frac{\beta - 1.2}{1.3 - \beta}$ or $\frac{ f(1.2) }{ f(1.3) } = \frac{\beta - 1.2}{1.3 - \beta}$
	Note Give M1M1 for the correct statement $\frac{1.3 f(1.2) + 1.2f(1.3)}{f(1.3) + f(1.2) }$
	Note Give M1M1 for the correct statement $\beta = \frac{1.3 + 1.2k}{k + 1}$, where $k = \frac{f(1.3)}{ f(1.2) } = \frac{0.116141\dots}{0.367892\dots} = 0.31569\dots$
	Note $\frac{\beta - 1.2}{1.3 - \beta} = \frac{"0.3678924937\dots"}{"0.1161410527\dots"} \Rightarrow \beta = 1.276$ with no intermediate working is B1 M1 dM1 A1
	Note $\frac{\beta - 1.2}{-0.3678924937\dots} = \frac{1.3 - \beta}{0.1161410527\dots} \Rightarrow \beta = 1.34613\dots = 1.346$ (3 dp) is B1 M0 dM0 A0
	Note $\frac{\beta - 1.2}{-0.3678924937\dots} = \frac{1.3 - \beta}{-0.1161410527\dots} \Rightarrow \beta = 1.276$ (3 dp) is B1 M1 dM1 A1

Question Number	Scheme	Notes	Marks
7.	$5x^2 - 4x + 3 = 0$ has roots α, β		
(a)	$\alpha + \beta = \frac{4}{5}, \alpha\beta = \frac{3}{5}$	Both $\alpha + \beta = \frac{4}{5}$ and $\alpha\beta = \frac{3}{5}$, seen or implied	B1
	$\frac{1}{\alpha^2} + \frac{1}{\beta^2} = \frac{\alpha^2 + \beta^2}{\alpha^2\beta^2}$	States or uses $\frac{1}{\alpha^2} + \frac{1}{\beta^2} = \frac{\alpha^2 + \beta^2}{\alpha^2\beta^2}$	M1
	$\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta = \dots$	Use of the correct identity for $\alpha^2 + \beta^2$ (May be implied by their work)	M1
	$\frac{1}{\alpha^2} + \frac{1}{\beta^2} = \frac{\alpha^2 + \beta^2}{\alpha^2\beta^2} = \frac{(\frac{4}{5})^2 - 2(\frac{3}{5})}{(\frac{3}{5})^2}$	Applies $\alpha^2\beta^2 = (\alpha\beta)^2$ correctly in the denominator of $\frac{\alpha^2 + \beta^2}{\alpha^2\beta^2}$ using their value of $\alpha\beta$	M1
	$= \frac{-(\frac{14}{25})}{(\frac{9}{25})} = -\frac{14}{9}$	dependent on ALL previous marks being awarded $-\frac{14}{9}$ or $-1\frac{5}{9}$ or -1.5 from correct working	A1 cso
			(5)
(b) Way 1	{Sum =} $\frac{3}{\alpha^2} + \frac{3}{\beta^2} = 3\left(-\frac{14}{9}\right) \left\{ = -\frac{14}{3} \text{ or } -\frac{42}{9} \right\}$	Simplifies $\frac{3}{\alpha^2} + \frac{3}{\beta^2}$ to give 3(their answer to (a))	M1
	{Product =} $\left(\frac{3}{\alpha^2}\right)\left(\frac{3}{\beta^2}\right) = \frac{9}{(\frac{3}{5})^2} \{ = 25 \}$	Applies $\frac{9}{(\text{their } \alpha\beta)^2}$ using their value of $\alpha\beta$	M1
	$x^2 + \frac{14}{3}x + 25 = 0$	Applies $x^2 - (\text{sum})x + \text{product}$ (can be implied), where sum and product are numerical values. Note: "=0" is not required for this mark	M1
	$3x^2 + 14x + 75 = 0$	Any integer multiple of $3x^2 + 14x + 75 = 0$, including the "=0"	A1
			(4)
			9
Question 7 Notes			
7. (a)	Note	Writing a correct $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$ without attempting to substitute at least one of either their $\alpha + \beta$ or their $\alpha\beta$ into $(\alpha + \beta)^2 - 2\alpha\beta$ is 2 nd M0	
	Note	Give B0M1M1M1A0 for $\alpha + \beta = -\frac{4}{5}, \alpha\beta = \frac{3}{5}$ leading to $\frac{1}{\alpha^2} + \frac{1}{\beta^2} = \frac{(-\frac{4}{5})^2 - 2(\frac{3}{5})}{(\frac{3}{5})^2} = -\frac{14}{9}$	
	Note	Writing down $\alpha, \beta = \frac{2 + \sqrt{11}i}{5}, \frac{2 - \sqrt{11}i}{5}$ and then stating $\alpha + \beta = \frac{4}{5}, \alpha\beta = \frac{3}{5}$ or applying $\alpha + \beta = \frac{2 + \sqrt{11}i}{5} + \frac{2 - \sqrt{11}i}{5} = \frac{4}{5}$ and $\alpha\beta = \left(\frac{2 + \sqrt{11}i}{5}\right)\left(\frac{2 - \sqrt{11}i}{5}\right) = \frac{3}{5}$ scores B0	
	Note	Those candidates who then apply $\alpha + \beta = \frac{4}{5}, \alpha\beta = \frac{3}{5}$, having written down/applied $\alpha, \beta = \frac{2 + \sqrt{11}i}{5}, \frac{2 - \sqrt{11}i}{5}$, can only score the M marks in part (a)	
	Note	Give B0M0M0M0A0 for $\frac{1}{\alpha^2} + \frac{1}{\beta^2} = \frac{1}{\left(\frac{2 + \sqrt{11}i}{5}\right)^2} + \frac{1}{\left(\frac{2 - \sqrt{11}i}{5}\right)^2} = -\frac{14}{9}$	

Question 7 Notes Continued		
7. (a)	Note	Give BOM1MOM0A0 for $\frac{1}{\alpha^2} + \frac{1}{\beta^2} = \frac{\alpha^2 + \beta^2}{\alpha^2 \beta^2} = \frac{\left(\frac{2+\sqrt{11}i}{5}\right)^2 + \left(\frac{2-\sqrt{11}i}{5}\right)^2}{\left(\frac{2+\sqrt{11}i}{5}\right)^2 \left(\frac{2-\sqrt{11}i}{5}\right)^2} = -\frac{14}{9}$
	Note	Give BOM1MOM0A0 for $\frac{1}{\alpha^2} + \frac{1}{\beta^2} = \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha^2 \beta^2} = \frac{\left(\frac{2+\sqrt{11}i}{5} + \frac{2-\sqrt{11}i}{5}\right)^2 - 2\left(\frac{2+\sqrt{11}i}{5}\right)\left(\frac{2-\sqrt{11}i}{5}\right)}{\left(\frac{2+\sqrt{11}i}{5}\right)^2 \left(\frac{2-\sqrt{11}i}{5}\right)^2} = -\frac{14}{9}$
	Note	Allow B1 for both $S = \frac{4}{5}$ and $P = \frac{3}{5}$ or for $\sum = \frac{4}{5}$ and $\prod = \frac{3}{5}$
	Note	Give final A0 for e.g. -1.55 or -1.5556 without reference to $-\frac{14}{9}$ or $-1\frac{5}{9}$ or $-1.\dot{5}$
	Note	Give 2 nd M1 for applying their $\alpha + \beta = \frac{4}{5}$ on $5\alpha^2 - 4\alpha + 3 = 0, 5\beta^2 - 4\beta + 3 = 0 \Rightarrow 5(\alpha^2 + \beta^2) - 4(\alpha + \beta) + 6 = 0$ to give $5(\alpha^2 + \beta^2) - 4\left(\frac{4}{5}\right) + 6 = 0 \left\{ \Rightarrow \alpha^2 + \beta^2 = \frac{-6 + \frac{16}{5}}{5} = -\frac{14}{25} \right\}$
(b)	Note	A correct method leading to $a=3, b=14, c=75$ without writing a final answer of $3x^2 + 14x + 75 = 0$ is final M1A0
	Note	Using $\frac{2+\sqrt{11}i}{5}, \frac{2-\sqrt{11}i}{5}$ explicitly , to find the sum and product of $\frac{3}{\alpha^2}$ and $\frac{3}{\beta^2}$ to give $x^2 + \frac{14}{3}x + 25 = 0 \Rightarrow 3x^2 + 14x + 75 = 0$ scores M0M0M1A0 in part (b)
	Note	Using $\frac{2+\sqrt{11}i}{5}, \frac{2-\sqrt{11}i}{5}$ to find $\alpha + \beta = \frac{4}{5}, \alpha\beta = \frac{3}{5}, \frac{1}{\alpha^2} + \frac{1}{\beta^2} = -\frac{14}{9}$ and applying $\left\{ \alpha + \beta = \frac{4}{5}, \right\} \alpha\beta = \frac{3}{5}, \frac{1}{\alpha^2} + \frac{1}{\beta^2} = -\frac{14}{9}$ can potentially score full marks in part (b). E.g. <ul style="list-style-type: none">• Sum = $\frac{3}{\alpha^2} + \frac{3}{\beta^2} = 3\left(-\frac{14}{9}\right) = -\frac{14}{3}$• Product = $\left(\frac{3}{\alpha^2}\right)\left(\frac{3}{\beta^2}\right) = \frac{9}{\left(\frac{3}{5}\right)^2} = 25$• $x^2 + \frac{14}{3}x + 25 = 0 \Rightarrow 3x^2 + 14x + 75 = 0$
	Note	Finding $\frac{1}{\alpha^2} + \frac{1}{\beta^2} = -\frac{14}{9}$ and correctly writing $x^2 - 3\left(\frac{1}{\alpha^2} + \frac{1}{\beta^2}\right)x + \frac{9}{(\alpha\beta)^2} = 0$ followed by $x^2 - \frac{14}{3}x + 25 = 0 \Rightarrow 3x^2 - 14x + 75 = 0$ (incorrect substitution of $\frac{1}{\alpha^2} + \frac{1}{\beta^2} = -\frac{14}{9}$) is M0M1M1A0

Question Number	Scheme	Notes	Marks
7.	$5x^2 - 4x + 3 = 0$ has roots α, β		
(b) Way 2	$y = \frac{3}{x^2} \Rightarrow x = \frac{3}{y^2} \Rightarrow 5\left(\frac{3}{y}\right) - 4\sqrt{\frac{3}{y}} + 3 = 0$	Substitutes $x^2 = \frac{3}{y}$ into $5x^2 - 4x + 3 = 0$	M1
	$\frac{15}{y} + 3 = 4\sqrt{\frac{3}{y}} \Rightarrow \left(\frac{15}{y} + 3\right)^2 = \left(4\sqrt{\frac{3}{y}}\right)^2$	dependent on the previous M mark Correct method for squaring both sides of their equation	dM1
	$\frac{225}{y^2} + \frac{45}{y} + \frac{45}{y} + 9 = 16\left(\frac{3}{y}\right)$		
	$\frac{225}{y^2} + \frac{42}{y} + 9 = 0$		
	$9y^2 + 42y + 225 = 0$	dependent on the previous M mark Obtains an expression of the form $ay^2 + by + c, a, b, c \neq 0$ Note: "= 0" not required for this mark	dM1
		<i>Any integer multiple</i> of $3y^2 + 14y + 75 = 0$, or $3x^2 + 14x + 75 = 0$, including the "= 0"	A1
			(4)

Question Number	Scheme	Notes	Marks
8.	$\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}^n = \begin{pmatrix} a^n & 0 \\ \frac{a^n - b^n}{a-b} & b^n \end{pmatrix}; n \in \mathbb{Z}^+; a \neq b$		
	$n=1$, LHS = $\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}$, RHS = $\begin{pmatrix} a & 0 \\ \frac{a-b}{a-b} & b \end{pmatrix} = \begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}$	Shows or states that either LHS = RHS = $\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}$ or LHS = $\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}$ or $\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}^1$, RHS = $\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}$	B1
	(Assume the result is true for $n=k$)		
	$\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}^{k+1} = \begin{pmatrix} a^k & 0 \\ \frac{a^k - b^k}{a-b} & b^k \end{pmatrix} \begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}$ or $\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix} \begin{pmatrix} a^k & 0 \\ \frac{a^k - b^k}{a-b} & b^k \end{pmatrix}$	$\begin{pmatrix} a^k & 0 \\ \frac{a^k - b^k}{a-b} & b^k \end{pmatrix}$ multiplied by $\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}$ (either way round)	M1
	$= \begin{pmatrix} a^{k+1} & 0 \\ \frac{a(a^k - b^k)}{a-b} + b^k & b^{k+1} \end{pmatrix}$ or $\begin{pmatrix} a^{k+1} & 0 \\ a^k + \frac{b(a^k - b^k)}{a-b} & b^{k+1} \end{pmatrix}$ or e.g. $\begin{pmatrix} a^{k+1} & 0 \\ \frac{a(a^k - b^k)}{a-b} + \frac{b^k(a-b)}{(a-b)} & b^{k+1} \end{pmatrix}$	Multiplies out to give a correct un-simplified matrix	A1
	$= \begin{pmatrix} a^{k+1} & 0 \\ \frac{a^{k+1} - b^{k+1}}{a-b} & b^{k+1} \end{pmatrix}$	dependent on the previous A mark Achieves this result with no algebraic errors	A1
	If the result is <u>true for $n=k$</u> , then it is <u>true for $n=k+1$</u> . As the result has been shown to be <u>true for $n=1$</u> , then the result is true for all $n \in \mathbb{Z}^+$		A1 cso
			(5)
			5
Question 8 Notes			
8.	Note	Final A1 is dependent on all previous marks being scored. It is gained by candidates conveying the ideas of all four underlined points either at the end of their solution or as a narrative in their solution.	
	Note	Give B0 for stating LHS = RHS by itself with no reference to LHS = RHS = $\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}$	
	Note	Give B0 for just stating $\begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}^1 = \begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix}$	
	Note	E.g. $\begin{pmatrix} a^k & 0 \\ \frac{a^k - b^k}{a-b} & b^k \end{pmatrix} \begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix} = \begin{pmatrix} a^{k+1} & 0 \\ \frac{a^{k+1} - b^{k+1}}{a-b} & b^{k+1} \end{pmatrix}$ with no intermediate working is M1A0A0A0	
	Note	Writing $\begin{pmatrix} a^k & 0 \\ \frac{a^k - b^k}{a-b} & b^k \end{pmatrix} \begin{pmatrix} a & 0 \\ 1 & b \end{pmatrix} = \begin{pmatrix} a^{k+1} & 0 \\ \frac{a(a^k - b^k)}{a-b} + b^k & b^{k+1} \end{pmatrix} = \begin{pmatrix} a^{k+1} & 0 \\ \frac{a^{k+1} - b^{k+1}}{a-b} & b^{k+1} \end{pmatrix}$ is M1A1A1	

Question Number	Scheme	Notes	Marks
9.	(a) $\frac{z-ki}{z+3i} = i$ (b)(i) $k=4$ (ii) $k=1$		
(a) Way 1	$z - ki = i(z + 3i) \Rightarrow z - ki = iz - 3$ $\Rightarrow z - iz = -3 + ki \Rightarrow z(1 - i) = -3 + ki$ $\Rightarrow z = \frac{-3 + ki}{(1 - i)}$	Complete method of making z the subject	M1
		Correct expression for $z = \dots$	A1
	$z = \frac{(-3 + ki)(1 + i)}{(1 - i)(1 + i)} \left\{ = \frac{(-3 + ki)(1 + i)}{2} \right\}$	dependent on the previous M mark Multiplies numerator and denominator by the conjugate of the denominator	dM1
	$z = -\frac{(k+3)}{2} + \frac{(k-3)}{2}i$ *	Achieves the correct answer with no errors seen	A1* cs0
			(4)
(a) Way 2	$z - ki = i(z + 3i)$ $(x + yi) - ki = i(x + yi + 3i)$ $x + (y - k)i = -y - 3 + xi$ {Real \Rightarrow } $x = -y - 3$ {Imaginary \Rightarrow } $y - k = x$	Multiplies both sides by $(z + 3i)$, applies $z = x + yi$, o.e., multiplies out and attempts to equate both the real part and the imaginary part of the resulting equation	M1
		Both correct equations which can be simplified or un-simplified	A1
	$\left\{ \begin{array}{l} x + y = -3 \\ x - y = -k \end{array} \right\} \Rightarrow x = \frac{-k-3}{2}, y = \frac{k-3}{2}$	dependent on the previous M mark Obtains two equations both in terms of x and y and solves them simultaneously to give at least one of $x = \dots$ or $y = \dots$	dM1
	$\Rightarrow z = -\frac{(k+3)}{2} + \frac{(k-3)}{2}i$ *	Finds $x = \frac{-k-3}{2}, y = \frac{k-3}{2}$ and writes down the given result	A1* cs0
			(4)
(b)(i)	$\{k=4 \Rightarrow\} z = -\frac{(4+3)}{2} + \frac{(4-3)}{2}i \left\{ = -\frac{7}{2} + \frac{1}{2}i \right\}$ $\{ z = \} \sqrt{\left(-\frac{7}{2}\right)^2 + \left(\frac{1}{2}\right)^2}$	Some evidence of substituting $z = 4$ into the given expression for z and a full attempt at applying Pythagoras to find $ z $	M1
	$= \sqrt{\frac{50}{4}}, \sqrt{12.5}, \frac{\sqrt{50}}{2}, \frac{5}{2}\sqrt{2}$ or $\frac{5}{\sqrt{2}}$ or $\sqrt{\frac{25}{2}}$	Correct exact answer	A1
(ii)	$\{k=1 \Rightarrow\} z = -\frac{(1+3)}{2} + \frac{(1-3)}{2}i \left\{ = -2 - i \right\}$ $\arg z = -\pi + \tan^{-1}\left(\frac{1}{2}\right)$	Some evidence of substituting $z = 1$ into the given expression for z and uses trigonometry to find an expression for $\arg z$ in the range $(-3.14\dots, -1.57\dots)$ or $(-180^\circ, -90^\circ)$ or $(3.14\dots, 4.71\dots)$ or $(180^\circ, 270^\circ)$	M1
	$\{\arg z = -\pi + 0.463647\dots \Rightarrow\} \arg z = -2.677945\dots \left\{ = -2.678 \text{ (3 dp)} \right\}$	awrt -2.678	A1
			(4)
			8

Question Number	Scheme	Notes	Marks
9.	(a) $\frac{z - ki}{z + 3i} = i$ (b)(i) $k = 4$ (ii) $k = 1$		
(a) Way 3	$\frac{z - ki}{i} = z + 3i \Rightarrow \frac{iz + k}{(-1)} = z + 3i$	Complete method of making z the subject	M1
	$\Rightarrow -iz - k = z + 3i \Rightarrow -k - 3i = z + iz$ $\Rightarrow -k - 3i = z(1 + i)$ $\Rightarrow z = \frac{-k - 3i}{(1 + i)}$	Correct expression for $z = \dots$	A1
	$z = \frac{(-k - 3i)(1 - i)}{(1 + i)(1 - i)}$	dependent on the previous M mark Multiplies numerator and denominator by the conjugate of the denominator	dM1
	$z = -\frac{(k + 3)}{2} + \frac{(k - 3)}{2}i$ *	Achieves the correct answer with no errors seen	A1* cs0
			(4)
Question 9 Notes			
9. (a)	Note	Condone any of e.g. $z = -\frac{k + 3}{2} + \frac{k - 3}{2}i$ or $z = -\frac{(3 + k)}{2} + \frac{(-3 + k)}{2}i$ for the final A mark	
(b)(i)	Note	M1 can be implied by awrt 3.54 or truncated 3.53	
	Note	Give A0 for 3.5355... without reference to $\sqrt{\frac{50}{4}}$, $\sqrt{12.5}$, $\frac{\sqrt{50}}{2}$, $\frac{5}{2}\sqrt{2}$ or $\frac{5}{\sqrt{2}}$ or $\sqrt{\frac{25}{2}}$	
(b)(ii)	Note	Allow M1 (implied) for awrt -2.7 , truncated -2.6 , awrt -153° or awrt 207° or awrt 3.6	

Question Number	Scheme	Notes	Marks
10.	$H: xy = 144; P\left(12p, \frac{12}{p}\right), p \neq 0, \text{ lies on } H.$ Normal to H at P crosses positive x -axis at Q and negative y -axis at R		
(a)	$y = \frac{144}{x} = 144x^{-1} \Rightarrow \frac{dy}{dx} = -144x^{-2} \text{ or } -\frac{144}{x^2}$	$\frac{dy}{dx} = \pm kx^{-2}; k \neq 0$	M1
	$xy = 144 \Rightarrow x\frac{dy}{dx} + y = 0$	Uses product rule to give $\pm x\frac{dy}{dx} \pm y$	
	$x = 12t, y = \frac{12}{t} \Rightarrow \frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = -\left(\frac{12}{t^2}\right)\left(\frac{1}{12}\right)$	their $\frac{dy}{dt} \times \frac{1}{\text{their } \frac{dy}{dt}}$; Condone $t \equiv p$	
	So at $P, m_T = -\frac{1}{p^2}$	Correct calculus work leading to $m_T = -\frac{1}{p^2}$	A1
	So, $m_N = p^2$	Applies $m_N = \frac{-1}{m_T}$, where m_T is found using calculus	M1
	<ul style="list-style-type: none"> $y - \frac{12}{p} = "p^2"(x - 12p) \text{ or}$ $\frac{12}{p} = "p^2"(12p) + c \Rightarrow y = "p^2"x + \text{their } c$ 	Correct straight line method for an equation of a normal where $m_N (\neq m_T)$ is found by using calculus.	M1
	Correct algebra leading to $y = p^2x + \frac{12}{p} - 12p^3$ *	Correct solution only	A1 *
Note: m_N must be a function of p for the 2 nd M1 and 3 rd M1 mark			(5)
(b)	$y = 0 \Rightarrow x_Q = 12p - \frac{12}{p^3}$	Puts $y = 0$ and finds x or puts $x = 0$ and finds y	M1
	$x = 0 \Rightarrow y_R = \frac{12}{p} - 12p^3$	At least one of x_Q or y_R correct, o.e.	A1
	$\left(12p - \frac{12}{p^3}, 0\right)$ and $\left(0, \frac{12}{p} - 12p^3\right)$	Both sets of coordinates correct. {Ignore labelling of coordinates}	A1
			(3)
(c)	Area $OQR = \frac{1}{2}\left(12p - \frac{12}{p^3}\right)\left(\frac{12}{p} - 12p^3\right) = 512$	$\frac{1}{2} \times (\pm \text{their } x_Q)(\pm \text{their } y_R) = 512$	M1
		Correct equation which can be un-simplified or simplified	A1
	$144p^4 - 1312 + \frac{144}{p^4} = 0$		
	$144p^8 - 1312p^4 + 144 = 0$ { $\Rightarrow 9p^8 - 82p^4 + 9 = 0$ }	Correct 3 term quadratic in p^4 Note: $144p^8 + 144 = 1312p^4$ is acceptable for this mark	A1
	$(9p^4 - 1)(p^4 - 9) = 0 \Rightarrow p^4 = \dots$	dependent on the previous M mark Uses a 3TQ in p^4 (or an implied 3TQ in p^4) to find at least one value of $p^4 = \dots$	dM1
$p = \sqrt{3}$ and $p = -\frac{1}{\sqrt{3}}$	Obtains both $p = \sqrt{3}$ and $p = -\frac{1}{\sqrt{3}}$ only Note: Allow $p = -\frac{\sqrt{3}}{3}$ in place of $p = -\frac{1}{\sqrt{3}}$	A1	
			(5)
			13

Question Number	Scheme	Notes	Marks
10. (c)	$\text{Area } OQR = \frac{1}{2} \left(12p - \frac{12}{p^3} \right) \left \left(\frac{12}{p} - 12p^3 \right) \right = 512$	$\frac{1}{2} \times (\pm \text{ their } x_Q)(\pm \text{ their } y_R) = 512$	M1
		Correct equation which can be un-simplified or simplified	A1
	$144 \left(p - \frac{1}{p^3} \right) \left(p^3 - \frac{1}{p} \right) = 1024 \Rightarrow p^4 - 2 + \frac{1}{p^4} = \frac{1024}{144}$		
	$\left(p^2 - \frac{1}{p^2} \right)^2 = \frac{64}{9} \Rightarrow p^2 - \frac{1}{p^2} = \pm \frac{8}{3}$		
	$3p^4 - 8p^2 - 3 = 0 \text{ and } 3p^4 + 8p^2 - 3 = 0$	Both correct 3 term quadratics in p^2 Note: Both $p^4 - 1 = \frac{8}{3}p^2$ and $3p^4 + 8p^2 = 3$ is acceptable for this mark	A1
	$(3p^2 + 1)(p^2 - 3) = 0 \Rightarrow p^2 = \dots$ or $(3p^2 - 1)(p^2 + 3) = 0 \Rightarrow p^2 = \dots$	dependent on the previous M mark Uses a 3TQ in p^2 (or an implied 3TQ in p^2) to find at least one value of $p^2 = \dots$	dM1
	$p = \sqrt{3} \text{ and } p = -\frac{1}{\sqrt{3}}$	Obtains both $p = \sqrt{3}$ and $p = -\frac{1}{\sqrt{3}}$ only	A1

(5)

Question 10 Notes

10. (a)	Note	Allow $y = p^2x - 12p^3 + \frac{12}{p}$ {order of terms interchanged in $y = \dots$ } for final A1
(b)	Note	For the accuracy marks in part (b) allow equivalents such as <ul style="list-style-type: none"> • $x = 12p - \frac{12}{p^3}$ or $x = \frac{12p^4 - 12}{p^3}$ or $x = \frac{12(p^2 - 1)(p^2 + 1)}{p^3}$ • $y = \frac{12}{p} - 12p^3$ or $y = \frac{12 - 12p^4}{p}$
(c)	Note	Give 1 st M1, 1 st A1 for <ul style="list-style-type: none"> • $\frac{1}{2} \left(12p - \frac{12}{p^3} \right) \left \left(\frac{12}{p} - 12p^3 \right) \right = 512$ {correct use of modulus} • $\frac{1}{2} \left(12p - \frac{12}{p^3} \right) \left(12p^3 - \frac{12}{p} \right) = 512$ {modulus has been applied here} • $-\frac{1}{2} \left(12p - \frac{12}{p^3} \right) \left(\frac{12}{p} - 12p^3 \right) = 512$ {modulus has been applied here}
	Note	Give 1 st M1, 1 st A0 for $\frac{1}{2} \left(12p - \frac{12}{p^3} \right) \left(\frac{12}{p} - 12p^3 \right) = 512$ {modulus has not been applied on y_R }
	Note	Writing a correct $144p^4 - 1312 + \frac{144}{p^4} = 0$ o.e. followed by a correct e.g. $p^4 = 9$ with no intermediate working is 2 nd A0, 2 nd M1
	Note	Writing a correct $144p^4 - 1312 + \frac{144}{p^4} = 0$ o.e. followed by $p^4 = 9$ and $p^4 = \frac{1}{9}$ with no intermediate working is 2 nd A1 (implied), 2 nd M1

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom