



Pearson  
Edexcel

Mark Scheme (Provisional)

Summer 2021

Pearson Edexcel International Advanced Level  
In Further Pure Mathematics F1  
(WFM01/01)

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2021

Question Paper Log Number P66060A

Publications Code WFM01\_01\_2106\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2021

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

# PEARSON EDEXCEL IAL MATHEMATICS

## General Instructions for Marking

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\surd$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - o.e. – or equivalent (and appropriate)
  - d... or dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper or ag- answer given
  - $\square$  or d... The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao), unless shown, for example, as A1ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
  
6. If a candidate makes more than one attempt at any question:
  - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
  
7. Ignore wrong working or incorrect statements following a correct answer.

## General Principles for Further Pure Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles)

### **Method mark for solving 3 term quadratic:**

#### **1. Factorisation**

$(x^2 + bx + c) = (x + p)(x + q)$ , where  $|pq| = |c|$ , leading to  $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$ , where  $|pq| = |c|$  and  $|mn| = |a|$ , leading to  $x = \dots$

#### **2. Formula**

Attempt to use the correct formula (with values for  $a$ ,  $b$  and  $c$ ).

#### **3. Completing the square**

Solving  $x^2 + bx + c = 0$ :  $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$ ,  $q \neq 0$ , leading to  $x = \dots$

### **Method marks for differentiation and integration:**

#### **1. Differentiation**

Power of at least one term decreased by 1. ( $x^n \rightarrow x^{n-1}$ )

#### **2. Integration**

Power of at least one term increased by 1. ( $x^n \rightarrow x^{n+1}$ )

## **Use of a formula**

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values but may be lost if there is any mistake in the working.

## **Exact answers**

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Question Number	Scheme	Notes	Marks
<b>1</b>	$f(x) = x^3 + 4x - 6$		
<b>(i)(a)</b>	$f(1) = -1$ $f(1.5) = 3.375 \left( = \frac{27}{8} \right)$	Attempts to evaluate at both end points. If substitution not seen accept $f(1) = -1$ <b>or</b> $f(1.5) = 3.375$ as evidence, with any value for the other end.	M1
	Sign change and $f(x)$ continuous therefore $\alpha$ is between $x = 1$ and $x = 1.5$	$f(1) = -1$ <b>and</b> $f(1.5) = 3.375$ <u>both correct</u> and mentions/indicates <u>sign change</u> , <u>continuous</u> and <u>conclusion</u> .	A1
			(2)
<b>(i)(b)</b>	$f'(x) = 3x^2 + 4$	Correct derivative. Can be implied by a correct expression seen later such as $3(1.5)^2 + 4$	B1
	$x_2 = 1.5 - \frac{f(1.5)}{f'(1.5)} = 1.5 - \frac{3.375}{10.75}$	Attempt Newton-Raphson using the <b>correct</b> formula.	M1
	$x_2 = 1.186\dots$	Accurate first application either awrt $1.186$ , $\frac{51}{43}$ or a correct numerical expression.	A1
	$x_3 = \left( 1.186\dots - \frac{0.412\dots}{8.220\dots} \right) = 1.1358\dots$		
	$\alpha \approx 1.136$	cao	A1
			(4)
<b>(ii)</b>	$g(1.4) = 3.442116\dots$ <b>or</b> $g(1.5) = -3.601419\dots$	Evidence of at least one value correct to 3 d.p. or better. May be implied by correct answer if never seen.	B1
	$\frac{1.5 - \beta}{\beta - 1.4} = \frac{0 - g(1.5)}{g(1.4) - 0}$ <b>or</b> $\frac{\beta - 1.4}{0 - 1.4} = \frac{1.5 - 1.4}{g(1.5) - g(1.4)}$ <b>oe</b>	A correct linear interpolation statement as shown <b>oe</b> (with correct signs). May omit the zeroes or use evaluated values (e.g 0.1 instead of $1.5 - 1.4$ ). Other forms are possible.	B1
	$\Rightarrow \beta \approx 1.448869\dots$	Evaluates $\beta$ from an attempt at a linear interpolation statement, allow if signs are incorrect. $\beta \approx$ awrt 1.449	M1 A1
			(4)
			<b>Total 10</b>



Qn No	Scheme	Notes	Marks	
<b>2. (a)</b>	$\frac{z_2 z_3}{z_1} = \frac{(p-i)(p+i)(2+i)}{(2-i)(2+i)}$	Multiply top and bottom by complex conjugate of their denominator. (The two M's may be scored if the given numbers are wrongly placed.)	M1	
	$= \frac{(p^2+1)(2+i)}{5}$	Simplifies numerator with evidence that $i^2 = -1$ and denominator real. Accept any equivalent form in the numerator as long as there are not $i^2$ terms if expanded.	M1	
	$= \frac{2(p^2+1)}{5} + \frac{(p^2+1)}{5}i$	Correct real +imaginary form with i factored out. Accept as single fraction with numerator in correct form. Accept 'a=' and 'b='.	A1	
			<b>(3)</b>	
<b>ALT</b>	$\frac{z_2 z_3}{z_1} = \frac{(p-i)(p+i)}{(2-i)} = a+bi$ $p^2+1 = (a+bi)(2-i)$ $\left. \begin{aligned} 2a+b &= p^2+1 \\ 2b-a &= 0 \end{aligned} \right\}$	Cross multiplies by $2-i$ (or their denominator), expands and equates real and imaginary parts. (The two M's may be scored if the given numbers are wrongly placed.)	M1	
	$\left. \begin{aligned} 2a+b &= p^2+1 \\ 2b-a &= 0 \end{aligned} \right\} \Rightarrow a = \dots, b = \dots$	Attempt to solve their equations.	M1	
	$a+bi = \frac{2(p^2+1)}{5} + \frac{(p^2+1)}{5}i$	Correct real +imaginary form with i factored out. Accept as single fraction with numerator in correct form. Accept 'a=' and 'b='.	A1	
			<b>(3)</b>	
	<b>(b)</b>	$\left  \frac{z_2 z_3}{z_1} \right ^2 = \frac{4(p^2+1)^2}{25} + \frac{(p^2+1)^2}{25}$	Correct attempt at the modulus or modulus squared. Accept with their answers to part (a). Any erroneous i or $i^2$ is M0.	M1
		$\frac{4(p^2+1)^2}{25} + \frac{(p^2+1)^2}{25} = (2\sqrt{5})^2$	Their $\left  \frac{z_2 z_3}{z_1} \right ^2 = (2\sqrt{5})^2$	dM1
$(p^2+1)^2 = 100 \Rightarrow p = \pm 3$		Attempt to solve and achieves $p = \dots$ (may be scored from use of $ \dots ^2 = 2\sqrt{5}$ ) $p = \pm 3$	M1 A1	
			<b>(4)</b>	
<b>ALT 1</b>	$\left  \frac{z_2 z_3}{z_1} \right  = 2\sqrt{5} \Rightarrow  z_2 z_3  = \sqrt{4+1} \times 2\sqrt{5}$	Cross multiplies and attempts $ z_1 $	M1	
	$\Rightarrow  z_2 ^2 = \sqrt{4+1} \times 2\sqrt{5} \Rightarrow p^2+1 = \dots$	Attempts $ z_2 z_3 $ either directly or using $ z_2 z_2^*  =  z_2 ^2$ to get an equation in p.	dM1	
	$(p^2+1) = 10 \Rightarrow p = \pm 3$	Attempt to solve and achieves $p = \dots$ $p = \pm 3$	M1 A1	
			<b>(4)</b>	
			<b>Total 7</b>	

Question Number	Scheme	Notes	Marks
<b>3(a)</b>	$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 2 & 2 & 0 \\ 1 & 3 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 3 & 1 \\ -2 & -2 & 0 \end{pmatrix}$	Attempt to multiply in the correct order with at least four correct elements. May be done as three separate calculations, so look for at least 4 correct values.	M1
	(1,-2), (3,-2) and (1,0)	Accept as individual column vectors but not as a single 2x3 matrix.	A1
			(2)
<b>(b)</b>	Rotation	Accept rotate or turn oe.	B1
	<b>270° (anticlockwise) about the origin</b>	Accept -90° (anticlockwise) or 90° clockwise (must be stated) and (0,0) or O. <i>Assume anticlockwise unless otherwise stated.</i>	B1
			(2)
<b>(c)</b>	$\mathbf{Q} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \mathbf{R} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$	One correct, both correct	B1,B1
			(2)
<b>(d)</b>	$\mathbf{RQ} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	Multiplication in correct order for their matrices and at least 1 row or 1 column correct.	M1
	$= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	Correct matrix	A1
			(2)
<b>(e)</b>	Reflection	Correct type identified.	B1
	in (the line) $y = x$	Correct line of reflection specified, accepting equivalent forms (e.g. line at angle 45° (anticlockwise) to the (positive) x-axis).	B1
			(2)
			<b>Total 10</b>

Question Number	Scheme	Notes	Marks
<b>4(a)</b>	$y = 25x^{-1} \Rightarrow \frac{dy}{dx} = -25x^{-2}$ $xy = 25 \Rightarrow y + x \frac{dy}{dx} = 0$ $\frac{dy}{dx} = \frac{dy}{dt} \frac{dt}{dx} = -\frac{5}{t^2} \cdot \frac{1}{5}$	Attempts a derivative expression, such as $\frac{dy}{dx} = kx^{-2}$ or $x \frac{dy}{dx} = ky$ or $\frac{dy}{dx} = \frac{\text{their } \frac{dy}{dt}}{\text{their } \frac{dx}{dt}}$	M1
	$\frac{dy}{dx} = -25x^{-2}$ or $\frac{-y}{x}$ or $\frac{-1}{t^2}$	Correct derivative.	A1
	$y - \frac{5}{t} = -\frac{1}{t^2}(x - 5t)$	Uses $y - \frac{5}{t} = (\text{their gradient}) \times (x - 5t)$ or $y = (\text{their gradient})x + c$ using $x = 5t, y = \frac{5}{t}$ in an attempt to find $c$ . Their gradient must be a function of $t$ for marks to be awarded.	M1
	$t^2y + x = 10t$ *	cso	A1
			(4)
<b>Alt (a)</b>	Intersect when $(10t - t^2y)y = 25$	Substitutes given line equation into the curve equation to find intersections	M1
	$\Rightarrow t^2y^2 - 10ty + 25 = 0$	Correct equation	A1
	$\Rightarrow (ty - 5)^2 = 0 \Rightarrow y = \frac{5}{t}$ is single root.	Shows the equation has a single root.	M1
	Single root when $y = \frac{5}{t}$ means the line is the tangent to $H$ at $P$ .	All work correct, with correct conclusion.	A1
			(4)
<b>(b)</b>	$t^2(-5) + 15 = 10t$	Substitute $(15, -5)$ into equation of tangent	M1
	$(t+3)(t-1) = 0 \Rightarrow t = ..$	Solves via any valid means	M1
	$\Rightarrow t = -3, t = 1$ (or one correct point)	Both values of $t$ or one correct coordinate pair.	A1
	$\left(-15, -\frac{5}{3}\right)$ and $(5, 5)$	Both correct sets of coordinates.	A1
			(4)
<b>ALT</b>	$t_1^2y + x = 10t_1$ $t_2^2y + x = 10t_2$	Form equations with their $t_1, t_2$ and attempt to solve for $x$ and $y$	M1
	$(x =) \frac{10t_1t_2}{(t_1+t_2)} = 15, (y =) \frac{10}{(t_1+t_2)} = -5$	Equates to coordinate and attempt to solve for $t_1, t_2$	M1
	$t_1 = -3, t_2 = 1$ (or one correct point)	Both values of $t$ or one correct coordinate pair.	A1
	$\left(-15, -\frac{5}{3}\right)$ and $(5, 5)$	Both correct sets of coordinates.	A1
			(4)
			<b>Total 8</b>

Question Number	Scheme	Notes	Marks
	$f(x) = (9x^2 + d)(x^2 - 8x + (10d + 1))$		
<b>5(a)</b>	$9x^2 + d = 0 \Rightarrow x = \pm\sqrt{\frac{-d}{9}}$ or $\pm\frac{i\sqrt{d}}{3}$	or exact equivalents	B1
	or $x = \frac{8 \pm \sqrt{64 - 4(10d + 1)}}{2}$ or $(x - 4)^2 - 16 + 10d + 1 = 0 \Rightarrow x = \dots$	Solve $x^2 - 8x + (10d + 1) = 0$ by formula or completing the square. Must have complete constant term.	M1
	$x = 4 + \sqrt{15 - 10d}$ and $x = 4 - \sqrt{15 - 10d}$	oe with discriminant simplified. Mark final answer, do not isw.	A1
			<b>(3)</b>
<b>(b)</b>	$x = \pm\frac{2i}{3}$	Correct roots for their answer for the $9x^2 + d$	B1ft
	$x = 4 \pm 5i$	Correct roots for their 3TQ	B1ft
	SC Award B1ftB0 if only one of <b>each</b> pair is given.		
			<b>(2)</b>
<b>(c)</b>		Two roots on imaginary axis the same distance from $O$ . Follow through their <i>imaginary</i> roots from (b). B0 if real roots found.	B1ft
		<p>Their two <i>complex</i> roots with real and imaginary parts, one the conjugate of the other, so reflected in the real axis. Must be correct relative scale compared with the imaginary roots <b>if the first B1 in (c) has been awarded</b> (ie clearly further from <math>O</math> if correct, or f.t. their answers). But if first B0 has been given, ignore scales.</p> <p>Accept points or vectors.</p> <p>Complex numbers must be labelled in some way e.g. via scales or coordinates or vectors.</p>	B1ft
			<b>(2)</b>
			<b>Total 7</b>

Question Number	Scheme	Notes	Marks
<b>6(a)</b>	$y = \sqrt{8x^2} \Rightarrow \frac{dy}{dx} = \frac{1}{2} \sqrt{8x}^{-\frac{1}{2}} = \sqrt{2x}^{-\frac{1}{2}}$ $y^2 = 8x \Rightarrow 2y \frac{dy}{dx} = 8$ or $\frac{dy}{dx} = \frac{dy}{dt} \frac{dt}{dx} = 4 \cdot \frac{1}{4t}$	Attempts a derivative expression, such as $\frac{dy}{dx} = kx^{-\frac{1}{2}} \text{ or } ky \frac{dy}{dx} = c$ or their $\frac{dy}{dt} \times \left( \frac{1}{\text{their } \frac{dx}{dt}} \right)$	M1
	$\frac{dy}{dx} = \frac{1}{2} \sqrt{8x}^{-\frac{1}{2}} \left( = \sqrt{2x}^{-\frac{1}{2}} \right) \text{ or } 2y \frac{dy}{dx} = 8 \text{ or}$ $\frac{dy}{dx} = 4 \cdot \frac{1}{4t} \left( = \frac{1}{t} \right)$	Correct differentiation (need not have substituted for $t$ etc)	A1
	At $P$ , gradient of normal is $m_N = -p$	Correct gradient for the normal.	A1
	$y - 4p = -p(x - 2p^2)$	$y - 4p = (\text{their } m_N) \times (x - 2p^2) \text{ or}$ $y = (\text{their } m_N)x + c \text{ using}$ $x = 2p^2, y = 4p \text{ in an attempt to find } c.$ Their gradient must be a function of $p$ for marks to be awarded. Must use a changed gradient, not tangent gradient.	M1
	$y + px = 2p^3 + 4p^*$	cso	A1
			(5)
<b>(b)</b>	$y + qx = 2q^3 + 4q$	oe	B1
			(1)
<b>(c)</b>	<b>Note: part (c) appears as MMAAA on ePEN but is being marked MAMAA.</b>		
	$y + px = 2p^3 + 4p$ $y + qx = 2q^3 + 4q$	Attempt to solve simultaneous equations.	M1
	$px - qx = 2p^3 + 4p - 2q^3 - 4q$	A correct equation in only one variable	A1
	$(p - q)x = 2((p - q)(p^2 + pq + q^2) + 2(p - q))$	Attempts to simplify the expression to required form. E.g. factorise difference of two cubes, $p^3 - q^3 = (p - q)(p^2 + pq + q^2) \text{ or}$ equivalent work to enable $p - q$ to cancel.	M1
	$x = 2(p^2 + pq + q^2 + 2)^*$	cso for reaching the correct $x$ coordinate.	A1
	$y + 2p^3 + 2p^2q + 2pq^2 + 4p = 2p^3 + 4p$ $y = -2pq(p + q)^*$	cso for reaching both coordinates correctly.	A1
			(5)

<b>ALT</b>	$qy + pqx = 2p^3q + 4pq$ $py + pqx = 2pq^3 + 4pq$ $py - qy = 2pq^3 - 2p^3q$	Attempt to solve simultaneous equations.  A correct equation in only one variable	M1  A1
	$(p - q)y = -2pq(p^2 - q^2)$ $(p - q)y = -2pq(p - q)(p + q)$	Attempts to simplify the expression to required form. E.g. Attempt to factorise difference of two squares. $p^2 - q^2 = (p - q)(p + q)$ or equivalent work to enable $p - q$ to cancel.	M1
	$y = -2pq(p + q)^*$	cso for reaching the correct $y$ coordinate.	A1
	$-2p^2q^2 - 2pq^3 + pqx = 2p^3q + 4pq$ $x = 2(p^2 + pq + q^2 + 2)^*$	cso for reaching both coordinates correctly.	A1
			(5)
<b>ALT 2</b>	$-2pq(p + q) + 2p(p^2 + pq + q^2 + 2) =$ $-2p^2q - 2pq^2 + 2p^3 + 2p^2q + 2pq^2 + 4p$	Substitutes both coordinates into the equation for normal at $P$ and expands brackets.	M1
	$= 2p^3 + 4p$ so $N$ is on normal at $P$	Simplifies correctly to $2p^3 + 4p$ and deduces $N$ is on the normal at $P$	A1
	$-2pq(p + q) + 2q(p^2 + pq + q^2 + 2) =$ $-2p^2q - 2pq^2 + 2p^2q + 2pq^2 + 2q^3 + 4q$	Substitutes both coordinates into the equation for normal at $Q$ and expands brackets.	M1
	$= 2q^3 + 4q$ so $N$ is on normal at $Q$	Simplifies correctly to $2q^3 + 4q$ and deduces $N$ is on the normal at $Q$ but penalise only the first instance for missing the deduction.	A1
	As $N$ is on both normals it is therefore the intersection point of them.	Makes a suitable conclusion	A1
			(5)
<b>(d)</b>	$\text{Grad } PQ = \frac{4q - 4p}{2q^2 - 2p^2} \left( = \frac{2}{p + q} \right)$	Use of $\frac{y_2 - y_1}{x_2 - x_1}$ and substituting.	M1A1
	$\text{Grad } ON = \frac{-2pq(p + q)}{2(p^2 + pq + q^2 + 2)}$	Can be unsimplified.	B1
	$\text{Grad } PQ \times \text{Grad } ON$ $= \frac{2}{p + q} \times \frac{-2pq(p + q)}{2(p^2 + pq + q^2 + 2)} = -1$	Using product of their gradients = -1	M1
	$(p + q)^2 - 3pq = -2$	-2	A1
			(5)
			<b>Total 16</b>

Question Number	Scheme	Notes	Marks
7	$\sum_{r=1}^n r^2 = \frac{n}{6}(n+1)(2n+1)$		
(a)	$n = 1, \text{LHS} = 1^2 = 1, \text{RHS} = \frac{1}{6} \cdot 2 \cdot 3 = 1$	Shows both LHS=1 and RHS=1 Accept LHS = 1 but must see at least $\frac{1}{6} \cdot 2 \cdot 3 = 1$ for RHS.	B1
	Assume true for $n = k$		
	When $n = k + 1$ $\sum_{r=1}^{k+1} r^2 = \frac{k}{6}(k+1)(2k+1) + (k+1)^2$	Adds $(k+1)^2$ to result for $n = k$	M1
	$= \frac{(k+1)}{6}(k(2k+1) + 6(k+1))$	Attempt to factorise by $\frac{(k+1)}{6}$	dM1
	$= \frac{(k+1)}{6}(k+2)(2k+3)$ $= \frac{(k+1)}{6}((k+1)+1)((2(k+1)+1))$	Either factorised form.	A1
	True for $n = 1$ . If true for $n = k$ then true for $n = k + 1$ therefore true for all $n$ .	Complete proof with no errors and these 4 statements seen anywhere. Depends on both M's and the A, but may be scored if the B is lost as long as some indication of true for $n = 1$ is given.	A1cso
			(5)
(b)	$\sum_{r=1}^n (r^2 + 2) = \sum_{r=1}^n r^2 + \sum_{r=1}^n 2$ $= \frac{n}{6}(n+1)(2n+1) + \dots$	Split into the addition of 2 sums and applies the result of (a).	M1
	$= \frac{n}{6}(n+1)(2n+1) + 2n$	Correct expression.	A1
	$= \frac{n}{6}(2n^2 + 3n + 13)$	Factorises out the $\frac{n}{6}$ - must have a common factor $n$ to achieve this mark; Simplifies to correct answer.	M1; A1
	$(a = 2, b = 3, c = 13)$		
			(4)
(c)	$\sum_{r=10}^{25} (r^2 + 2) = S_{25} - S_9$ $= \frac{25}{6} \cdot (2 \times 25^2 + 3 \times 25 + 13) - \frac{9}{6} \cdot (2 \times 9^2 + 3 \times 9 + 13)$	Attempts $S_{25} - S_9$ or $S_{25} - S_{10}$ with some substitution.	M1
	$= \frac{25}{6} \times 1338 - \frac{9}{6} \times 202 = 5575 - 303 = 5272$	For 5272	A1
	Note: Answer only (from calculator) is M0A0 as question requires use of part (b).		
			<b>Total 11</b>

Question Number	Scheme	Notes	Marks
	$f(n) = 4^{n+2} + 5^{2n+1}$ divisible by 21		
8	$n = 1, 4^3 + 5^3 = 189 = 9 \times 21$ (Or $n = 0, 4^2 + 5^1 = 21$ )	$f(1) = 21 \times 9$ Accept $f(0) = 21$ as an alternative starting point.	B1
	Assume that for $n = k, f(k) = (4^{k+2} + 5^{2k+1})$ is divisible by 21 for $k \in \mathbb{Z}^+$ .		
	$f(k+1) - f(k) = 4^{k+3} + 5^{2k+3} - (4^{k+2} + 5^{2k+1})$	Applies $f(k+1)$ with at least 1 power correct. May be just as $f(k+1)$ , or as part of an expression in $f(k+1)$ and $f(k)$ .	M1
	$= 4 \cdot 4^{k+2} + 25 \cdot 5^{2k+1} - 4^{k+2} - 5^{2k+1}$	For a correct expression in $f(k+1)$ , and possibly $f(k)$ , with powers reduced to those of $f(k)$ .	A1
	$= 3 \cdot 4^{k+2} + 24 \cdot 5^{2k+1}$		
	$= 3f(k) + 21 \cdot 5^{2k+1}$ or $= 24f(k) - 21 \cdot 4^{k+2}$	For one of these expression or equivalent with obvious factor of 21 in each.	A1
	$f(k+1) = 4f(k) + 21 \cdot 5^{2k+1}$	Makes $f(k+1)$ the subject or gives clear reasoning of each term other than $f(k+1)$ being divisible by 21. Dependent on at least one of the previous accuracy marks being awarded.	dM1
	{ $f(k+1)$ is divisible by 21 as both $f(k)$ and 21 are both divisible by 21 }		
	If the result is <b>true for <math>n = k</math></b> , then it is now <b>true for <math>n = k + 1</math></b> . As the result has shown to be <b>true for <math>n = 1</math> (or 0)</b> , then the result is <b>true for all <math>n \in \mathbb{Z}^+</math></b> .	Correct conclusion seen at the end. Condone true for $n = 1$ stated earlier. Depends on both M's and A's, but may be scored if the B is lost as long as at least $f(1) = 189$ was reached (so e.g. if the $21 \times 9$ was not shown)	A1 cso
			(6)
<b>ALT for first 4 marks</b>	$n = 1, 4^3 + 5^3 = 189 = 9 \times 21$ (Or $n = 0, 4^2 + 5^1 = 21$ )	As main scheme.	B1
	$f(k+1) - \alpha f(k) = 4^{k+3} + 5^{2k+3} - \alpha(4^{k+2} + 5^{2k+1})$	Attempts $f(k+1)$ in any equation (as main scheme).	M1
	$f(k+1) - \alpha f(k) = (4 - \alpha)4^{k+2} + (25 - \alpha)5^{2k+1}$	For a correct expression with any $\alpha$ , with powers reduced to match $f(k)$ .	A1
	$f(k+1) - \alpha f(k) = (4 - \alpha)(4^{k+2} + 5^{2k+1}) + 21 \cdot 5^{2k+1}$ $f(k+1) - \alpha f(k) = (25 - \alpha)(4^{k+2} + 5^{2k+1}) - 21 \cdot 4^{k+2}$	Any suitable equation with powers sorted appropriately to match $f(k)$	A1
	NB: $\alpha = 0, \alpha = 4, \alpha = 25$ will make relevant terms disappear, but marks should be awarded accordingly.		
			<b>Total 6</b>



Pearson Education Limited. Registered company number 872828  
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom