

Mark Scheme (Results)
January 2012

GCE Physics (6PH02) Paper 01
Physics at Work

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our qualifications website at www.edexcel.com.

For information about our BTEC qualifications, please call 0844 576 0026, or visit our website at www.btec.co.uk.

If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

Alternatively, you can contact our Science Subject Advisor directly by sending an email to ScienceSubjectAdvisor@EdexcelExperts.co.uk.

You can also telephone 0844 576 0037 to speak to a member of our subject advisor team.

(If you are calling from outside the UK please dial + 44 1204 770 696 and state that you would like to speak to the Science subject specialist).

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2012

Publications Code US030786

All the material in this publication is copyright

© Pearson Education Ltd 2012

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Mark scheme notes

Underlying principle

The mark scheme will clearly indicate the concept that is being rewarded, backed up by examples. It is not a set of model answers.

For example:

(iii) Horizontal force of hinge on table top

66.3 (N) or 66 (N) and correct indication of direction [no ue] ✓ 1
[Some examples of direction: acting from right (to left) / to the left / West / opposite direction to horizontal. May show direction by arrow. Do not accept a minus sign in front of number as direction.]

This has a clear statement of the principle for awarding the mark, supported by some examples illustrating acceptable boundaries.

1. Mark scheme format

- 1.1 You will not see 'wtte' (words to that effect). Alternative correct wording should be credited in every answer unless the ms has specified specific words that must be present. Such words will be indicated by underlining e.g. 'resonance'
- 1.2 Bold lower case will be used for emphasis.
- 1.3 Round brackets () indicate words that are not essential e.g. "(hence) distance is increased".
- 1.4 Square brackets [] indicate advice to examiners or examples e.g. [Do not accept gravity] [ecf].

2. Unit error penalties

- 2.1 A separate mark is not usually given for a unit but a missing or incorrect unit will normally mean that the final calculation mark will not be awarded.
- 2.2 Incorrect use of case e.g. 'Watt' or 'w' will not be penalised.
- 2.3 There will be no unit penalty applied in 'show that' questions or in any other question where the units to be used have been given, for example in a spreadsheet.
- 2.4 The same missing or incorrect unit will not be penalised more than once within one question (one clip in open).
- 2.5 Occasionally, it may be decided not to penalise a missing or incorrect unit e.g. the candidate may be calculating the gradient of a graph, resulting in a unit that is not one that should be known and is complex.
- 2.6 The mark scheme will indicate if no unit error penalty is to be applied by means of [no ue].

3. Significant figures

- 3.1 Use of an inappropriate number of significant figures in the theory papers will normally only be penalised in 'show that' questions where use of too few significant figures has resulted in the candidate not demonstrating the validity of the given answer.
- 3.2 The use of $g = 10 \text{ m s}^{-2}$ or 10 N kg^{-1} instead of 9.81 m s^{-2} or 9.81 N kg^{-1} will be penalised.

4. Calculations

- 4.1 Bald (i.e. no working shown) correct answers score full marks unless in a 'show that' question.
- 4.2 If a 'show that' question is worth 2 marks then both marks will be available for a reverse working; if it is worth 3 marks then only 2 will be available.
- 4.3 use of the formula means that the candidate demonstrates substitution of physically correct values, although there may be conversion errors e.g. power of 10 error.
- 4.4 recall of the correct formula will be awarded when the formula is seen or implied by substitution.
- 4.5 The mark scheme will show a correctly worked answer for illustration only.
- 4.6 Example of mark scheme for a calculation:

'Show that' calculation of weight

Use of $L \times W \times H$ ✓

Substitution into density equation with a volume and density ✓

Correct answer [49.4 (N)] to at least 3 sig fig. [No ue] ✓
[If 5040 g rounded to 5000 g or 5 kg, do not give 3rd mark; if conversion to kg is omitted and then answer fudged, do not give 3rd mark]
[Bald answer scores 0, reverse calculation 2/ 3]

3

Example of answer:

$$80 \text{ cm} \times 50 \text{ cm} \times 1.8 \text{ cm} = 7200 \text{ cm}^3$$

$$7200 \text{ cm}^3 \times 0.70 \text{ g cm}^{-3} = 5040 \text{ g}$$

$$5040 \times 10^{-3} \text{ kg} \times 9.81 \text{ N/ kg}$$

$$= 49.4 \text{ N}$$

5. Quality of Written Communication

- 5.1 Indicated by QoWC in mark scheme. QWC – Work must be clear and organised in a logical manner using technical wording where appropriate.
- 5.2 Usually it is part of a max mark, the final mark not being awarded unless the QoWC condition has been satisfied.

6. Graphs

- 6.1 A mark given for axes requires both axes to be labelled with quantities and units, and drawn the correct way round.
- 6.2 Sometimes a separate mark will be given for units or for each axis if the units are complex. This will be indicated on the mark scheme.
- 6.3 A mark given for choosing a scale requires that the chosen scale allows all points to be plotted, spreads plotted points over more than half of each axis and is not an awkward scale e.g. multiples of 3, 7 etc.
- 6.4 Points should be plotted to within 1 mm.
 - Check the two points furthest from the best line. If both OK award mark.
 - If either is 2 mm out do not award mark.
 - If both are 1 mm out do not award mark.
 - If either is 1 mm out then check another two and award mark if both of these OK, otherwise no mark.
- 6.5 For a line mark there must be a thin continuous line which is the best-fit line for the candidate's results.

Question Number	Answer	Mark
1	B	1
2	A	1
3	C	1
4	D	1
5	C	1
6	B	1
7	A	1
8	D	1
9	D	1
10	D	1

Question Number	Answer	Mark
11	<p>Indication that 500 W is 15% of incident radiation / Apply 15% efficiency to incident flux of 210 W m⁻² (i.e. find useful input) (1)</p> <p>Use of radiation flux is power per unit area (1)</p> <p>Answer = 16 m² (1)</p> <p>[2.38 m² is the answer without applying 15% → 1 mark]</p> <p><u>Example of calculation</u></p> <p>Input power = (500 × 100)/15 = 3300 W</p> <p>Area = input power/radiation flux = 3300 W / 210 W m⁻² = 16 m²</p> <p>OR</p> <p>(15/100) × 210 W m⁻² = 31.5 W m⁻²</p> <p>Area = 500 W / 31.5 W m⁻² = 16 m²</p>	3
Total for question 11		3

Question Number	Answer	Mark
12(a)	Use of $v = f\lambda$ (1) $f = 7.3 \text{ Hz}$ [accept 7.3 s^{-1} , do not accept fractions] (1) <u>Example of calculation</u> $f = 330 \text{ m s}^{-1} / 45 \text{ m}$ $f = 7.3 \text{ Hz}$	2
12(b)	Diffraction / it diffracts (1) Either an explanation of diffraction in general: Idea that the waves spread out (not bending) OR a diagram showing diffraction OR An explanation of why the tiger is heard: diffraction is significant for an obstacle (not a gap) of a size similar to the wavelength OR a diagram showing diffraction over a hill (1)	2
Total for question 12		4

Question Number	Answer	Mark
13(a)	Tick in Ultrasound box only (1)	1
13 (b)	A polarised wave is when the oscillations/vibrations are in one plane only which includes direction of travel (of the wave). Or A polarised wave is when the oscillations/vibrations are in one direction only which is perpendicular to the direction of travel (of the wave). Or Describes polarisation as a process where oscillations/vibrations in many planes are reduced to oscillations/vibrations in one plane [References to displacement are only acceptable in the context of varying displacement] (1) Longitudinal waves oscillate/vibrate in one direction which is the direction of travel of the wave / parallel to the direction of travel of the wave. (1)	2
Total for question 13		3

Question Number	Answer	Mark
14(a)	Resistance of parallel combination much less than resistance of V_1 (1) (Therefore) voltage of parallel combination is much less than voltage of V_1 (1) Or Identifies current (nearly) zero (because of resistance of V_1 very large) (1) (So) p.d. across 10Ω is zero by $V = IR$ (1) (Credit for each marking point may be obtained by completing a calculation.)	2
14(b)	Identifies resistance of parallel combination is $5 M\Omega$ (1) Use of resistors in parallel formula (1) $R = 10 M\Omega$ (1)	3
Total for question 14		5

Question Number	Answer	Mark
15	(QWC – Work must be clear and organised in a logical manner using technical wording where appropriate) MAX 6 Mention of energy levels (ignore electron shells) (1) Electrons in atoms can only occupy certain (discrete) energy levels (1) Ground state is the lowest energy/level electron(s)/atom can occupy (1) Energy is transferred in the collisions (1) Electrons/atoms move to higher level / become excited (when they gain energy) (1) These electrons return (later) to lower level/ground state (1) By emitting energy in the form of photons / reducing their energy by emitting photons (1) Photons have a specific energy or frequency or reference to $E = hf$ or $E = E_2 - E_1$ (1)	Max 6
Total for question 15		6

Question Number	Answer	Mark
16(a)	State ($V = E - Ir$)	(1)
	Correct substitution	(1)
	p.d. = 11V	(1)
	OR	
	Use of ($V = E - Ir$) to attempt to find lost volts	(1)
	Subtraction from E	(1)
	p.d. = 11V	(1)
	OR	
	Use of $E = I(R+r)$ to attempt to find R	(1)
Use of $V = IR$ with the value of R calculated	(1)	
p.d. = 11V	(1)	
	<u>Example of calculation</u>	
	$V = 12 \text{ V} - 3 \times 10^{-3} \Omega \times 400 \text{ A}$	
	p.d. = 10.8 V	
16(b)	To prevent large energy dissipation / wire heating up / wire melting / large pd across the wires OR to allow a large current	(1)
	Resistance of cable low	(1)
	(cross-sectional) area large [Not surface area]	(1)
	[Reverse argument in terms of a thin wire acceptable for all points]	
	Total for question 16	6

Question Number	Answer	Mark
17(a)	($\Omega = \text{V A}^{-1}$) OR ($\Omega = \text{V/A}$) OR $R = V/I$ [OR volt in alternative equivalent units divided by ampere in alternative equivalent units, as long as Ω isn't part of it] [Units and quantities must not be mixed.]	(1)
17(b)(i)	Use of $R = V/I$ with values feasibly from the graph	(1)
	$R = 6.8 \Omega$ to 8.0Ω (marks not awarded if using a gradient)	(1)
17(b)(ii)	resistance of metal remains constant	(1)
	resistance of thermistor decreases (as p.d. increases)	(1)
17(b)(iii)	(Increasing) current leads to temperature increase / leads to thermistor 'heating up'	(1)
	More conduction electrons / more electrons released / more free electrons / more charge carriers / charge carrier density increased / n increases	(1)
	Total for question 17	7

Question Number	Answer	Mark
18(a)	<p>Use of $\sin i \times v_2 = \sin r \times v_1$ (1) $r = 90^\circ$ at critical angle (1) critical angle = 75° (1)</p> <p>Acceptable alternative: Use of ${}_1\mu_2 = v_1 / v_2$ (1) State $\sin c = 1 / \mu$ (1) $c = 75^\circ$ (1) ($\mu = 1.036$, but look out for effects of rounding on calculated angle)</p> <p><u>Example of calculation</u> $\sin c / 1 = 1.96 / 2.03$ $c = 75^\circ$</p>	3
18(b)	<p>It will be reflected (back into the core) / totally internally reflected (1)</p> <p>Reflection back into core may be shown on the diagram (allow e.c.f for value of c from (a))</p>	1
18(c)	<p>Most of the light will undergo repeated (total internal) reflection Or most light continually strikes at greater than the critical angle Or minimal light is lost through refraction (1)</p> <p>Light reaches the bottom of the curtain Or Rays hitting the bottom will escape Or light hits the bottom at less than the critical angle (1)</p>	2
Total for question 18		6

Question Number	Answer	Mark
19(a)	Use of $W = QV$ (1) Energy of electron = 1.6×10^{-14} (J) (1) <u>Example of calculation</u> Energy = $1.6 \times 10^{-19} \times 100\,000$ J Energy of electron = 1.6×10^{-14} J	2
19(b)	Use of energy = power \times time (1) Energy = 2.88×10^7 (J) (1) <u>Example of calculation</u> Energy = $1000 \times 8 \times 3600$ Energy = 2.88×10^7 J	2
19(c)	Max 2 eV is very much smaller than Joule (1) kW h is very much bigger than Joule (1) in these units, answers more easily obtained from information available OR answers can be found without conversions (1)	Max 2
Total for question 19		6

Question Number	Answer	Mark
20(a)	<p>(QWC – Work must be clear and organised in a logical manner using technical wording where appropriate)</p> <p>Reference to photons (may be descriptive, e.g. quantum of energy / light arrives in small packets / light particles ...) (1)</p> <p>Energy of photon greater than or equal to work function (of zinc) / $hf \geq \phi$ (1)</p> <p>Results in electron being emitted (1)</p> <p>So (electroscope) loses charge / charge decreases (and leaf falls) (1)</p>	4
20(b)	<p>Photon energy (for visible light) is less than the work function OR frequency (of visible light) less than threshold frequency (1)</p>	1
20(c)	<p>Use of $c = f\lambda$ to find frequency (award if hc/λ used) (1)</p> <p>Use of $hf = \Phi + \frac{1}{2} m v^2$ to find KE (1)</p> <p>Use of ke equation with m_e (1)</p> <p>$v = 8.20 \times 10^5 \text{ m s}^{-1}$ (1)</p> <p><u>Example of calculation</u> $\text{KE} = (6.63 \times 10^{-34} \times 3 \times 10^8) / 200 \times 10^{-9} - 6.88 \times 10^{-19}$ $\text{KE} = 3.07 \times 10^{-19} \text{ J}$ $v = \sqrt{(2 \times 3.07 \times 10^{-19}) / 9.11 \times 10^{-31}}$ $v = 8.20 \times 10^5 \text{ m s}^{-1}$</p>	4
20(d)	<p>No change (1)</p> <p>Photon energy doesn't change (with distance) Or photon energy depends (only) on frequency/wavelength (1)</p>	2
Total for question 20		11

BLANK PAGE

Question Number	Answer	Mark
21(a)	Units of LHS m s^{-1} (1) (1) Units of T = kg m s^{-2} (1) (1) Units of $\mu = \text{kg m}^{-1}$ (1) (1)	3
21(b)(i)	Waves travel in both directions along wire OR reference to being reflected (not bounce) Waves superpose / interference effect / superposition occurs (not superimpose) Producing nodes and antinodes OR node is produced where waves are 180° out of phase / antiphase OR antinode is produced where waves are in phase OR node produced at a point of destructive interference OR antinode produced at a point of constructive interference OR produces points/ positions of constructive interference and points/ positions of destructive interference	3
21(b)(ii)	$\lambda = 4 \text{ m}$	1
21(b)(iii)	Substitution into the formula ignoring powers of ten $v = 173 \text{ m s}^{-1}$ <u>Example of calculation</u> $v = \sqrt{(150 \text{ N} / 0.005 \text{ kg m}^{-1})}$ $v = 173 \text{ m s}^{-1}$	2

<p>21(b)(iv)</p>	<p>Some of the marks may be gained from diagrams which show length remaining constant. Nodes and antinodes do not need to be labelled. Assume a sequence of diagrams shows increasing frequency.</p> <ul style="list-style-type: none"> • Wave speed const (1) • (As frequency increases) wavelength decreases (1) <p>Then max 3 from</p> <ul style="list-style-type: none"> • At most frequencies there is no standing wave / as frequency changes from a standing wave the wave no longer occurs / Standing waves only occur at some frequencies (1) • At higher frequencies there are more nodes / antinodes / loops (Not 'more waves') (1) • There is always a node at either end Or No of nodes = no of antinodes plus one (1) • Amplitude is less if there is a greater number of nodes (1) • Length = $n \lambda/2$ / after first standing wave, they occur when frequency x 2, x 3, x 4 etc / for frequency $n f_0$ (1) 	<p style="text-align: right;">Max 4</p>
	<p>Total for question 21</p>	<p style="text-align: right;">13</p>

Further copies of this publication are available from
International Regional Offices at www.edexcel.com/international

For more information on Edexcel qualifications, please visit
www.edexcel.com

Alternatively, you can contact Customer Services at
www.edexcel.com/ask or on + 44 1204 770 696

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual




Llywodraeth Cynulliad Cymru
Welsh Assembly Government

