

Write your name here

Surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

| | | | | | |
|--|--|--|--|--|--|
| | | | | | |
|--|--|--|--|--|--|

Candidate Number

| | | | | | |
|--|--|--|--|--|--|
| | | | | | |
|--|--|--|--|--|--|

Physics

Advanced

Unit 6: Experimental Physics

Wednesday 28 January 2015 – Morning
Time: 1 hour 20 minutes

Paper Reference

WPH06/01

You must have:
Ruler

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided – *there may be more space than you need.*

Information

- The total mark for this paper is 40.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*
- The list of data, formulae and relationships is printed at the end of this booklet.
- Candidates may use a scientific calculator.

Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P45040A

©2015 Pearson Education Ltd.

6/6/6/2/



PEARSON

Answer ALL questions in the spaces provided.

1 A student made measurements to determine the density of a single microscope slide.

(a) The dimensions of the slide were approximately 8 cm long, 3 cm wide and 1 mm thick.

Complete the table below to show the instruments you would use to make these measurements to an appropriate precision.

(4)

| Measurement | Instrument | Precision of instrument |
|-------------|------------|-------------------------|
| length | metre rule | 1 mm |
| width | | |
| thickness | | |

(b) The student recorded the following measurements.

| Measurement | Reading | Mean |
|----------------|--------------------------|------|
| length / mm | 75.8 75.9 75.7 75.8 | 75.8 |
| width / mm | 25.8 25.8 25.8 25.8 | 25.8 |
| thickness / mm | 1.01 1.02 0.98 0.99 1.00 | 1.00 |

Use these measurements to estimate the percentage uncertainty in the readings for length and thickness.

(2)

.....

.....

.....

.....

Percentage uncertainty in length = %

Percentage uncertainty in thickness = %



(c) The mass of the microscope slide is recorded as 4.82 g with an uncertainty of 0.03 g.

(i) Calculate a value for the density of the slide.

(2)

Density =

(ii) Estimate the percentage uncertainty in your value for the density.

You may assume the uncertainty in the measurement for the width is negligible.

(2)

Percentage uncertainty = %

(iii) The student researched the density of different types of glass and found a value for 'Crown glass' of $2600 \pm 100 \text{ kg m}^{-3}$.

Use this information to decide if the slide is made from Crown glass.

(2)

(d) Measuring the thickness of a stack of 10 slides would produce a better value for the thickness of one slide.

Explain why.

(2)

(Total for Question 1 = 14 marks)



(Total for Question 2 = 9 marks)




- 3 In an electrical circuit, as the frequency f of a signal generator is varied the potential difference V across a component also varies. This causes an electrical resonance effect which is similar to mechanical resonance.

The magnitude of V is a maximum at the resonant frequency.

In an experiment using a particular circuit the following measurements were recorded.

| f/kHz | V/mV |
|----------------|---------------|
| 5 | 3 |
| 10 | 5 |
| 15 | 11 |
| 20 | 22 |
| 25 | 27 |
| 27 | 19 |
| 30 | 15 |

- (a) A graph of V against f is plotted.



(i) Draw a line of best fit on the graph.

(2)

(ii) Estimate the maximum value of V .

(1)

(iii) State the value of the resonant frequency.

(1)

(b) Suggest one way in which the experiment could be improved to obtain a more accurate value of the resonant frequency.

(1)


(Total for Question 3 = 5 marks)



BLANK PAGE



- 4 A pendulum is made by tying a small mass on to a length of thread. A paper cone is placed over the mass as shown.



- (a) The pendulum is pulled to one side and released so that it oscillates. The amplitude of oscillation decreases because the cone provides damping.

- (i) Explain how you would use the apparatus above to measure the amplitude. You should add to the diagram.

(2)

.....

.....

.....

- (ii) State the uncertainty you would expect in the measurements of amplitude.

(1)

.....

.....



(b) It is predicted that the amplitude A is related to the number of oscillations n by

$$A = A_0 e^{-kn}$$

where A_0 is the initial amplitude and k is a constant.

Show that a graph of $\ln A$ against n should be a straight line.

(2)

.....

.....

.....

(c) In an experiment to measure A and n the following measurements were recorded.

| n | A/cm | |
|-----|---------------|--|
| 0 | 20.0 | |
| 2 | 18.2 | |
| 4 | 16.2 | |
| 6 | 14.4 | |
| 8 | 13.2 | |
| 10 | 11.9 | |

(i) Use the grid opposite to draw a graph of $\ln A$ against n . Use the column in the table for your processed data.

(5)

(ii) Use your graph to determine a value for k .

(2)


.....

.....

.....

$k =$





(Total for Question 4 = 12 marks)

TOTAL FOR PAPER = 40 MARKS



List of data, formulae and relationships

| | | |
|------------------------------|---|----------------------------|
| Acceleration of free fall | $g = 9.81 \text{ m s}^{-2}$ | (close to Earth's surface) |
| Boltzmann constant | $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$ | |
| Coulomb's law constant | $k = 1/4\pi\epsilon_0$ $= 8.99 \times 10^9 \text{ N m}^2 \text{ C}^{-2}$ | |
| Electron charge | $e = -1.60 \times 10^{-19} \text{ C}$ | |
| Electron mass | $m_e = 9.11 \times 10^{-31} \text{ kg}$ | |
| Electronvolt | $1 \text{ eV} = 1.60 \times 10^{-19} \text{ J}$ | |
| Gravitational constant | $G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$ | |
| Gravitational field strength | $g = 9.81 \text{ N kg}^{-1}$ | (close to Earth's surface) |
| Permittivity of free space | $\epsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1}$ | |
| Planck constant | $h = 6.63 \times 10^{-34} \text{ J s}$ | |
| Proton mass | $m_p = 1.67 \times 10^{-27} \text{ kg}$ | |
| Speed of light in a vacuum | $c = 3.00 \times 10^8 \text{ m s}^{-1}$ | |
| Stefan-Boltzmann constant | $\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$ | |
| Unified atomic mass unit | $u = 1.66 \times 10^{-27} \text{ kg}$ | |

Unit 1

Mechanics

| | |
|-------------------------------|--|
| Kinematic equations of motion | $v = u + at$ $s = ut + \frac{1}{2}at^2$ $v^2 = u^2 + 2as$ |
| Forces | $\Sigma F = ma$ $g = F/m$ $W = mg$ |
| Work and energy | $\Delta W = F\Delta s$ $E_k = \frac{1}{2}mv^2$ $\Delta E_{\text{grav}} = mg\Delta h$ |

Materials

| | |
|-----------------------|--|
| Stokes' law | $F = 6\pi\eta rv$ |
| Hooke's law | $F = k\Delta x$ |
| Density | $\rho = m/V$ |
| Pressure | $p = F/A$ |
| Young modulus | $E = \sigma/\epsilon$ where Stress $\sigma = F/A$ Strain $\epsilon = \Delta x/x$ |
| Elastic strain energy | $E_{\text{el}} = \frac{1}{2}F\Delta x$ |



Unit 2

Waves

Wave speed $v = f\lambda$

Refractive index ${}_1\mu_2 = \sin i / \sin r = v_1 / v_2$

Electricity

Potential difference $V = W/Q$

Resistance $R = V/I$

Electrical power, energy and efficiency
 $P = VI$
 $P = I^2R$
 $P = V^2/R$
 $W = VI t$

$$\% \text{ efficiency} = \frac{\text{useful energy output}}{\text{total energy input}} \times 100$$

$$\% \text{ efficiency} = \frac{\text{useful power output}}{\text{total power input}} \times 100$$

Resistivity $R = \rho l/A$

Current
 $I = \Delta Q / \Delta t$
 $I = nqvA$

Resistors in series $R = R_1 + R_2 + R_3$

Resistors in parallel $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3}$

Quantum physics

Photon model $E = hf$

Einstein's photoelectric equation
 $hf = \phi + \frac{1}{2}mv_{\max}^2$



Unit 4

Mechanics

| | |
|---|--|
| Momentum | $p = mv$ |
| Kinetic energy of a non-relativistic particle | $E_k = p^2/2m$ |
| Motion in a circle | $v = \omega r$ $T = 2\pi/\omega$ $F = ma = mv^2/r$ $a = v^2/r$ $a = r\omega^2$ |

Fields

| | |
|----------------------------|--|
| Coulomb's law | $F = kQ_1Q_2/r^2$ where $k = 1/4\pi\epsilon_0$ |
| Electric field | $E = F/Q$ $E = kQ/r^2$ $E = V/d$ |
| Capacitance | $C = Q/V$ |
| Energy stored in capacitor | $W = \frac{1}{2}QV$ |
| Capacitor discharge | $Q = Q_0 e^{-t/RC}$ |
| In a magnetic field | $F = BIl \sin \theta$ $F = Bqv \sin \theta$ $r = p/BQ$ |
| Faraday's and Lenz's Laws | $\epsilon = -d(N\phi)/dt$ |

Particle physics

| | |
|-----------------------|---------------------------|
| Mass-energy | $\Delta E = c^2 \Delta m$ |
| de Broglie wavelength | $\lambda = h/p$ |



Unit 5

Energy and matter

Heating $\Delta E = mc\Delta\theta$

Molecular kinetic theory $\frac{1}{2}m\langle c^2 \rangle = \frac{3}{2}kT$

Ideal gas equation $pV = NkT$

Nuclear Physics

Radioactive decay $dN/dt = -\lambda N$

$$\lambda = \ln 2/t_{1/2}$$

$$N = N_0 e^{-\lambda t}$$

Mechanics

Simple harmonic motion

$$a = -\omega^2 x$$
$$a = -A\omega^2 \cos \omega t$$
$$v = -A\omega \sin \omega t$$
$$x = A \cos \omega t$$
$$T = 1/f = 2\pi/\omega$$

Gravitational force $F = Gm_1 m_2 / r^2$

Observing the universe

Radiant energy flux $F = L/4\pi d^2$

Stefan-Boltzmann law

$$L = \sigma T^4 A$$
$$L = 4\pi r^2 \sigma T^4$$

Wien's Law $\lambda_{\max} T = 2.898 \times 10^{-3} \text{ m K}$

Redshift of electromagnetic radiation $z = \Delta\lambda/\lambda \approx \Delta f/f \approx v/c$

Cosmological expansion $v = H_0 d$



BLANK PAGE

