

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Friday 5 June 2020

Afternoon (Time: 1 hour 20 minutes)

Paper Reference **WPH16/01**

Physics

International Advanced Level

Unit 6: Practical Skills in Physics II

You must have:

Scientific calculator, Ruler

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - *there may be more space than you need.*
- **Show all your working out in calculations and include units where appropriate.**

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
 - *use this as a guide as to how much time to spend on each question.*
- The list of data, formulae and relationships is printed at the end of this booklet.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

P65751A

©2020 Pearson Education Ltd.

1/1/1/1/1/1/1/1/1




P 6 5 7 5 1 A 0 1 2 0



Pearson

Answer ALL questions.

- 1 A student investigated the conservation of momentum using the apparatus as shown.



- (a) The air track provides a cushion of air which reduces friction between the gliders and the track.

Describe how the student would show that the air track is horizontal before starting the investigation.

(1)

- (b) The student pushed glider A. The first light gate recorded the time t_1 for the card on glider A to pass through it.

The gliders collided and stuck together. The second light gate recorded the time t_2 for the card on glider A to pass through it.

The student recorded t_1 and t_2 for three separate collisions.

t_1 /s	0.34	0.21	0.28
t_2 /s	0.70	0.39	0.55

The masses of the gliders were identical. If momentum is conserved then $t_2 = 2t_1$.

Show that momentum was conserved in this investigation.

(3)

.....
.....
.....
.....
.....
.....
.....

- (c) Another student suggested that using a piece of card twice as long would improve the investigation.

Assess this suggestion.


(3)

.....
.....
.....
.....
.....
.....
.....

(Total for Question 1 = 7 marks)



- 2 A student wrote the following plan to investigate the distance travelled by alpha particles in air.



- Place the Geiger-Müller tube in front of the source.
- Measure the distance d from the source to the tube.
- Measure the count.
- Change d and take more readings.

Devise a more detailed plan for this investigation.

(Total for Question 2 = 6 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

BLANK PAGE




P 6 5 7 5 1 A 0 5 2 0

- 3 A bar magnet was suspended from a wooden retort stand as shown.



The magnet lined up with the magnetic field of the Earth.


The magnet was given a small angular displacement from its equilibrium position and oscillated in a horizontal plane about the string as shown.



- (a) Describe how the time period of these oscillations should be measured to make the readings as accurate as possible.

(3)

- (b) A loop of wire was placed vertically around the centre of the oscillating magnet as shown.



When the switch was closed, there was a current I in the loop of wire and the time period T of the oscillations decreased.

A student predicted that the relationship between T and I is

$$T = I^n$$

where n is a constant.

- (i) State an additional component required in the circuit that would allow this relationship to be investigated.

(1)

- (ii) Explain why plotting a graph of $\log T$ against $\log I$ would test the validity of this relationship.

(2)



(c) The student processed his results and produced the table below.

T/s	I/A		
0.813	1.20		
0.754	1.40		
0.706	1.60		
0.663	1.80		
0.631	2.00		
0.593	2.20		

(i) Plot a graph of $\log T$ against $\log I$ on the grid opposite.

Use the additional columns in the table to record your processed data.

(6)

(ii) Use your graph to determine a value for n .

(3)


.....
.....
.....
.....
 $n = \dots$



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 5 7 5 1 A 0 9 2 0

- (iii) After plotting the graph, the student modified his prediction. He suggested that the relationship between T and I is

$$T = kI^n$$

where k is a constant.


Justify this suggestion.

(4)

(Total for Question 3 = 19 marks)



- 4 The diagram shows a 100 g slotted mass drawn approximately to size.
A student determined the density of the metal from which the slotted mass was made.



- (a) (i) State the most appropriate measuring instrument for the student to use to measure the width a and the length b of the slot.

(1)

.....
.....
.....

- (ii) Explain one technique she should use when measuring a and b .

(2)

.....
.....
.....

- (iii) Calculate the area of the slot and its uncertainty in cm^2 .
Assume the slot is rectangular.

$$a = 0.47 \pm 0.01 \text{ cm}$$

$$b = 2.19 \pm 0.005 \text{ cm}$$

(3)

.....
.....
.....

$$\text{Area of the slot} = \dots \pm \dots \text{ cm}^2$$



(b) The student made a single measurement of the diameter of the slotted mass as 3.81 cm.

(i) Calculate the shaded area of the slotted mass in cm^2 .

(2)

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

$$\text{Shaded area} = \dots \text{cm}^2$$

(ii) Calculate the uncertainty in the value of the shaded area.

(3)

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

$$\text{Uncertainty} = \dots \text{cm}^2$$

(c) The student used a micrometer screw gauge to measure the thickness t of the slotted mass. She obtained the following results.

t/mm				$\text{mean } t/\text{mm}$
11.39	11.36	11.35	11.38	11.37

(i) Calculate the density ρ of the metal in g cm^{-3} . Assume the value of mass is 100 g with negligible uncertainty.

(2)

.....
.....
.....
.....
.....
.....
.....
.....
$$\rho = \dots \text{g cm}^{-3}$$



(ii) Calculate the percentage uncertainty in the value of ρ .

(3)

.....
.....
.....
.....
Percentage uncertainty =

(d) The student thinks that the slotted mass is made from brass, which has a density of 8.5 g cm^{-3} .

Determine whether the slotted mass could be made of brass.

(2)

.....
.....
.....
.....
(Total for Question 4 = 18 marks)

TOTAL FOR PAPER = 50 MARKS



List of data, formulae and relationships

Acceleration of free fall	$g = 9.81 \text{ m s}^{-2}$	(close to Earth's surface)
Boltzmann constant	$k = 1.38 \times 10^{-23} \text{ J K}^{-1}$	
Coulomb's law constant	$k = 1/4\pi\epsilon_0$ $= 8.99 \times 10^9 \text{ N m}^2 \text{ C}^{-2}$	
Electron charge	$e = -1.60 \times 10^{-19} \text{ C}$	
Electron mass	$m_e = 9.11 \times 10^{-31} \text{ kg}$	
Electronvolt	$1 \text{ eV} = 1.60 \times 10^{-19} \text{ J}$	
Gravitational constant	$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$	
Gravitational field strength	$g = 9.81 \text{ N kg}^{-1}$	(close to Earth's surface)
Permittivity of free space	$\epsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1}$	
Planck constant	$h = 6.63 \times 10^{-34} \text{ J s}$	
Proton mass	$m_p = 1.67 \times 10^{-27} \text{ kg}$	
Speed of light in a vacuum	$c = 3.00 \times 10^8 \text{ m s}^{-1}$	
Stefan-Boltzmann constant	$\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$	
Unified atomic mass unit	$u = 1.66 \times 10^{-27} \text{ kg}$	

Unit 1

Mechanics

Kinematic equations of motion

$$s = \frac{(u + v)t}{2}$$

$$v = u + at$$

$$s = ut + \frac{1}{2}at^2$$

$$v^2 = u^2 + 2as$$

Forces

$$\Sigma F = ma$$

$$g = \frac{F}{m}$$

$$W = mg$$

Momentum

$$p = mv$$

Moment of force

$$\text{moment} = Fx$$

Work and energy

$$\Delta W = F\Delta s$$

$$E_k = \frac{1}{2}mv^2$$

$$\Delta E_{\text{grav}} = mg\Delta h$$

Power

$$P = \frac{E}{t}$$

$$P = \frac{W}{t}$$



Efficiency

$$\text{efficiency} = \frac{\text{useful energy output}}{\text{total energy input}}$$

$$\text{efficiency} = \frac{\text{useful power output}}{\text{total power input}}$$

Materials**Density**

$$\rho = \frac{m}{V}$$

Stokes' law

$$F = 6\pi\eta rv$$

Hooke's law

$$\Delta F = k\Delta x$$

Elastic strain energy

$$\Delta E_{\text{el}} = \frac{1}{2} F\Delta x$$

Young modulus

$$E = \frac{\sigma}{\varepsilon} \text{ where}$$

$$\text{Stress } \sigma = \frac{F}{A}$$

$$\text{Strain } \varepsilon = \frac{\Delta x}{x}$$



Unit 2*Waves*

Wave speed

$$v = f\lambda$$

Speed of a transverse wave
on a string

$$v = \sqrt{\frac{T}{\mu}}$$

Intensity of radiation

$$I = \frac{P}{A}$$

Refractive index

$$n_1 \sin \theta_1 = n_2 \sin \theta_2$$

$$n = \frac{c}{v}$$

Critical angle

$$\sin C = \frac{1}{n}$$

Diffraction grating

$$n\lambda = d \sin \theta$$

Electricity

Potential difference

$$V = \frac{W}{Q}$$

Resistance

$$R = \frac{V}{I}$$

Electrical power, energy

$$P = VI$$

$$P = I^2 R$$

$$P = \frac{V^2}{R}$$

$$W = VIt$$

Resistivity

$$R = \frac{\rho l}{A}$$

Current

$$I = \frac{\Delta Q}{\Delta t}$$

$$I = nqvA$$

Resistors in series

$$R = R_1 + R_2 + R_3$$

Resistors in parallel

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3}$$

Quantum physics

Photon model

$$E = hf$$

Einstein's photoelectric
equation

$$hf = \phi + \frac{1}{2} mv_{\max}^2$$

de Broglie wavelength

$$\lambda = \frac{h}{p}$$



Unit 4*Mechanics*

Impulse

$$F\Delta t = \Delta p$$

Kinetic energy of a
non-relativistic particle

$$E_k = \frac{p^2}{2m}$$

motion in a circle

$$v = \omega r$$

$$T = \frac{2\pi}{\omega}$$

$$a = \frac{v^2}{r}$$

Centripetal force

$$F = ma = \frac{mv^2}{r}$$

$$F = m\omega^2 r$$

Electric and magnetic fields

Electric field

$$E = \frac{F}{Q}$$

Coulomb's law

$$F = \frac{Q_1 Q_2}{4\pi\epsilon_0 r^2}$$

$$E = \frac{Q}{4\pi\epsilon_0 r^2}$$

$$E = \frac{V}{d}$$

Electrical Potential

$$V = \frac{Q}{4\pi\epsilon_0 r}$$

Capacitance

$$C = \frac{Q}{V}$$

Energy stored in capacitor

$$W = \frac{1}{2} QV$$

$$W = \frac{1}{2} CV^2$$

$$W = \frac{1}{2} \frac{Q^2}{C}$$

Capacitor discharge

$$Q = Q_0 e^{-t/RC}$$



Resistor capacitor discharge

$$I = I_0 e^{-t/RC}$$

$$V = V_0 e^{-t/RC}$$

$$\ln Q = \ln Q_0 - \frac{t}{RC}$$

$$\ln I = \ln I_0 - \frac{t}{RC}$$

$$\ln V = \ln V_0 - \frac{t}{RC}$$

In a magnetic field

$$F = Bqv \sin \theta$$

$$F = BIl \sin \theta$$

Faraday's and Lenz's laws

$$\mathcal{E} = \frac{-d(N\phi)}{dt}$$

Nuclear and particle physics

In a magnetic field

$$r = \frac{p}{BQ}$$

Mass-energy

$$\Delta E = c^2 \Delta m$$



Unit 5*Thermodynamics*

Heating

$$\Delta E = mc\Delta\theta$$

$$\Delta E = L\Delta m$$

Ideal gas equation

$$pV = NkT$$

Molecular kinetic theory

$$\frac{1}{2}m \langle c^2 \rangle = \frac{3}{2}kT$$

Nuclear decay

Mass-energy

$$\Delta E = c^2 \Delta m$$

Radio-active decay

$$A = -\lambda N$$

$$\frac{dN}{dt} = -\lambda N$$

$$\lambda = \frac{\ln 2}{t_{1/2}}$$

$$N = N_0 e^{-\lambda t}$$

$$A = A_0 e^{-\lambda t}$$

Oscillations

Simple harmonic motion

$$F = kx$$

$$a = -\omega^2 x$$

$$x = A \cos \omega t$$

$$v = -A\omega \sin \omega t$$

$$a = A\omega^2 \cos \omega t$$

$$T = \frac{1}{f} = \frac{2\pi}{\omega}$$

$$\omega = 2\pi f$$

Simple harmonic oscillator

$$T = 2\pi \sqrt{\frac{m}{k}}$$

$$T = 2\pi \sqrt{\frac{l}{g}}$$



Astrophysics and Cosmology

Gravitational field strength
$$g = F/m$$

Gravitational force
$$F = \frac{Gm_1m_2}{r^2}$$

Gravitational field
$$g = \frac{Gm}{r^2}$$

Gravitational potential
$$V_{grav} = \frac{-Gm}{r}$$

Stephan-Boltzman law
$$L = \sigma T^4 A$$

Wein's law
$$\lambda_{\max} T = 2.898 \times 10^{-3} \text{ m K}$$

Intensity of radiation
$$I = \frac{L}{4\pi d^2}$$

Redshift of electromagnetic radiation
$$z = \frac{\Delta\lambda}{\lambda} \approx \frac{\Delta f}{f} \approx \frac{v}{c}$$

Cosmological expansion
$$v = H_0 d$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

