Mark Scheme (Results)

Summer 2013

International GCSE Chemistry (4CH0) Paper 1CR

Science Double Award (4SC0)
Paper 1CR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.btec.co.uk for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.

Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at www.edexcel.com/ask. You will need an Edexcel username and password to access this service.

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2013
Publications Code UG035595
All the material in this publication is copyright
© Pearson Education Ltd 2013

Question number	Answer	Accept	Reject	Marks
1 (a)	can <u>easily/quickly</u> identify each gas OR <u>less likely</u> to make a mistake in identification I GNORE just to identify the gas			1
(b) (i)	argon/Ar <u>and</u> helium/He			1
(ii)	oxygen/O ₂ I GNORE O			1
(iii)	air/it is a mixture (of gases) OR air/it is not a single substance I GNORE mixture of elements			1
(iv)	not flammable/not explosive/does not burn			
(c) (i)	hydrogen/H ₂ I GNORE H	air		1
(ii)	carbon dioxide/CO ₂			1
(iii)	carbon dioxide/CO ₂			1
			_	
			Total	8

Question number	Answer	Accept	Reject	Marks
2 (a)	D			1
(b)	M1 before heating – colourless (solution/liquid) I GNORE clear/transparent/looks like water M2 after heating – milky/chalky/cloudy/white (precipitate)/turbid	no colour	white solution/liquid any colour other than white	1
	I GNORE references to goes clear OWTTE			
(c)	M1 (sulfur dioxide/it) dissolves in/reacts with (rain) water	$SO_2 + H_2O \rightarrow H_2SO_3$ OR $SO_2 + H_2O +$ $1/2O_2 \rightarrow H_2SO_4$ for both M1 and M2		1
	M2 to form an acidic solution/an acid/sulfurous acid /acid rain I GNORE references to any other products whether correct or not	sulfuric acid		1
	M3 which reacts with/corrodes the marble/calcium carbonate	chemical weathering dissolves correct equation for reaction with either sulfurous or sulfuric acid		
		SO ₂ reacts with marble for M3 only		
	I GNORE erodes / weathers / melts / eats into			
			Total	6

Question number		Ans	sw er		Accept	Reject	Marks
3 (a)							
	Name of barium salt	Formula of barium salt	Solubility in water	Poisonous			
	barium chloride	BaCl ₂					1
	barium nitrate						
	barium carbonate	BaCO₃					1
	barium sulfate						
(b)	M1 (it forms) bai	rium chloride/Ba	Cl ₂ /a soluble (bari	um) salt			1
	M2 by reaction/	with hydrochloric	acid/stomach aci	d	by neutralisation	any suggestion that barium chloride is reacting	1
					word or chemical equation for 2 marks (equation can be unbalanced)	J	
(c)	barium sulfate/Ba	aSO ₄			be unbalanceu)		1

Question number	Answer	Accept	Reject	Marks
3 (d)	M1 barium sulfate is formed	'products', provided shown correctly in word equation		1
	M2 which is not poisonous/not toxic/harmless I GNORE references to magnesium hydroxide not poisonous	is insoluble		1
	M2 dep on M1			
	M3 barium hydroxide + magnesium sulfate → barium sulfate + magnesium hydroxide	$Ba(OH)_2 + MgSO_4$ $\rightarrow BaSO_4 +$ $Mg(OH)_2$		1
	OR barium ions + sulfate ions → barium sulfate	OR		
	Darium ions + Sunate ions → Darium Sunate	$Ba^{2+} + SO_4^{2-} \rightarrow BaSO_4$		
(e) (i)	M1 water – (reacts) <u>very/extremely</u> quickly/more quickly <u>than</u> <u>strontium</u> /quickest I GNORE rapidly/vigorously	explosively/violently		1
	M2 air – (reacts) <u>very/extremely</u> quickly/more quickly <u>than</u> <u>strontium</u> /quickest	explosively/violently		1
	(without heating) I GNORE rapidly/vigorously			1
(ii)		in a vacuum		1
(iii)	in/under any one of the following: (paraffin/mineral) oil/petroleum (oil)/(liquid) paraffin I GNORE in an air tight container reactivity <u>increases</u> as atomic number <u>increases</u>	reactivity increases with atomic number/down the group OWTTE reverse argument		1

	positive correlation		
		Total	12

Question number	Answer	Accept	Reject	Marks
4 (a)	M1 (negative electrode) – graphite	carbon		2
	M2 (positive electrode) – graphite	carbon		
(b) (i)	it/aluminium oxide/alumina has a (very) high m.pt I GNORE high b.pt/references to strong bonding/bauxite has a high m.pt/lot of energy		aluminium has a high melting point	1
(ii)	needed to melt it aluminium oxide/alumina is dissolved in/mixed with (molten/liquid) cryolite	added to Na ₃ AlF ₆ for cryolite cryolite is used as the solvent (for aluminium oxide/alumina)	aluminium is dissolved in cryolite	1
	I GNORE cryolite lowers the m.pt of aluminium oxide/alumina	oxide, didiffinia)		
(c)	M1 reduction		redox for M1 only	1
	M2 (it/aluminium ions/Al ³⁺) gain of electron(s) I GNORE references to loss of oxygen M2 dep on M1	reacts with/combines with decrease in oxidation number/oxidation number changes from +3 to 0	Al/aluminium gains electrons	1
(d)	M1 oxygen formed/produced (at the positive electrode/anode) I GNORE oxygen from the aluminium oxide	oxygen from the electrolysis	any indication that the oxygen is from the air for M1 only	1
	M2 reacts with the carbon/the (positive) electrode	anode / graphite	cathode/negative electrode	1
	M2 not dep on M1, but must mention oxygen			
(e)	Any two from:			2
	M1 malleable	easy to shape/easy to bend/easy to extrude bend		
	M2 low density			

M3 does not react with food/drink(s)	non-toxic/does not corrode		
I GNORE light(er)/high strength to weight ratio/references to cost/lightweight/does not ru	st		
		Total	10

Question number	Answer	Accept	Reject	Marks
5 (a)	M1 (molecules/compounds/substances) with the same molecular formula/number of each type of atoms	hydrocarbons	elements/atoms general formula/empirical formula for M1 only	1
	I GNORE chemical formula/same compound	atoms arranged differently		1
	M2 (but) different structural formulae/different displayed formulae/different structures			
(b)	D D			1
(c) (i)	M1 C _n H _{2n}	letters other than n, e.g. x	C _n +H _{2n}	1
(ii)	M1 double bond between two left hand end carbon atoms			1
	M2 single bond between each pair of rest of carbon atoms			1
	Penalise max 1 mark for any extra bond shown			
(d)	M1 addition	additional		1
	M2 orange	yellow/brown	red, either on its own or in combination with	
	M3 colourless I GNORE clear/transparent/looks like water		any other colour	1
(e)	M1 saturated – <u>all</u> (carbon to carbon) bonds are single /contains <u>only</u> (carbon to carbon)	does not contain any multiple/double bonds	5	1

single bonds		1
M2 unsaturated - contains (carbon to carbon) double/multiple bond(s)		
	Total	11

Question number	Answer	Accept	Reject	Marks
6 (a) (i)	7			1
(ii)	M1 solid			1
	M2 black	very dark grey		1
(iii)	M1 (formula) – HAt	AtH		1
	M2 (name) – hydrogen astatide	astatine hydride	hydrogen astati <u>n</u> e	1
(iv)		iodine is more reactive	_	1
	iodine, I) I GNORE astatine is unreactive	reverse argument	any references to astatide or iodide	1
	M2 – elements get less reactive with increasing	Astatine (atom) has more	louide	
	atomic number/as group is <u>descended</u> /the lower	(electron) shells/outer shell of astatine is further		
	they are	from nucleus so attracts		
	in the group	an <u>electron</u> less readily		
(b) (i)	4 (1) (1) 2 (1)	multiples/halves		1
(ii)	(paper) turns white/bleaches	(litmus) turns colourless		1
	I GNORE turns red			
(c) (i)	acid	correct formula		1
	I GNORE hydrogen ions/names of acids			1
(ii)	to displace (all of) the bromine / to react all of the bromide (ions)	bromine (an)ions for bromide to complete the reaction		
(iii)		to complete the reaction		2
	$Br_2 + SO_2 + 2H_2O \rightarrow 2HBr + H_2SO_4$	multiples and halves		
	M1 all formulae correct			
	M2 balanced			
(iv)	$2HBr + Cl_2 \rightarrow Br_2 + 2HCl$	multiples and halves		1

(d)	M1 colourless I GNORE clear/transparent/looks like water			1
	M2 brown (solution) / (dark) grey/black solid/precipitate	red- brown/orange/orange- brown	red on its own	1
			Total	16

Question number	Answ er	Accept	Reject	Marks
7 (a)	M1 (reactants) s aq	capital letters		1
	M2 (products) aq I g			1
(b) (i)	to prevent acid escaping/spraying out/spitting out I GNORE to prevent water escaping	solution/liquid/HCl		1
(ii)	С			1
(c) (i)	M1 A			1
	M2 gas produced/collected more quickly / experiment over	reaction is faster		1
	in shorter time / (gradient of) line steeper M2 dep on M1			1
(ii)	M1 0.1(0) M2 volume of gas is half/40 \div 80 = $\frac{1}{2}$ / 80 = 40	Half the products are produced		1
	x 2 M2 dep on M1			
(d) (i)	M1 & M2 - all points plotted to nearest gridline deduct 1 mark			2
	for each incorrect plot up to a max. of 2			1
	M3 suitable straight line of best fit (csq on plotted points)			
(ii)	(must be drawn with the aid of a ruler). Line need not beextrapolated.	(show a) positive correlation		1
	M1 as concentration increases rate increases	as one doubles the other doubles/directly proportional		1

	M2 proportional / in proportion	for 2 marks		
(iii)			molecules/atoms	1
, ,				1
	M1 more ions/particles (in a given volume) I GNORE more reactants			1
	M2 collide (successfully)		any reference to greater energy	
	M3 more per second/more frequently			
	Must be reference to frequency or number of collisions per unit time I GNORE greater chance of collision		5	
	TONOTIL greater chance of comsion		Total	16

Question	Answer	Accept	Reject	Marks
number		7100001	110,000	Warko
8 (a) (i)	Impurities/chemicals/substances may affect the colour/flame I GNORE affect the result/test			1
(ii)	colour can (easily) be seen (in a non-luminous flame)	a luminous flame may mask the colour		1
(iii)	I GNORE references to temperature yellow/orange/gold(en)	any combination of the acceptable colours, e.g. golden-yellow		1
(b) (i)	Li ⁺ and Ca ²⁺	lithium and calcium/Li and Ca	Ca ⁺ etc	1
(ii)	M1 – ammonia/NH ₃			1
	M2 – (water is needed) to form hydroxide ions/OH ⁻	to form an alkali/an alkaline solution/ammonium hydroxide		1
(iii)	M1 - iron(III)/Fe ³⁺	to dissolve the ammonia ammonia needs to be aqueous	any other oxidation states/ferrous	1
	M2 – ammonium/NH ₄ ⁺	ferric	ammonia	1
	If both names and formulae given both must be correct			
			Total	8


ct Marks
1
1
1
1
1
1
1
1
1
Total 9

Question number	Answer	Accept	Reject	Marks
10 (a)	 any two from: forward and backward reactions (still) occurring concentrations/amounts of reactants/products/components remain constant rate of forward reaction = rate of reverse reaction I GNORE concentrations/amounts of reactants and products are the same I GNORE reaction is reversible/goes both ways, OWTTE I GNORE references to le Chatelier 	both reactions (still) occurring stay the same in place of remain constant		2
(b) (i)	M1 - (increase in temperature) decrease(s) M2 - (increase in pressure) increase(s) M1 - (forward) reaction is exothermic/gives out heat	less/low <u>er(s)/drop(s)/gets</u> small <u>er</u> more/raise(s)/high <u>er</u> /gets bigg <u>er</u>		1 1 1
	OR reverse reaction is endothermic/takes in heat M2 – fewer (gas) molecules/particles on right hand side OR fewer moles (of gas) on right hand side I GNORE references to volumes I GNORE references to le Chatelier's principle I GNORE references to reverse reaction lowers the temperature I GNORE references to forward reaction reduces the pressure	reverse argument shifts to side with fewer (gas) molecules/fewer moles (of gas)	atoms	1

10	(c)	(i)	(the position of) equilibrium is not established/reached			1
		(ii)	M1 – (the mixture of gases is) cooled	temperature is decreased		1
		(iii)	M2 – ammonia liquefies / condenses recycled / <u>re</u> used / recirculated	put (back) into the reaction chamber used <u>again</u> (in the process)		1
	(d)		heat(ing) / energy costs would be higher	yield (of ammonia) would decrease		1
	(e)	(i)	M1 $M_r(N_2) = 28$	28 anywhere in the calculation		1
			M2 112 000 ÷ 28 (= 4 000) / 112 000 ÷ M1			1
			M3 8 000 / M2 x 2	112 ÷ 28) x 2 = 8 for 2 marks		1
				(112 000 ÷ 14) x 2 = 16 000 for 2 marks		
				Correct final answer without working for 3 marks		
		(ii)	1 200 / 15% of M3			1
					Total	15

Question number	Answer	Accept	Reject	Marks
11 (a)	(produces) most heat/energy per gram / per unit mass	highest temperature rise per gram / per unit mass	per amount	1
		most energy for smallest number of grams / mass		
(b)	(produces) most heat/energy per mole/per molecule /per amount	highest temperature rise per mole / per molecule		1
		most energy for smallest number of moles / molecules / amount		
(c)	Any two from: • heat/energy losses (e.g. by convection, by conduction, to air, to surroundings) • incomplete combustion • evaporation of water • copper / can / beaker / thermometer /apparatus absorbs heat • flame moves around because of draughts	• non-standard conditions		2
(d) (i)	A			1
(ii)	В			1
(e)	M1 breaking bonds is endothermic / takes in heat/energy	more energy is given out when bonds are made than is taken in		1
	M2 making bonds is exothermic / gives out	when bonds are broken for 3 marks		1
	heat/energy	more energy is given out when bonds are made than when bonds		1
	M3 more heat/energy given out than taken in	are broken for 1 mark		

I GNORE references to numbers/strengths of bonds		
	Total	9



Further copies of this publication are available from Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email <u>publication.orders@edexcel.com</u>
Order Code xxxxxxxx Summer 2013

For more information on Edexcel qualifications, please visit our website $\underline{www.edexcel.com}$

Pearson Education Limited. Registered company number 872828 with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE





