

Write your name here

Surname

Other names

Pearson Edexcel Certificate
Pearson Edexcel
International GCSE

Centre Number

--	--	--	--	--	--

Candidate Number

--	--	--	--	--

Chemistry

Unit: KCH0/4CH0

Paper: 2C

Thursday 16 January 2014 – Afternoon

Time: 1 hour

Paper Reference

KCH0/2C

4CH0/2C

You must have:

Ruler

Calculator

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- Show all the steps in any calculations and state the units.
- Some questions must be answered with a cross in a box (☒). If you change your mind about an answer, put a line through the box (☒) and then mark your new answer with a cross (☒).

Information

- The total mark for this paper is 60.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P42864A

©2014 Pearson Education Ltd.

1/1/1/



PEARSON

THE PERIODIC TABLE

Period 1 2 3 4 5 6 7 0

Group

	¹ H Hydrogen 1								⁴ He Helium 2																													
1																																						
2	⁷ Li Lithium 3	⁹ Be Beryllium 4							¹⁹ K Potassium 19	²⁰ Ca Calcium 20	³⁹ Sc Scandium 21	⁴⁰ Ti Titanium 22	⁴¹ V Vanadium 23	⁴² Cr Chromium 24	⁴³ Mn Manganese 25	⁴⁴ Fe Iron 26	⁴⁵ Co Cobalt 27	⁴⁶ Ni Nickel 28	⁴⁷ Cu Copper 29	⁴⁸ Zn Zinc 30	⁴⁹ Ga Gallium 31	⁵⁰ Ge Germanium 32	⁵¹ As Arsenic 33	⁵² Se Selenium 34	⁵³ Br Bromine 35	⁵⁴ Kr Krypton 36												
3	¹¹ Na Sodium 11	¹² Mg Magnesium 12							⁸¹ Rb Rubidium 37	⁸² Sr Strontium 38	⁸³ Y Yttrium 39	⁸⁴ Zr Zirconium 40	⁸⁵ Nb Niobium 41	⁸⁶ Mo Molybdenum 42	⁸⁷ Tc Technetium 43	⁸⁸ Ru Ruthenium 44	⁸⁹ Rh Rhodium 45	⁹⁰ Pd Palladium 46	⁹¹ Ag Silver 47	⁹² Cd Cadmium 48	⁹³ In Indium 49	⁹⁴ Sn Tin 50	⁹⁵ Sb Antimony 51	⁹⁶ Te Tellurium 52	⁹⁷ I Iodine 53	⁹⁸ Xe Xenon 54												
4	¹⁹ Fr Francium 87	²⁰ Ra Radium 88							¹⁰¹ La Lanthanum 57	¹⁰² Ce Cerium 58	¹⁰³ Pr Praseodymium 59	¹⁰⁴ Nd Neodymium 60	¹⁰⁵ Pm Promethium 61	¹⁰⁶ Sm Samarium 62	¹⁰⁷ Eu Europium 63	¹⁰⁸ Gd Gadolinium 64	¹⁰⁹ Tb Terbium 65	¹¹⁰ Dy Dysprosium 66	¹¹¹ Ho Holmium 67	¹¹² Er Erbium 68	¹¹³ Tm Thulium 69	¹¹⁴ Yb Ytterbium 70	¹¹⁵ Lu Lutetium 71	¹¹⁶ Hf Hafnium 72	¹¹⁷ Ta Tantalum 73	¹¹⁸ W Tungsten 74	¹¹⁹ Re Rhenium 75	¹²⁰ Os Osmium 76	¹²¹ Ir Iridium 77	¹²² Pt Platinum 78	¹²³ Au Gold 79	¹²⁴ Hg Mercury 80	¹²⁵ Tl Thallium 81	¹²⁶ Pb Lead 82	¹²⁷ Bi Bismuth 83	¹²⁸ Po Polonium 84	¹²⁹ At Astatine 85	¹³⁰ Rn Radon 86
5	²⁷ Fr Francium 87	²⁸ Ra Radium 88							¹³⁷ La Lanthanum 57	¹³⁸ Ce Cerium 58	¹³⁹ Pr Praseodymium 59	¹⁴⁰ Nd Neodymium 60	¹⁴¹ Pm Promethium 61	¹⁴² Sm Samarium 62	¹⁴³ Eu Europium 63	¹⁴⁴ Gd Gadolinium 64	¹⁴⁵ Tb Terbium 65	¹⁴⁶ Dy Dysprosium 66	¹⁴⁷ Ho Holmium 67	¹⁴⁸ Er Erbium 68	¹⁴⁹ Tm Thulium 69	¹⁵⁰ Yb Ytterbium 70	¹⁵¹ Lu Lutetium 71	¹⁵² Hf Hafnium 72	¹⁵³ Ta Tantalum 73	¹⁵⁴ W Tungsten 74	¹⁵⁵ Re Rhenium 75	¹⁵⁶ Os Osmium 76	¹⁵⁷ Ir Iridium 77	¹⁵⁸ Pt Platinum 78	¹⁵⁹ Au Gold 79	¹⁶⁰ Hg Mercury 80	¹⁶¹ Tl Thallium 81	¹⁶² Pb Lead 82	¹⁶³ Bi Bismuth 83	¹⁶⁴ Po Polonium 84	¹⁶⁵ At Astatine 85	¹⁶⁶ Rn Radon 86
6	⁸³ Fr Francium 87	⁸⁴ Ra Radium 88							¹⁷³ La Lanthanum 57	¹⁷⁴ Ce Cerium 58	¹⁷⁵ Pr Praseodymium 59	¹⁷⁶ Nd Neodymium 60	¹⁷⁷ Pm Promethium 61	¹⁷⁸ Sm Samarium 62	¹⁷⁹ Eu Europium 63	¹⁸⁰ Gd Gadolinium 64	¹⁸¹ Tb Terbium 65	¹⁸² Dy Dysprosium 66	¹⁸³ Ho Holmium 67	¹⁸⁴ Er Erbium 68	¹⁸⁵ Tm Thulium 69	¹⁸⁶ Yb Ytterbium 70	¹⁸⁷ Lu Lutetium 71	¹⁸⁸ Hf Hafnium 72	¹⁸⁹ Ta Tantalum 73	¹⁹⁰ W Tungsten 74	¹⁹¹ Re Rhenium 75	¹⁹² Os Osmium 76	¹⁹³ Ir Iridium 77	¹⁹⁴ Pt Platinum 78	¹⁹⁵ Au Gold 79	¹⁹⁶ Hg Mercury 80	¹⁹⁷ Tl Thallium 81	¹⁹⁸ Pb Lead 82	¹⁹⁹ Bi Bismuth 83	²⁰⁰ Po Polonium 84	²⁰¹ At Astatine 85	²⁰² Rn Radon 86
7	¹¹⁹ Fr Francium 87	¹²⁰ Ra Radium 88							²²⁷ La Lanthanum 57	²²⁸ Ce Cerium 58	²²⁹ Pr Praseodymium 59	²³⁰ Nd Neodymium 60	²³¹ Pm Promethium 61	²³² Sm Samarium 62	²³³ Eu Europium 63	²³⁴ Gd Gadolinium 64	²³⁵ Tb Terbium 65	²³⁶ Dy Dysprosium 66	²³⁷ Ho Holmium 67	²³⁸ Er Erbium 68	²³⁹ Tm Thulium 69	²⁴⁰ Yb Ytterbium 70	²⁴¹ Lu Lutetium 71	²⁴² Hf Hafnium 72	²⁴³ Ta Tantalum 73	²⁴⁴ W Tungsten 74	²⁴⁵ Re Rhenium 75	²⁴⁶ Os Osmium 76	²⁴⁷ Ir Iridium 77	²⁴⁸ Pt Platinum 78	²⁴⁹ Au Gold 79	²⁵⁰ Hg Mercury 80	²⁵¹ Tl Thallium 81	²⁵² Pb Lead 82	²⁵³ Bi Bismuth 83	²⁵⁴ Po Polonium 84	²⁵⁵ At Astatine 85	²⁵⁶ Rn Radon 86

Key

Relative atomic mass
Symbol
Name
Atomic number



BLANK PAGE



Answer ALL questions.

1 The table shows the numbers of particles in two atoms, L and M.

	Atom L	Atom M
number of electrons	6	6
number of neutrons	8	6
number of protons	6	6

(a) Which particles are present in the nuclei of both atoms? (1)

- A** electrons and neutrons
- B** electrons and protons
- C** neutrons and protons
- D** neutrons, protons and electrons

(b) (i) The atomic number of atom L is (1)

(ii) The mass number of atom L is (1)

(c) Atoms L and M are neutral because (1)

- A** the numbers of electrons and neutrons are equal
- B** the numbers of electrons and protons are equal
- C** the numbers of neutrons and protons are equal
- D** the numbers of electrons, neutrons and protons are equal



(d) Use information from the table to explain why atoms L and M are isotopes of the same element.

(2)

.....

.....

.....

.....

(e) The electronic configuration of atom M is

(1)

- A 2.2.2
- B 2.4
- C 2.4.6
- D 4.2

(Total for Question 1 = 7 marks)




2 Bromine, chlorine, fluorine and iodine are elements in Group 7 of the Periodic Table.

(a) Which two of these elements have the darkest colours?

(1)

..... and

(b) The equation for the reaction between hydrogen and chlorine is



Different names are used for the product, depending on its state symbol.

(i) What are the names used for HCl(g) and HCl(aq)?

(2)

HCl(g)

HCl(aq)

(ii) The presence of HCl(g) can be confirmed by adding ammonia (NH₃) gas.

State the observation in the reaction between HCl(g) and ammonia gas and write a chemical equation for the reaction.

(2)

observation

.....

chemical equation

(iii) The presence of chloride ions in HCl(aq) can be shown by mixing it with silver nitrate solution and dilute nitric acid.

State the result of this test and complete the chemical equation for the reaction by adding the state symbols.

(3)

result

.....



(c) Solution X is made by dissolving HCl(g) in water.

Solution Y is made by dissolving HCl(g) in methylbenzene.

A student added magnesium ribbon and blue litmus paper to separate samples of each solution.

The table shows her results.

Test	Solution X	Solution Y
magnesium ribbon added	bubbles	no change
blue litmus paper added	goes red	stays blue

(i) What substance is responsible for the bubbles? (1)

(ii) State one change to the magnesium ribbon that could be seen after adding it to solution X. (1)

(iii) What does the colour change of the litmus paper show about solution X? (1)

(iv) Why does the litmus paper stay blue in solution Y? (1)

(Total for Question 2 = 12 marks)




3 Tungsten is a useful metal. It has the chemical symbol W.

(a) One method of extracting tungsten involves heating a tungsten compound (WO_3) with hydrogen.

(i) Suggest the chemical name of WO_3 (1)

(ii) Balance the equation for the reaction between WO_3 and hydrogen. (1)



(iii) Why is this reaction described as reduction? (1)

(b) Scheelite is an ore of tungsten.

The main compound in scheelite has the percentage composition by mass
Ca = 13.9%, W = 63.9%, O = 22.2%.


Calculate the empirical formula of this compound. (3)

empirical formula =



(c) Tungsten can also be obtained by reacting tungsten fluoride with hydrogen.

The equation for this reaction is



(i) In an experiment, a chemist used 59.6 g of tungsten fluoride.

What is the maximum mass of tungsten he could obtain from 59.6 g of tungsten fluoride?

Relative formula mass of tungsten fluoride = 298

(2)

maximum mass = g

(ii) Starting with a different mass of tungsten fluoride, he calculates that the mass of tungsten formed should be 52.0 g. In his experiment he actually obtains 47.5 g of tungsten.

What is the percentage yield of tungsten in this experiment?

(2)


percentage yield = %

(Total for Question 3 = 10 marks)



4 A student investigated the neutralisation of acids by measuring the temperature changes when alkalis were added to acids of known concentrations.

He used this apparatus to add different volumes of sodium hydroxide solution to a fixed volume of dilute nitric acid.



He used this method.

- measure the temperature of 25.0 cm^3 of the acid in the polystyrene cup
- add the sodium hydroxide solution in 5.0 cm^3 portions until a total of 30.0 cm^3 has been added

(a) State two properties of the sodium hydroxide solution that should be kept constant for each 5.0 cm^3 portion.


(2)

1


2



(b) The diagram shows the thermometer readings in one experiment.



before adding alkali



after adding alkali

Write down the thermometer readings and calculate the temperature change.

(3)

temperature after adding alkali°C

temperature before adding alkali°C

temperature change°C



(c) The student carried out the experiment three times.

The table shows his results.

Volume of alkali added in cm ³	Temperature in °C		
	experiment 1	experiment 2	experiment 3
0.0	17.4	16.6	15.9
5.0	18.5	21.0	18.0
10.0	19.6	24.5	20.0
15.0	20.5	23.6	22.2
20.0	21.4	22.7	23.6
25.0	22.5	21.4	22.8
30.0	23.4	20.5	22.0

The teacher said that only the results for experiment 3 showed the expected increase and decrease in temperature.

(i) Why was there no temperature decrease in experiment 1?

(1)

- A The alkali was added too quickly
- B The starting temperature of the acid was too high
- C The acid concentration was half what it should have been
- D The volume of acid used was 50.0 cm³ instead of 25.0 cm³

(ii) Why were the temperature increases in experiment 2 much greater than expected?

(1)


- A The starting temperature of the acid was too high
- B The acid concentration was double what it should have been
- C The volume of acid used was 50.0 cm³ instead of 25.0 cm³
- D The alkali was added in 10.0 cm³ portions but were recorded as 5.0 cm³ portions



(d) Plot the results of experiment 3 on the grid.

Draw a straight line of best fit through the first four points, and another straight line of best fit through the last three points. Make sure that the two lines cross.

(4)



(e) The point where the lines cross indicates the volume of alkali added to exactly neutralise the acid and also the maximum temperature reached.

Record these values.

(2)

volume of alkali..... cm³


maximum temperature..... °C



(f) Another student used sulfuric acid instead of nitric acid in her experiments. She started with 25.0 cm³ of sulfuric acid of concentration 0.650 mol/dm³.

She added 0.500 mol/dm³ sodium hydroxide solution until the acid was completely neutralised.

The equation for this reaction is



(i) Calculate the amount, in moles, of sulfuric acid used.

(2)

amount = mol

(ii) Calculate the amount, in moles, of sodium hydroxide needed to neutralise this amount of sulfuric acid.

(1)

amount = mol

(iii) Calculate the volume, in cm³, of sodium hydroxide solution needed to neutralise this amount of sulfuric acid.


(2)

volume = cm³

(Total for Question 4 = 18 marks)



5 The diagram shows some important conversion processes used in the oil industry.



(a) Process 1 is called

(1)

- A catalytic cracking
- B condensation polymerisation
- C fractional distillation
- D thermal decomposition

(b) Describe the differences between crude oil and kerosene. In your answer you should refer to

- the average size of the molecules in the two liquids
- the covalent bonding in the molecules
- the viscosities of the two liquids

(3)

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....



(c) The equation for one reaction that could occur in process 2 is



(i) Deduce the formula of C_xH_y (1)

(ii) Give the name of the compound C_5H_{12} (1)

(iii) Draw the displayed formula of C_2H_4 (1)

(d) The structural formula of chloroethene formed in process 3 is $CH_2=CHCl$

The polymer formed in process 4 is poly(chloroethene).

Draw the **displayed** formula for the repeat unit of poly(chloroethene). (2)



(e) Poly(chloroethene) is formed by addition polymerisation.

Nylon is formed by condensation polymerisation.

(i) How does condensation polymerisation differ from addition polymerisation?

(1)

.....

.....

(ii) Poly(chloroethene) and nylon do not biodegrade easily.

What is meant by the term **biodegrade**?

(2)

.....

.....

.....

.....

(iii) What feature of addition polymers makes it difficult for them to biodegrade?

(1)

.....

.....

(Total for Question 5 = 13 marks)

(TOTAL FOR PAPER = 60 MARKS)



BLANK PAGE



BLANK PAGE



BLANK PAGE

