



Pearson
Edexcel

Mark Scheme (Results)

January 2021

Pearson Edexcel International GCSE
In Chemistry (4CH1) Paper 1CR and Science
(Double Award) (4SD0) Paper 1CR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2021

Publications Code 4CH1_1CR_2101_MS

All the material in this publication is copyright

© Pearson Education Ltd 2021

General Marking Guidance


- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1	(a) copper		1
	(b) glucose/water		1
	(c) air		1
	(d) nitrogen/oxygen		1
	(e) copper		1
	(f) oxygen and sulfur	in either order	1

(Total for Question 1 = 6)

Question number	Answer	Notes	Marks
2 (a)	Any two from the following: M1 contains 3 dyes M2 contains (dye) A M3 contains (dye) B M4 does not contain (dye) C / contains an unknown dye OWTTE		2
(b) (i)	(Ink 2) is insoluble (in solvent/water)	ALLOW does not contain (dye) A/B/C	1
(b) (ii)	(repeat) using different solvent	ALLOW named alternative solvent eg alcohol/ethanol	1
(c)	M1 correct measurement of distance moved by spot AND correct measurement of distance moved by solvent M2 use and evaluation of $R_f = \frac{\text{distance moved by spot}}{\text{distance moved by solvent front}}$ M3 answer to 2 sig fig	ALLOW 5.4-5.6 ALLOW 7.9-8.1 Expected: $\frac{5.5}{8.0} = 0.6875$ $= 0.69$ ALLOW ECF from M1 M2 correct answer with no working scores 3	3

(Total for Question 2 = 7)

Question number	Answer	Notes	Marks
3 (a)	alkanes		1
(b) (i)	<p>A boiling point is the correct answer because fractional distillation depends on differences in boiling point</p> <p>B is not correct because fractional distillation does not depend on differences in density</p> <p>C is not correct because fractional distillation does not depend on differences in melting point</p> <p>D is not correct because fractional distillation does not depend on differences in solubility</p>		1
(b) (ii)	 <p>1 mark for each correct line from boxes on left</p>	<p>If more than one line from a box on left do not award mark for that box</p>	3
(c)	<p>explanation including the following points:</p> <p>M1 (common impurity in fuels is) sulfur</p> <p>M2 sulfur burns/combusts/reacts (in air/oxygen) to form sulfur dioxide/SO₂</p> <p>M3 sulfur dioxide/SO₂ dissolves in/reacts with rain/water to form acid rain</p>	<p>If M2 M3 not scored ALLOW 1 mark for reference to sulphur dioxide/SO₂ and acid rain</p>	3

Question number	Answer	Notes	Marks
3 (d) (i)	cracking		1
(ii)	M1 (catalyst) silica/alumina M2 (temperature) 600-700 (°C)	ALLOW silicon dioxide/aluminium oxide ALLOW formulae ALLOW zeolite	2
(iii)	$C_{13}H_{28} \rightarrow C_8H_{18} +$ M1 $C_3H_6 +$ M2 C_2H_4	in either order ALLOW structural formulae ALLOW 1 mark for single product C_5H_{10}	2

(Total for Question 3 = 13)

Question number	Answer	Notes	Marks							
4 (a)	(i)	(hydrated) iron(III) oxide	REJECT incorrect oxidation states	1						
	(ii)	M1 (barrier method involves) coating iron in paint/oil/grease/plastic	ALLOW coating in named metal below iron in reactivity series eg tin	2						
		M2 stops oxygen/air/water getting to the iron	ALLOW stops iron reacting with oxygen/air/water							
	(b) (i)	<table border="1"> <tr> <td>reading at start</td> <td>20.5</td> </tr> <tr> <td>reading at end</td> <td>33.5 (1)</td> </tr> <tr> <td>volume of oxygen used in cm³</td> <td>13.0 (1)</td> </tr> </table>		reading at start	20.5	reading at end	33.5 (1)	volume of oxygen used in cm ³	13.0 (1)	CSQ on reading
reading at start		20.5								
reading at end		33.5 (1)								
volume of oxygen used in cm ³	13.0 (1)									
(ii)	not all oxygen had reacted / not enough iron (wool)	ALLOW not left for long enough/ OWTTE	1							
(c)	<p>M1 calculation of volume oxygen used</p> <p>M2 correct expression for percentage of oxygen</p> <p>M3 correct evaluation</p> <p>Example calculation:</p> <p>M1 (35.5 - 20.0 =) 15.5</p> <p>M2 (15.5 ÷ 80.0) x 100</p> <p>M3 19.4(%)</p>	<p>ALLOW ECF from M1</p> <p>ACCEPT 19.375/19.38</p> <p>correct answer with no working scores 3</p>	3							

(Total for Question 4 = 9)

Question number	Answer	Notes	Marks
5 (a)	(thermal) decomposition (1)		1
(b)	any two of the following: M1 (use the same) amount of metal carbonate M2 (use the same) sized pieces/surface area M3 (use the same) volume of limewater M4 (use same) size flame / distance of flame from boiling tube OWTTE	ALLOW mass ALLOW amount	2
(c)	bubbles are air (from tube) / caused by air (expanding on heating)	ALLOW gas in tube expands (on heating)	1
(d)	explanation including M1 (when limewater turns milky/cloudy it) shows carbon dioxide produced M2 showing metal carbonate has reacted/decomposed	ALLOW carbon dioxide comes from carbonate (reacting/decomposing)	2
(e) (i)	M1 (from) green M2 (to) black	IGNORE qualifiers eg light	2
(ii)	$\text{CuCO}_3 \rightarrow \text{CuO} + \text{CO}_2$	ALLOW products in either order	1

Question number	Answer	Notes	Marks
5 (f) (i)	<p>M1 the lower the metal is (in the reactivity series)</p> <p>M2 the more easily the (metal) carbonate reacts/decomposes</p>	<p>ALLOW the less reactive a metal is</p> <p>ALLOW the more easily the (metal) carbonate produces carbon dioxide</p> <p>ALLOW references to the less time the (metal) carbonate takes to react/decompose</p> <p>ALLOW references to the faster the (metal) carbonate reacts/decomposes</p> <p>ACCEPT reverse arguments</p>	2
(ii)	repeat (the investigation) using different / other / more (metal) carbonates		1

(Total for Question 5 = 12)

Question number	Answer	Notes	Marks
6 (a) (i)	Zn + 2HCl → ZnCl ₂ + H ₂ M1 all symbols and formulae correct M2 correctly balanced	M2 DEP M1	2
(ii)	lighted splint (produces squeaky) pop		1
(b) (i)	52 (cm ³)	ALLOW 51.5-52.5	1
(ii)	M1 vertical line from 15 cm ³ acid to graph line M2 volume hydrogen from graph multiplied by 2 OR M1 vertical line from 30 cm ³ acid to graph line M2 volume hydrogen from graph	ALLOW extra point drawn at 15 cm ³ ALLOW 68-70 ALLOW extra point drawn at 30 cm ³ ALLOW 68-70	2
(c)	explanation linking M1 more (acid) particles/(hydrogen) ions/H ⁺ in same volume M2 more (successful) collisions per second/unit time M3 rate increases	REJECT molecules once only ALLOW particles are closer together ACCEPT more frequent collisions IGNORE more chance/probability of collision ALLOW reaction is faster /speeds up MAX 1 if refer to particles moving faster/having more energy	3




6 (d)	<p>explanation including</p> <p>M1 (increase/decrease) surface area</p> <p>M2 increase surface area by using smaller pieces of zinc</p> <p>M3 more (successful) collisions per second/unit time (so rate increases)</p>	<p>ACCEPT more frequent collisions</p> <p>IGNORE more chance/probability of collision</p> <p>M2 M3 ACCEPT reverse arguments</p>	3
-------	---	---	---


(Total for Question 6 = 12)

Question number	Answer	Notes	Marks
7 (a)	<p>description including</p> <p>(formation of ions in lithium chloride involves)</p> <p>M1 lithium (atom) losing electron</p> <p>M2 chlorine (atom) gaining an electron</p> <p>(formation of covalent bonds in hydrogen chloride involves)</p> <p>M3 sharing a pair of electrons (one electron from each atom)</p>	<p>ALLOW correct dot-and-cross diagrams for ions for M1 and M2</p> <p>ALLOW correct dot-and-cross diagram showing shared pair in hydrogen chloride for M3</p>	3

<p>7 (b)</p>	<p>explanation including five of the following points:</p> <p>(lithium chloride)</p> <p>M1 giant (ionic) structure</p> <p>M2 strong (electrostatic) forces of attraction</p> <p>M3 between oppositely charged ions</p> <p>(hydrogen chloride)</p> <p>M4 simple molecular structure</p> <p>M5 weak intermolecular forces of attraction</p> <p>M6 more (heat/thermal) energy needed to overcome forces/break bonds in lithium chloride (than intermolecular forces in hydrogen chloride)</p> <p>OWTTE</p>	<p>ALLOW giant lattice</p> <p>ALLOW strong bonds</p> <p>ACCEPT positive and negative ions</p> <p>If any reference to molecules/atoms/covalent bonds/intermolecular forces/metallic bonds cannot score M2 M3 M6</p> <p>ALLOW weak bonds between molecules</p> <p>ACCEPT reverse argument</p> <p>If description/implication of breaking covalent bonds in hydrogen chloride cannot score M5 M6</p>	<p>5</p>
--------------	--	---	----------

(Total for Question 7 = 8)

Question number	Answer	Notes	Marks
8 (a) (i)	<p>explanation linking</p> <p>M1 (molecules/compounds) having same molecular formulae</p> <p>M2 but different structural/displayed formulae</p>	<p>ALLOW different structures</p> <p>ALLOW different arrangement of atoms</p>	2
(ii)	<p>M1 displayed formula of but-1-ene</p>  <p>M2 displayed formula of but-2-ene</p> 	<p>IGNORE bond angles</p> <p>ALLOW displayed formula of methylpropene</p> 	2

Question number	Answer	Notes	Marks
8 (b) (i)	<p>explanation linking</p> <p>M1 molecule is unsaturated as contains (carbon to carbon) double bond</p> <p>M2 molecule not a hydrocarbon as contains oxygen</p>	ACCEPT does not contain hydrogen and carbon only	2
(ii)	addition		
(iii)	<div style="text-align: center;">  </div> <p>M1 correct repeat unit structure</p> <p>M2 extension bonds, brackets and n</p>	n can be anywhere after bracket extension bonds do not have to go through brackets M2 DEP M1 or near miss	2

Question number	Answer	Notes	Marks
8 (c) (i)	<p>M1 calculation of mass of octane</p> <p>M2 calculation of Mr of C₈H₁₈ and CO₂</p> <p>M3 link between mass/mol C₈H₁₈ and mass/mol CO₂</p> <p>M4 calculation of mass CO₂ in g</p> <p>M5 calculation of mass CO₂ in kg</p> <p>Example calculation:</p> <p>M1 50 x 700 = 35000 (g)</p> <p>M2 M_r of C₈H₁₈ = 114 and M_r CO₂ = 44</p> <p>M3 114g C₈H₁₈ produces (8 x 44 =) 352 (g) CO₂</p> <p>M4 35000 g C₈H₁₈ produces $\frac{35000 \times 352}{114}$ = 108070.2 (g) CO₂</p> <p>M5 = 108 (kg)</p> <p>Alternative method using mol:</p> <p>M1 50 x 700 = 35000 (g)</p> <p>M2 M_r of C₈H₁₈ = 114 and M_r CO₂ = 44</p> <p>M3 n (C₈H₁₈) = $\frac{35000}{114}$ = 307.0175 mol so n (CO₂) = 8 x 307.0175 = 2456.14 mol</p> <p>M4 mass CO₂ = 2456.14 x 44 = 108070.2 (g)</p> <p>M5 = 108 (kg)</p>	<p>ALLOW 2 sig fig or more in M4 and M5</p> <p>ALLOW 2 sig fig or more in M3 M4 and M5</p> <p>MARK CSQ in both methods</p> <p>correct answer without working scores 5</p>	5

(ii)	global warming/climate change	IGNORE greenhouse effect	1
------	-------------------------------	--------------------------	---

(Total for Question 8 = 15)

Question number	Answer	Notes	Marks
9 (a)	to keep out of contact/prevent reaction with air/oxygen/water/moisture	ALLOW they react with air/oxygen/water/moisture	1
(b) (i)	similarity: both fizz/move on surface/ produce flame difference caesium: faster/more violent reaction	ALLOW both melt/form a ball/produce a gas/ produce hydrogen/form an alkaline solution ACCEPT reverse argument for potassium ALLOW caesium explodes	2
(ii)	$2\text{Cs} + 2\text{H}_2\text{O} \rightarrow 2\text{CsOH} + \text{H}_2$ M1 all symbols/formulae correct M2 correctly balanced	ACCEPT fractions and multiples M2 DEP M1	2
(c) (i)	a lid/cover		1
(ii)	explanation linking either M1 stir solution M2 to obtain more accurate (maximum) temperature OR M1 measure temperature of sodium hydroxide M2 to check if different to/same as temperature of (hydrochloric) acid	ALLOW reference to even temperature throughout/ heat evenly distributed OWTTE ALLOW take an average of temperature of sodium hydroxide and temperature of (hydrochloric) acid for 2 marks	2

Question number	Answer	Notes	Marks
9 (d) (i)	<p>M1 correct temperature change/ΔT</p> <p>M2 correct substitution into $Q = m \times c \times \Delta T$</p> <p>M3 correct evaluation</p> <p>Example calculation:</p> <p>M1 $\Delta T = (26.5 - 19.9)$ OR 6.6</p> <p>M2 $Q = 100 \times 4.2 \times 6.6$</p> <p>M3 = 2800 (J)</p>	<p>M2 ECF M1</p> <p>M3 ECF M2</p> <p>IGNORE any sign ALLOW 2770, 2772</p> <p>correct answer without working scores 3</p>	3
(ii)	<p>M1 answer to M3 from (i) $\div 0.05$</p> <p>M2 correct evaluation in kJ/mol with negative sign</p> <p>expected answer</p> <p>M1 $2800 \div 0.05$ OR 56000 (J)</p> <p>M2 - 56 (kJ/mol)</p>	<p>2770 gives 55400 2772 gives 55440</p> <p>negative sign required</p> <p>ACCEPT -55.4 ACCEPT -55.44 ACCEPT -55</p> <p>correct answer without working scores 2</p>	2

(Total for Question 9 = 13)

Question number	Answer	Notes	Marks
10 (a) (i)	neutralisation	ALLOW acid - base	1
(ii)	acid donates proton(s)/base accepts proton(s)	ALLOW metal oxide for base	1
(b) (i)	<p>description including</p> <p>M1 appropriate use of at least three named pieces of apparatus</p> <p>AND any four of the following points</p> <p>M2 add copper(II) carbonate to (dilute sulfuric acid (a spatula/little at a time and stir after each addition)</p> <p>M3 until no more effervescence</p> <p>M4 filter (to remove excess copper(II) carbonate/to obtain (copper(II) sulfate) solution)</p> <p>M5 heat/warm filtrate/(copper(II) sulfate) solution until crystals start to appear (solution saturated) OWTTE</p> <p>M6 filter to obtain (the saturated) solution</p>	<p>ALLOW until no more reacts/dissolves</p> <p>ALLOW until in excess</p> <p>IGNORE if continue and prepare crystals instead of saturated solution</p>	5
(b) (ii)	<p>M1 calculation of actual mass of crystals obtained</p> <p>M2 division by expected mass of crystals (6.4) and multiplication by 100 to convert to percentage</p> <p>M3 correct to 1 dp</p> <p>Example calculation</p> <p>M1 (6.40 - 1.80 =) 4.6(0)</p> <p>M2 (% yield =) $\frac{4.6}{6.4} \times 100$ OR 71.875 (%)</p> <p>M3 = 71.9 (%)</p>	<p>M2 ECF M1</p> <p>M3 DEP M2</p>	3

Question number	Answer	Notes	Marks
10 (c) (i)	<p>M1 find percentage of water</p> <p>M2 divide each percentage by Mr to find number of moles</p> <p>M3 divide each answer by smallest to find ratio and value of x</p> <p>Expected calculation:</p> <p>M1 (79%) CaSO₄ 21% H₂O</p> <p>M2 $\frac{79}{136}$ (= 0.58) $\frac{21}{18}$ (= 1.17)</p> <p>M3 $\frac{0.58}{0.58}$ $\frac{1.17}{0.58}$ = 1 : 2</p> <p>so x = 2</p>		3
(ii)	<p>description including</p> <p>M1 do a flame test</p> <p>M2 orange-red flame</p>	<p>correct answer without working scores 3</p> <p>ALLOW description of flame test</p>	2

(Total for Question 10 = 15)

Total for paper = 110 marks

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom