



Pearson

Mark Scheme (Results)

January 2018

Pearson Edexcel International GCSE in
Further Pure Mathematics (4PM0)
Paper 02

edexcel 

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2018

Publications Code 4PM0_02_1801_MS

All the material in this publication is copyright

© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.
Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
 - M marks: method marks
 - A marks: accuracy marks
 - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
 - cao – correct answer only
 - ft – follow through
 - isw – ignore subsequent working
 - SC - special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - eeoo – each error or omission
- **No working**

If no working is shown then correct answers may score full marks
If no working is shown then incorrect (even though nearly correct) answers score no marks.
- **With working**

Always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.
If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks.
Any case of suspected misread loses 2A (or B) marks on that part, but can gain the M marks.
If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.

Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

- **Linear equations**

Full marks can be gained if the solution alone is given, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

General Principles for Further Pure Mathematics Marking

(but note that specific mark schemes may sometimes override these general principles)

Method mark for solving a 3 term quadratic equation:

1. Factorisation:

$$(x^2 + bx + c) = (x + p)(x + q), \text{ where } |pq| = |c| \text{ leading to } x = \dots$$

$$(ax^2 + bx + c) = (mx + p)(nx + q) \text{ where } |pq| = |c| \text{ and } |mn| = |a| \text{ leading to } x = \dots$$

2. Formula:

Attempt to use the **correct** formula (shown explicitly or implied by working) with values for a , b and c , leading to $x = \dots$

3. Completing the square:

$$x^2 + bx + c = 0: \left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0, \quad q \neq 0 \quad \text{leading to } x = \dots$$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \rightarrow x^{n-1}$)

2. Integration:

Power of at least one term increased by 1. ($x^n \rightarrow x^{n+1}$)

Use of a formula:

Generally, the method mark is gained by **either**

quoting a correct formula and attempting to use it, even if there are mistakes in the substitution of values

or, where the formula is not quoted, the method mark can be gained by implication from the substitution of correct values and then proceeding to a solution.

Answers without working:

The rubric states "Without sufficient working, correct answers may be awarded no marks".

General policy is that if it could be done "in your head" detailed working would not be required.

(Mark schemes may override this eg in a case of "prove or show....")

Exact answers:

When a question demands an exact answer, all the working must also be exact. Once a candidate loses exactness by resorting to decimals the exactness cannot be regained.

Rounding answers (where accuracy is specified in the question)

Penalise only once per question for failing to round as instructed - ie giving more digits in the answers. Answers with fewer digits are automatically incorrect, but the isw rule may allow the mark to be awarded before the final answer is given.

Question Number	Scheme	Marks
<p>1 (a)</p> <p>(b)</p> <p>ALT</p>	<p>$192 = \frac{\theta}{2} \times 12^2 \Rightarrow \theta = \frac{8}{3}$ (radians) accept 2.67 or better</p> <p>$l = 12 \times \frac{8}{3} = 32$ (cm)</p> <p>Use $A = \frac{1}{2}rl$ $192 = 6l \Rightarrow l = 32$ M1A1</p>	<p>M1A1 (2)</p> <p>M1A1 (2)</p> <p>[4]</p>
<p>(a)</p> <p>M1</p> <p>A1</p> <p>(b)</p> <p>M1</p> <p>A1</p>	<p>Use of correct formula. If the formula for use with degrees is used, must change to radians for the answer to gain this mark</p> <p>Correct answer</p> <p>Use of either correct (radian) formula with their angle from (a) or the degree formula</p> <p>Correct answer awrt 32</p>	

Question Number	Scheme	Marks
<p>2(a)</p> <p>ALT</p> <p>(b)</p> <p>ALT</p>	$\sum_{r=1}^n (3r+2) = \frac{n}{2}(2 \times 5 + (n-1)3) = \frac{n}{2}(7+3n) \quad *$ <p>Splitting terms:</p> $\sum_{r=1}^n (3r+2) = \sum_{r=1}^n 3r + \sum_{r=1}^n 2 = 3 \times \frac{n(n+1)}{2} + 2n = \frac{n}{2}(7+3n) \quad *$ $\sum_{r=10}^{20} (3r+2) = \sum_{r=1}^{20} (3r+2) - \sum_{r=1}^9 (3r+2)$ $= \frac{20}{2}(7+3 \times 20) - \frac{9}{2}(7+3 \times 9) = 517$ $\sum_{r=10}^{20} (3r+2) = \frac{11}{2}(32+62) = 517 \quad \text{M1A1A1}$	<p>M1A1cso (2)</p> <p>M1,A1 (2)</p> <p>M1</p> <p>A1A1 (3)</p>
<p>(a)</p> <p>M1</p> <p>A1cso</p> <p>ALT:</p> <p>M1</p> <p>A1cso</p> <p>(b)</p> <p>M1</p> <p>A1</p> <p>A1cao</p> <p>ALT</p> <p>M1</p> <p>A1</p> <p>A1cao</p>	<p>Use $S = \frac{n}{2}(2a + (n-1)d)$ or $\frac{n}{2}(a+l)$ showing the correct substitution</p> <p>Reach given result with no errors seen</p> <p>Split the sum into 2 parts and use either sum formula on $\sum_{r=1}^n 3r$ or use the standard result for the sum of the first n natural numbers. Allow if $\sum_{r=1}^n 2$ or $2n$ seen</p> <p>Reach given result with no errors seen</p> <p>Use the difference of two sums with upper limit 9 or 10 for second sum</p> <p>Substitute correct numbers ($n = 9$ now)</p> <p>517</p> <p>Use summation formula with $n = 11$ or $10, a = 32, l = 62$</p> <p>Substitute correct numbers ($n = 11$ now)</p> <p>517</p> <p>NB: (b) can be done by listing the terms and adding them $32 + 35 + \dots + 62$ with an answer seen is minimum for M1 Ignore any intermediate terms if shown. A2 correct answer</p> <p>Correct answer with no working shown scores 0/3</p>	<p>[5]</p>

Question Number	Scheme	Marks
<p>3</p> <p>ALT</p>	<p>$x = 1, y^2 = 4 \Rightarrow y = \pm 2$</p> <p>Volume = $\pi \int_{-2}^2 (5 - y^2) dy - \pi \int_{-2}^2 1 dy, = (\pi) \left(\left[5y - \frac{y^3}{3} \right]_{-2}^2 - [y]_{-2}^2 \right)$</p> <p>$= (\pi) \left\{ \left(10 - \frac{8}{3} \right) - \left(-10 - \frac{-8}{3} \right) - (2 - -2) \right\} = \frac{32\pi}{3} \text{ (units}^3\text{)}$</p> <p>B1 Limits as above</p> <p>Volume = $\pi \int_{-2}^2 (5 - y^2) dy - \pi \times 1 \times 4 = \pi \left[5y - \frac{y^3}{3} \right]_{-2}^2 - 4\pi$ M1M1</p> <p>$= \pi \left\{ \left(10 - \frac{8}{3} \right) - \left(-10 - \frac{-8}{3} \right) \right\} - 4\pi = \frac{32\pi}{3} \text{ (units}^3\text{)}$ M1A1</p>	<p>B1</p> <p>M1,M1</p> <p>dM1A1cao</p> <p>[5]</p>
<p>B1</p> <p>M1</p> <p>M1</p> <p>dM1</p> <p>A1cao</p>	<p>Notes cover either method</p> <p>Correct y coords for points of intersection (shown explicitly or only seen as limits)</p> <p>Use $\pi \int x^2 dy$ for volume of curve, with an attempt to obtain an integrand in terms of y, and cylinder by integral or standard volume formula. This mark can only be awarded when evidence of a difference of these volumes is seen. Limits not needed.</p> <p>Attempt the integration of their dimensionally correct curve integral. (ie not squared). Integration must be wrt y. Limits and π not needed.</p> <p>Algebraic integration must be seen.</p> <p>Substitute their limits in their integrated function and obtain a value for the volume of the cylinder using consistent values. π not needed. Depends on 2nd M mark (but not first)</p> <p>Correct volume (as shown or equivalent multiple of π eg 10.7π)</p> <p>NB: All marks are available if work is done without π but π included in the final answer.</p>	

Question Number	Scheme	Marks
<p>4(a)</p> <p>(b)</p>	<p>$b^2 - 4ac > 0$ $p^2 - 4 \times 3 \times 4 > 0$</p> <p>$\Rightarrow p^2 > 48 \Rightarrow$ critical values are $p = \pm\sqrt{48}$ ($= \pm 4\sqrt{3}$)</p> <p>So set of values; $p < -4\sqrt{3}, p > 4\sqrt{3}$ (accept 3dp or better inc $\pm\sqrt{48}$)</p> <p>$\pm 6, \pm 5, \pm 4, \pm 3, \pm 2, \pm 1, 0$</p>	<p>M1</p> <p>dM1A1</p> <p>ddM1A1 (5)</p> <p>B1 (1)</p> <p>[6]</p>
<p>(a)</p> <p>M1</p> <p>dM1</p> <p>A1</p> <p>dM1</p> <p>A1</p> <p>(b)</p> <p>B1</p>	<p>For the first 3 marks accept an equation or any inequality sign. For the first 4 marks accept the use of x instead of p</p> <p>Use discriminant</p> <p>Solve to find the CVs Depends on the first M mark.</p> <p>Correct CVs, exact or (min) 3 dp</p> <p>Form 2 inequalities for the outside regions using their CVs ie $p <$ smaller CV and $p >$ larger CV Depends on both previous M marks</p> <p>Correct set of values</p> <p>Answer as shown.</p>	

Question Number	Scheme	Marks
<p>5</p>	$\frac{dy}{dx} = 2e^x(3x^2 - 6) + 12xe^x$ $\frac{d^2y}{dx^2} = [2e^x(3x^2 - 6) + 12xe^x] + [12xe^x + 12e^x]$ $\frac{d^2y}{dx^2} = e^x(6x^2 + 24x)$ $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = 12e^x \quad *$ <p>ALT First derivative as above</p> $\frac{dy}{dx} = y + 12xe^x$ $\frac{d^2y}{dx^2} = \frac{dy}{dx} + 12xe^x + 12e^x$ <p>Substitute both derivatives in the given result</p> <p>Correct answer as given, no errors seen</p>	<p>M1A1A1</p> <p>M1A1</p> <p>dM1A1 (7) [7]</p> <p>M1A1A1</p> <p>M1A1</p> <p>dM1</p> <p>A1</p>
<p>M1 A1 A1 M1 A1 dM1 A1</p>	<p>Differentiate using the product rule. The sum of 2 terms reqd.</p> <p>Either term correct. No simplification needed. Ignore errors made when simplifying.</p> <p>Both terms correct. No simplification needed. Ignore errors made when simplifying.</p> <p>Differentiate again using the product rule <i>correctly</i> on at least one term. (The term the product rule is applied to need not be correct.)</p> <p>Any correct result for $\frac{d^2y}{dx^2}$ (ie their second derivative should be equivalent to the one shown)</p> <p>Complete, probably by substituting their derivatives in LHS of the given result. At least one intermediate step (eg full substitution or bracketed equation above must be seen) Depends on both previous M marks.</p> <p>Fully correct final result reached with no errors seen.</p>	

Question Number	Scheme	Marks
6(a)	$\overrightarrow{AB} = -\mathbf{a} + \mathbf{b}$	B1 (1)
(b)	$\overrightarrow{PQ} = \frac{\mathbf{a}}{4} + \frac{1}{2}(-\mathbf{a} + \mathbf{b}), = \frac{1}{4}(2\mathbf{b} - \mathbf{a})$ oe	M1,A1 (2)
(c)(i)	$\overrightarrow{QR} = \frac{\mu}{4}(2\mathbf{b} - \mathbf{a})$	B1 ft
(ii)	$\overrightarrow{QR} = \frac{1}{2}(\mathbf{b} - \mathbf{a}) + \lambda\mathbf{b}$	M1A1 (3)
(d)(i)	$\frac{2\mu}{4}\mathbf{b} - \frac{\mu}{4}\mathbf{a} = \frac{1}{2}\mathbf{b} - \frac{1}{2}\mathbf{a} + \lambda\mathbf{b} \Rightarrow -\frac{\mu}{4} = -\frac{1}{2} \Rightarrow \mu = 2$	M1M1A1
(ii)	$\frac{2\mu}{4} = \frac{1}{2} + \lambda \Rightarrow \lambda = \frac{1}{2}$	A1 (4)
[10]		
(a)B1	Correct vector	
(b)M1	$\overrightarrow{PQ} = \frac{\mathbf{a}}{4} + \frac{1}{2}(\text{their } \overrightarrow{AB})$ or $\overrightarrow{PO} + \overrightarrow{OB} + \overrightarrow{BQ} = -\frac{3}{4}\mathbf{a} + \mathbf{b} - \frac{1}{2}(\text{their } \overrightarrow{AB})$	
A1	Correct vector \overrightarrow{PQ}	
(c)		
(i)B1ft	$\frac{\mu}{4}(\text{their } \overrightarrow{PQ})$	
(ii)M1	$\overrightarrow{QR} = \frac{1}{2}(\text{their } \overrightarrow{AB}) + \lambda\mathbf{b}$	
A1	Correct vector as shown or with \mathbf{b} terms collected	
(d)		
M1	Equate their 2 forms for \overrightarrow{QR}	
M1	Equate coeffs of \mathbf{a} and obtain a value for μ or equate coefficients of \mathbf{a} and \mathbf{b} and get a value of λ	
(i)A1	Both versions of \overrightarrow{QR} must have an \mathbf{a} term and a \mathbf{b} term	
(i)A1	Correct value of μ (or λ)	
(ii)A1	Complete to obtain the correct value of the second unknown.	

Question Number	Scheme	Marks
7(i)	$\frac{(8^x)^x}{32^x} = 4 \Rightarrow \frac{2^{3x^2}}{2^{5x}} = 2^2 \Rightarrow 2^{(3x^2-5x)} = 2^2$ $\Rightarrow 3x^2 - 5x = 2 \Rightarrow (3x+1)(x-2) = 0$ $x = -\frac{1}{3}, 2$	M1A1 M1 A1 (4)
(ii)	$\log_x 64 - \log_x 4 = \log_x \left(\frac{64}{4} \right) = \log_x 16$ $\log_x 16 = \frac{\log_4 16}{\log_4 x} = \frac{2}{\log_4 x}$ $3\log_4 x + \frac{2}{\log_4 x} = 5 \Rightarrow 3(\log_4 x)^2 + 2 = 5\log_4 x$ $\Rightarrow 3(\log_4 x)^2 - 5\log_4 x + 2 = 0$ $\Rightarrow (3\log_4 x - 2)(\log_4 x - 1) = 0$ $\Rightarrow \log_4 x = \frac{2}{3}, \log_4 x = 1$ $\Rightarrow x = 4^{\frac{2}{3}} \left(= 2^{\frac{4}{3}} = \sqrt[3]{16} \right) = 2.5198421... \approx 2.52 \text{ or better } x = 4^1 = 4$	M1 M1 M1 dM1 A1 dM1A1 (7) [11]
ALT (ii)	$\log_x 64 - \log_x 4 = \log_x \left(\frac{64}{4} \right) = \log_x 16 = 2\log_x 4$ $\log_4 x = \frac{\log_x x}{\log_x 4} = \frac{1}{\log_x 4}$ $2\log_x 4 + \frac{3}{\log_x 4} = 5 \Rightarrow 2(\log_x 4)^2 + 2 = 5\log_x 4$ $\Rightarrow 2(\log_x 4)^2 - 5\log_x 4 + 3 = 0$ $\Rightarrow (2\log_x 4 - 3)(\log_x 4 - 1) = 0$ $\Rightarrow \log_x 4 = \frac{3}{2}, \log_x 4 = 1 \Rightarrow 4 = x^{\frac{3}{2}}, 4 = x^1$ $\Rightarrow x = 4^{\frac{2}{3}} = \left(\sqrt[3]{16} \right) = 2.5198421... \approx 2.52 \text{ or better, } x = 4^1 = 4$	M1 M1 M1 dM1 A1 dM1A1 (7)

Question Number	Scheme	Marks
<p>(i)</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>Change all terms of equation to powers of 2 (or possibly 4)</p> <p>Correct 2 term equation with powers of 2</p> <p>Equate the powers in their equation and solve the resulting 3 term quadratic</p> <p>Correct values for x (both needed)</p> <p>Special Case: Using factor theorem: Substitute $x = 2$ and show correct M1A1M0A0</p> <p>If (unlikely) same done with $x = -\frac{1}{3}$ - send to Review!</p>	
<p>(ii)</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>dM1</p> <p>A1</p> <p>dM1</p> <p>A1</p> <p>ALT</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>dM1</p> <p>A1</p> <p>dM1</p> <p>A1</p>	<p>The work for the first 3 M marks may appear in a different order.</p> <p>Enter the marks in the order shown here.</p> <p>Combine the two logs base x or combine the equivalent logs after changing base. Award for combining the 2 equivalent numbers after multiplying through by their denominators</p> <p>Change all logs base x to logs base 4 (or all logs to the same base)</p> <p>Obtain a 3 term quadratic, terms in any order.</p> <p>Solve their 3 term quadratic to $\log_4 x = \dots$ or $\log_p x = \dots$ Depends on all previous M marks</p> <p>Two correct values for $\log_4 x$ or $\log_p x$</p> <p>"Undo" their logs to get at least one value for x (not nec correct) Depends on all previous M marks.</p> <p>Two correct values for x. Accept accurate answers or min 3 sf</p> <p>Combine the two logs base x</p> <p>Change all log base 4 to log base x</p> <p>Obtain a 3 term quadratic, terms in any order.</p> <p>Solve their 3 term quadratic to $\log_x 4 = \dots$ or $\log_x p = \dots$ Depends on all previous M marks</p> <p>Two correct values for $\log_x 4$ or $\log_x p$</p> <p>"Undo" their logs to get at least one value for x Depends on all previous M marks</p> <p>Two correct values for x</p>	


Question Number	Scheme	Marks
<p>8 (a)</p> <p>(b)</p> <p>(c)</p>	$355 = \pi r^2 h \Rightarrow h = \frac{355}{\pi r^2} \text{ or } \pi r h = \frac{355}{r}$ $S = 2\pi r^2 + 2\pi r h \Rightarrow S = 2\pi r^2 + 2\pi r \left(\frac{355}{\pi r^2} \right) = 2\pi r^2 + \frac{710}{r} \quad *$ $\frac{dS}{dr} = 4\pi r - 710r^{-2}$ $4\pi r - 710r^{-2} = 0 \Rightarrow 4\pi r = \frac{710}{r^2} \Rightarrow r^3 = \frac{710}{4\pi}$ $r = \sqrt[3]{\frac{710}{4\pi}} \quad (r = 3.837215\dots) \text{ cm}$ $S = 2\pi \times \left(\sqrt[3]{\frac{710}{4\pi}} \right)^2 + \frac{710}{\sqrt[3]{\frac{710}{4\pi}}} = 277.5450\dots \approx 278 \text{ (cm}^2\text{)}$ $\frac{d^2S}{dr^2} = 4\pi + \frac{1420}{r^3} \quad \left\{ = 4\pi + \frac{1420}{3.837215^3} = 37.699 \right\}$ <p>(r positive so) $\frac{d^2S}{dr^2} > 0 \therefore S$ is minimum</p>	<p>B1</p> <p>M1A1 A1cso (4)</p> <p>M1</p> <p>dM1</p> <p>A1</p> <p>dM1A1cao (5)</p> <p>M1</p> <p>A1ft (2)</p> <p>[11]</p>
<p>(a)</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1cso</p> <p>(b)</p> <p>M1</p> <p>dM1</p> <p>A1</p> <p>dM1</p> <p>A1cao</p> <p>(c)</p> <p>M1</p> <p>A1ft</p>	<p>$h = \frac{355}{\pi r^2} \text{ or } \pi r h = \frac{355}{r}$..seen explicitly</p> <p>Use a correct formula for the surface area and substitute their expression for h which must have been seen explicitly. (eg $S = 2\pi r^2 + 2\pi r h \Rightarrow S = 2\pi r^2 + \left(\frac{355 \times 2}{r} \right)$ alone scores M0 as does any other re-arrangement of the answer.)</p> <p>Correct expression for h used</p> <p>Obtain the given result from a fully correct solution. Must see $S = \dots$</p> <p>Differentiate the given expression for S - power of either term to decrease</p> <p>Equate their derivative to 0 and solve for r Depends on the first M mark</p> <p>Correct value for r, exact or decimal (3 sf sufficient) seen explicitly or used to calculate the minimum value of S.</p> <p>Substitute their value for r in the given expression for S Depends on the previous 2 M marks. 278</p> <p>Obtain the second derivative (or use an other method to test for a min value of S). Methods involving testing value of S on either side of their value of r or looking at the change of sign of the first derivative must include evaluating S or dS/dt</p> <p>Concluding (correct) statement. No need to evaluate the second derivative provided their value of r is positive and the second derivative is algebraically correct. (Ignore incorrect evaluation unless negative.)</p>	

Question Number	Scheme	Marks
<p>9(a)</p> <p>(i)</p> <p>(ii)</p>	$\frac{x^3 - 2x^2 - 5x + 6}{x + 2} = x^2 - 4x + 3$ $x^2 - 4x + 3 = (x - 3)(x - 1) \Rightarrow x = 1, 3$ <p>so $a = 1$ *</p> <p>$b = 3$</p>	<p>M1</p> <p>M1(NB A1 on e- PEN)</p> <p>A1cso A1 (B1 on e-PEN)</p> <p>(4)</p>
<p>(b)</p>	<p>Correct answers w/o working scores 0/4</p> $\frac{dy}{dx} = 3x^2 - 4x - 5 \quad \text{when } x = 2, \quad \frac{dy}{dx} = -1$ <p>$x = 2, y = -4$</p> $(y - -4) = -1(x - 2)$ <p>when $y = 0, x = -2$ *</p>	<p>M1A1</p> <p>B1</p> <p>M1A1</p> <p>A1cso (6)</p>
<p>ALT</p>	<p>$x = 2, y = -4$</p> $\frac{dy}{dx} = 3x^2 - 4x - 5 \quad \text{when } x = 2, \quad \frac{dy}{dx} = -1$ <p>Grad of line from P to $(-2, 0) = -1$</p> <p>Same gradient so l passes through $(-2, 0)$</p>	<p>B1</p> <p>M1A1</p> <p>M1A1</p> <p>A1cso</p>
<p>(c)</p>	$\text{Area} = \int_{-2}^2 (x^3 - 2x^2 - 5x + 6) dx - \int_{-2}^2 (-x - 2) dx = \int_{-2}^2 (x^3 - 2x^2 - 4x + 8) dx$ $= \left[\frac{x^4}{4} - \frac{2x^3}{3} - 2x^2 + 8x \right]_{-2}^2$ $= \left(4 - \frac{16}{3} - 8 + 16 \right) - \left(4 + \frac{16}{3} - 8 - 16 \right)$ $= \frac{64}{3}$	<p>M1</p> <p>M1</p> <p>dM1</p> <p>A1 (4)</p> <p>[14]</p>
<p>ALT</p> <p>NB</p>	<p>By splitting the area:</p> $\text{Area} = \int_{-2}^1 (x^3 - 2x^2 - 5x + 6) dx + \Delta [(-2, 0), (2, 0), P] - \left \int_1^2 (x^3 - 2x^2 - 5x + 6) dx \right $ <p>Integrate curve equation (ignore limits) and attempt area of triangle by formula or integration</p> <p>Substitute correct limits</p> $= \frac{64}{3}$ <p>No algebraic integration – only first M mark available</p>	<p>M1</p> <p>M1</p> <p>dM1</p> <p>A1</p>

Question Number	Scheme	Marks
<p>(a) (i)M1 M1 (A1 on e-PEN) A1cso (ii)A1 (B1 on e-PEN)</p>	<p>Obtain the quadratic factor by division or inspection Factor theorem allowed only if values for a and b are found.</p> <p>Factorise the quadratic factor</p> <p>Correct given value for <i>a</i></p> <p>Correct value for <i>b</i></p> <p>By Factor Theorem: M1 Test $x = 1$ M1 Test another value which is > 1 A1 $a = 1$ A1 $b = 3$</p>	
<p>(b) M1 A1 B1 M1 A1 A1cso ALT</p>	<p>Differentiate and substitute $x = 2$ to find the gradient of the tangent to <i>C</i> at <i>P</i></p> <p>Correct gradient of tangent</p> <p>$y = -4$ seen explicitly or used in the equation of the tangent</p> <p>Any complete method for the equation of the tangent at $(2, \text{their } y)$.</p> <p>Use of $y = mx + c$ must include an attempt at finding a value for <i>c</i></p> <p>Correct numbers in their (unsimplified) equation</p> <p>Correct <i>x</i> coordinate of the point where the tangent crosses the <i>x</i>-axis. No errors seen</p> <p>for the last 3 marks: Find the gradient of the line from <i>P</i> to $(-2, 0)$ M1 Any correct method; A1correct gradient All work correct and a conclusion. A1cso</p>	
<p>(c) M1 M1 dM1 A1 ALT M1 M1 dM1 A1</p>	<p>Using area = \int curve – line or \int line – curve with their line equation limits are needed</p> <p>Attempt to integrate the single function or two functions (ie all the integration needed) limits not needed. $\int_{-2}^2 (-x - 2) dx$ may be obtained by triangle formula.</p> <p>Substitute correct limits in their integrated function(s) Depends on both M marks</p> <p>Correct final answer. Must be positive.</p> <p>By splitting the area</p> <p>Suitable split eg as shown Limits are needed</p> <p>Attempt all the nec integration (ignore limits) and area of triangle by integration or formula</p> <p>Substitute correct limits in their integrated function(s) Depends on both M marks</p> <p>Correct final answer</p>	

Question Number	Scheme	Marks
10(a)	$\frac{y - -4}{-4 - 1} = \frac{x - -6}{-6 - 4} \Rightarrow y + 4 = \frac{1}{2}(x + 6) \quad \text{oe} \quad \text{eg } y = \frac{1}{2}x - 1$	M1A1 (2)
(b)	$\left(\frac{3 \times 4 + 2 \times -6}{5}, \frac{3 \times 1 + 2 \times -4}{5} \right) \Rightarrow (0, -1)$	M1A1 (2)
(c)	<p>Gradient of perpendicular = -2 Allow all following work if x, y used instead of m, n</p> $-2 = \frac{n - -1}{m - 0} \quad (\Rightarrow -2m = n + 1)$ $(3\sqrt{5})^2 = (m - 0)^2 + (n - -1)^2 \Rightarrow 45 = m^2 + (n + 1)^2$ $45 = m^2 + 4m^2 \Rightarrow 45 = 5m^2 \Rightarrow m = \pm 3 \text{ negative required } m = -3$	B1 B1ft M1A1
	$\Rightarrow n = -2m - 1 \Rightarrow n = -2 \times -3 - 1 = 5 \quad \text{coordinates are } (-3, 5)$	A1 (5)
(d)(i)	$RQ = \sqrt{(-13 - -3)^2 + (0 - 5)^2} = 5\sqrt{5}$ $AB = \sqrt{(4 - -6)^2 + (1 - -4)^2} = 5\sqrt{5}$ <p>With conclusion</p>	M1 A1cso
(ii)	$\left(\text{Gradient of } AB = \frac{1}{2} \right) \quad \text{Gradient of } RQ = \frac{5 - 0}{-3 - -13} = \frac{1}{2}$ <p>With conclusion *</p>	M1 A1cso (4)
ALT	<p>By vectors – combines both parts:</p> $\overrightarrow{AB} = 10\mathbf{i} + 5\mathbf{j} \quad \text{or equivalent column vector}$ $\overrightarrow{RQ} = 10\mathbf{i} + 5\mathbf{j} \quad \text{or equivalent column vector}$ <p>So same length and parallel (provided both vectors are correct)</p>	M1 M1 A1A1
(e)	<p>Area is base \times height $A = 3\sqrt{5} \times 5\sqrt{5} = 75 \quad (\text{units})^2$</p>	M1A1 (2) [15]
ALT:	$A = \frac{1}{2} \begin{vmatrix} -3 & 4 & -6 & -13 & -3 \\ 5 & 1 & -4 & 0 & 5 \end{vmatrix}$ $= \frac{1}{2} [(-3 - 20) + (-16 + 6) + (0 - 52) - (65 - 0)] = -75 \Rightarrow 75$	M1 A1

Question Number	Scheme	Marks
(a)		
M1	Any complete method for obtaining an equation of l	
A1	Correct equation in any form inc unsimplified	
(b)		
M1	Obtaining at least one of the coords of P . Must be correct. Can be by formula or diagram.	
A1	Both coords correct.	
	NB: If both coords are just written down, award M1A1 if both correct; M0A0 otherwise	
(c)		
B1	Correct gradient of the perpendicular	
B1ft	Correct equation connecting m and n from equating their gradient to -2 Can be unsimplified.	
	Follow through their gradient of the perpendicular but must be negative reciprocal of gradient of l	
M1	Use Pythagoras (with $+$ sign as shown oe)to find the length of PQ , equate this to $3\sqrt{5}$ and	
	solve to $m = \dots$	
A1	Correct value for m ± 3 allowed here	
A1	Correct value for n Values do not have to be written in coordinate brackets. Only one final	
	answer or this mark is lost.	
(d)		
(i)M1	Use Pythagoras to find the length of RQ or AB	
A1cso	Lengths of both lines correct with working for each and a conclusion shown	
(ii)M1	Find the gradient of RQ Must show working	
A1cso	Correct gradient of both lines and a conclusion shown	
ALT	M1M1 one M mark for each vector correct or working shown but slip made	
	A1A1 one A mark for each conclusion provided the vectors are correct.	
(e)		
M1	Obtaining the area of $ABPQ$ by using the formula for the area of a parallelogram	
A1	Correct area	
ALT:	Use the "determinant" method.	
M1	Formula must be correct ie $\frac{1}{2}$ needed, 5 pairs of coordinates with first and last the same,	
	coordinates to be in order round the quadrilateral (clockwise or anticlockwise). An attempt to	
	evaluate also needed.	
A1	Correct area - must be positive.	

Question Number	Scheme	Marks
11(a)	$AC = \sqrt{12^2 + 8^2} \quad (= \sqrt{208} = 4\sqrt{13})$ or AO or $OC = \sqrt{6^2 + 4^2} \quad (= 2\sqrt{13})$ $h = \sqrt{10^2 - 52} = \sqrt{48} = 4\sqrt{3} \quad *$	M1 M1A1cso (3)
(b)	$\angle OCE = \cos^{-1}\left(\frac{2\sqrt{13}}{10}\right) = 43.8537\dots \approx 43.9^\circ$ or $\sin^{-1}\left(\frac{4\sqrt{3}}{10}\right)$ or $\tan^{-1}\left(\frac{4\sqrt{3}}{2\sqrt{13}}\right)$	M1A1 (2)
(c)	Let M be the midpoint of BC . $EM = \sqrt{10^2 - 4^2} = 2\sqrt{21}$ $\cos \theta^\circ = \frac{4\sqrt{3}}{2\sqrt{21}} = \frac{2\sqrt{7}}{7} \quad *$ or cosine rule $\cos \theta^\circ = \frac{(4\sqrt{3})^2 + (2\sqrt{21})^2 - 6^2}{2 \times 4\sqrt{3} \times 2\sqrt{21}} = \frac{2\sqrt{7}}{7}$	M1 M1A1cso (3)
(d)	Let N be the midpoint of EM  $NO = \sqrt{(\sqrt{21})^2 + (4\sqrt{3})^2 - 2 \times \sqrt{21} \times 4\sqrt{3} \times \frac{2\sqrt{7}}{7}} \dots = \sqrt{21}$ hence triangle NEO is isosceles, so required angle ($\angle ENO$) $\angle ENO = 180 - 2 \times 40.8933\dots = 98.2134\dots \approx 98.2^\circ$	M1A1ftA1 B1 (4) [12]
ALT	Based on symmetry: $\tan \frac{\theta}{2} = \frac{\left(\frac{h}{2}\right)}{\frac{3}{2}} = \frac{2\sqrt{3}}{3}$ $\frac{\theta}{2} = 49.1066\dots$ $\theta = 98.2^\circ$	M1A1 A1 A1(B1 on e-PEN)

Question Number	Scheme	Marks
<p>(a) M1 M1 A1cso</p>	<p>Use Pythagoras with a + sign to find AC or AO or OC Use Pythagoras with a – sign to find the height Given height obtained from correct working. No decimals used but allow $2\sqrt{13} = 7.2\dots$ followed by $7.2\dots^2 = 52$</p>	
<p>(b) M1 A1</p>	<p>Use any trig function to obtain angle OCE Correct size of angle OCE Must be 1 dp</p>	
<p>(c) M1 M1</p>	<p>Use Pythagoras with a – sign to obtain the length of EM (need not be correct) $\cos \theta^\circ = \frac{4\sqrt{3}}{EM}$ with their EM or cosine rule as shown. Must reach $\cos \theta = \dots$ if other form used at start. (NB not dependent)</p>	
<p>A1cso</p>	<p>Correct completion to the given answer</p>	
<p>(d) M1 A1ft A1 B1</p>	<p>Use of cosine rule in $\triangle EON$ to obtain ON Correct numbers follow through their EM Correct length ON, exact or awrt 4.58 Correct size of angle, must be 1 dp unless already penalised in (b). (Can be obtained by the isos triangle as shown or by cosine or sine rule in $\triangle EON$)</p>	
	<p>NB: No A1ft in alt method as h is given in (a)</p>	