

Please check the examination details below before entering your candidate information

Candidate surname

Other names

**Pearson Edexcel
International GCSE**

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--

Thursday 20 June 2019

Morning (Time: 2 hours)

Paper Reference **4PM1/02R**

**Further Pure Mathematics
Paper 2R**



Calculators may be used.

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
 - *there may be more space than you need.*
- You must **NOT** write anything on the formulae page.
Anything you write on the formulae page will gain **NO** credit.

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
 - *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶

P61883A

©2019 Pearson Education Ltd.

1/1/1/1/1/1/



P 6 1 8 8 3 A 0 1 3 6



Pearson

International GCSE in Further Pure Mathematics Formulae sheet

Mensuration

Surface area of sphere = $4\pi r^2$

Curved surface area of cone = $\pi r \times \text{slant height}$

Volume of sphere = $\frac{4}{3}\pi r^3$

Series

Arithmetic series

Sum to n terms, $S_n = \frac{n}{2}[2a + (n - 1)d]$

Geometric series

Sum to n terms, $S_n = \frac{a(1 - r^n)}{(1 - r)}$

Sum to infinity, $S_\infty = \frac{a}{1 - r}$ $|r| < 1$

Binomial series

$(1 + x)^n = 1 + nx + \frac{n(n - 1)}{2!}x^2 + \dots + \frac{n(n - 1)\dots(n - r + 1)}{r!}x^r + \dots$ for $|x| < 1, n \in \mathbb{Q}$

Calculus

Quotient rule (differentiation)

$$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$$

Trigonometry

Cosine rule

In triangle ABC : $a^2 = b^2 + c^2 - 2bc \cos A$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

Logarithms

$$\log_a x = \frac{\log_b x}{\log_b a}$$



Answer all ELEVEN questions.

Write your answers in the spaces provided.

You must write down all the stages in your working.

$$f(x) = (x - 3)[x^2 + (p - 2)x + q]$$

Given that $f(0) = -12$

(a) find the value of q .

(2)

(b) Find the range of values of p for which the cubic equation $f(x) = 0$ has only one real root.

(5)

(Total for Question 1 is 7 marks)






Diagram **NOT**
accurately drawn

Figure 1

Figure 1 shows part of the curve S with equation $y = \frac{ax + b}{x + c}$ where a , b and c are integers.

The asymptote to S that is parallel to the x -axis has equation $y = -2$

The asymptote to S that is parallel to the y -axis has equation $x = -3$

The curve crosses the x -axis at the point with coordinates $(4, 0)$

The curve crosses the y -axis at the point with coordinates $(0, p)$ where p is a rational number.

Find

- (i) the value of a ,
- (ii) the value of b ,
- (iii) the value of c ,
- (iv) the value of p .

(4)



Question 2 continued

(Total for Question 2 is 4 marks)



3




Diagram NOT
accurately drawn

Figure 2

Figure 2 shows a solid right circular cylindrical metal rod.

The diameter of the rod is x cm and the length of the rod is $10x$ cm.

The rod is being heated so that the length of the rod is increasing at a rate of 0.005 cm/s.

Find the rate of increase, in cm^3/s to 2 significant figures, of the volume of the rod when $x = 3$

(6)

Question 3 continued

(Total for Question 3 is 6 marks)



- 4 A particle P moves along the x -axis. At time t seconds ($t \geq 0$) the acceleration, a m/s 2 , of P is given by $a = 6t - 12$

When $t = 0$, P is at rest at the origin.

- (a) Find the velocity of P when $t = 2$

(3)

At time T seconds, $T > 0$, P is instantaneously at rest.

- (b) Find the value of T .

(2)

- (c) Find the distance travelled by P in the first 8 seconds of its motion.

(3)



Question 4 continued

(Total for Question 4 is 8 marks)



- 5 (a) On the grid opposite, draw the graphs of the lines with equations

$$2x + 3y = 24 \quad y = 2x \quad 3y = 2x - 12$$

(3)

- (b) Show, by shading on the grid, the region R defined by the inequalities

$$2x + 3y \leq 24 \quad y \leq 2x \quad 3y \geq 2x - 12 \quad y \geq 0$$

(1)


For all points in R , with coordinates (x, y)

$$F = 2x + 5y$$

- (c) Find the greatest value of F .

(3)



Question 5 continued

Turn over for a spare grid if you need to redraw your graphs.



DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 5 continued

Only use this grid if you need to redraw your graphs.



(Total for Question 5 is 7 marks)



6 Given that $\sqrt{9-x}$ can be expressed in the form $p(1+qx)^{\frac{1}{2}}$ where p and q are constants

(a) find the value of p and the value of q .

(2)

(b) Hence expand $\sqrt{9-x}$ in ascending powers of x up to and including the term in x^3 expressing each coefficient as an exact fraction in its lowest terms.

(3)

Using the expansion you found in part (b) with a suitable value of x ,

(c) find an estimate to 5 decimal places for the value of $\sqrt{\frac{31}{4}}$

(3)



Question 6 continued



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 6 continued

(Total for Question 6 is 8 marks)



- 7 The n th term of a geometric series G is u_n

The first term of G is a and the common ratio of G is r , where $r > 0$

Given that $u_3 = 4$ and that $u_7 = 16$

(a) (i) show that $r = \sqrt{2}$

(ii) find the value of a .

(3)

(b) Find the least value of n for which $u_n > 500$

(4)

The sum of the first n terms of G is S_n

(c) Find S_{20}

Give your answer in the form $p(1 + \sqrt{2})$ where p is an integer.

(4)



Question 7 continued



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 7 continued

(Total for Question 7 is 11 marks)



Diagram NOT
accurately drawn




Figure 3

Figure 3 shows a right prism \$ABCDEF\$. The cross section \$BCF\$ of the prism is a triangle.

$$AB = DC = 12\text{ cm} \quad BC = AD = 8\text{ cm} \quad BF = AE = 10\text{ cm} \quad \angle FBC = \angle EAD = 60^\circ$$

The point \$N\$ lies on \$BC\$ such that \$FN\$ is perpendicular to \$BC\$.

(a) Show that \$BN = 5\text{ cm}\$.

(2)

(b) Find, in cm to 3 significant figures, the length of \$EN\$.

(3)

The midpoint of \$BF\$ is \$X\$ and the midpoint of \$FC\$ is \$Y\$.

(c) Find, in degrees to one decimal place, the size of the angle between the plane \$ABCD\$ and the plane \$AXYD\$.

(2)

(d) Find, in degrees to one decimal place, the size of the angle \$AYE\$.

(6)



Question 8 continued



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 8 continued

(Total for Question 8 is 13 marks)



- 9 The finite region R enclosed by the y -axis, the straight line with equation $y + 2x = 13$ and the curve with equation $y = x^2 - 2$, is defined for points with coordinates (x, y) with $x \geq 0$

The region R is rotated through 360° about the y -axis.

Use algebraic integration to find the volume of the solid generated.

Give your answer in terms of π .

(9)



Question 9 continued



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 9 continued

(Total for Question 9 is 9 marks)



10 (a) Use the formula for $\cos(A + B)$ to show that $\cos 2A = 2\cos^2 A - 1$ (2)

(b) Show that $\cos 4A = 8\cos^4 A - 8\cos^2 A + 1$ (4)

(c) Solve the equation $\cos^2\left(\frac{\theta}{4} + \frac{\pi}{24}\right)\left[\cos^2\left(\frac{\theta}{4} + \frac{\pi}{24}\right) - 1\right] = -\frac{1}{16}$ $0 \leq \theta < 2\pi$

Give your answers in terms of π . (5)

$$f(A) = 4\cos^4 A - 4\cos^2 A + 1$$

(d) Using calculus, find the exact value of $\int_{\frac{\pi}{6}}^{\frac{\pi}{2}} f(A) dA$

Give your answer in the form $a\pi - b\sqrt{c}$ where a and b are fractions in their lowest terms and c is a prime number.

(4)



Question 10 continued



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 10 continued

(Total for Question 10 is 15 marks)



11 The quadratic equation $x^2 - px + q = 0$ where $p > 0$, has roots α and β .

Given that $2\alpha\beta = 3$ and that $4(\alpha^2 + \beta^2) = k^2 - 6k - 3$ where $k > 3$

- (a) (i) write down the value of q ,
- (ii) find an expression, in terms of k , for p .

(5)

Given also that $7\alpha\beta = 3(\alpha + \beta)$

- (b) find the value of k .

(2)

- (c) Hence form an equation, with integer coefficients, which has roots

$$\frac{\alpha}{\alpha + \beta} \text{ and } \frac{\beta}{\alpha + \beta}$$

(5)



Question 11 continued



Question 11 continued

(Total for Question 11 is 12 marks)

TOTAL FOR PAPER IS 100 MARKS