

Please check the examination details below before entering your candidate information

Candidate surname

Other names

**Pearson Edexcel
International GCSE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Monday 20 January 2020

Morning (Time: 2 hours)

Paper Reference **4PM1/02**

Further Pure Mathematics

**Level 2
Paper 2**



Calculators may be used.

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
 - *there may be more space than you need.*
- You must **NOT** write anything on the formulae page.
Anything you write on the formulae page will gain **NO** credit.

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
 - *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶

P59939A

©2020 Pearson Education Ltd.

1/1/1/1/



P 5 9 9 3 9 A 0 1 3 6



Pearson

International GCSE in Further Pure Mathematics Formulae sheet

Mensuration

Surface area of sphere = $4\pi r^2$

Curved surface area of cone = $\pi r \times \text{slant height}$

Volume of sphere = $\frac{4}{3}\pi r^3$

Series

Arithmetic series

Sum to n terms, $S_n = \frac{n}{2}[2a + (n - 1)d]$

Geometric series

Sum to n terms, $S_n = \frac{a(1 - r^n)}{(1 - r)}$

Sum to infinity, $S_\infty = \frac{a}{1 - r} \quad |r| < 1$

Binomial series

$(1 + x)^n = 1 + nx + \frac{n(n - 1)}{2!}x^2 + \dots + \frac{n(n - 1)\dots(n - r + 1)}{r!}x^r + \dots$ for $|x| < 1, n \in \mathbb{Q}$

Calculus

Quotient rule (differentiation)

$$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$$

Trigonometry

Cosine rule

In triangle ABC : $a^2 = b^2 + c^2 - 2bc \cos A$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

Logarithms

$$\log_a x = \frac{\log_b x}{\log_b a}$$



Answer all ELEVEN questions.

Write your answers in the spaces provided.

You must write down all the stages in your working.

- 1 A particle P is moving along a straight line that passes through the fixed point O .
At time t seconds, $t \geq 0$, the displacement, s metres, of P from O is given by

$$s = t^3 + 4t^2 - 27t + 4$$

Find the value of t at the instant when the velocity of P is 8 m/s.

(4)

(Total for Question 1 is 4 marks)



P 5 9 9 3 9 A 0 3 3 6

- 2 Find the set of values of x for which
- (a) $3 + 2x \leqslant x + 2$ (1)
- (b) $8x^2 + 10x < 3$ (4)
- (c) **both** $3 + 2x \leqslant x + 2$ **and** $8x^2 + 10x < 3$ (1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 2 continued

(Total for Question 2 is 6 marks)






Diagram **NOT**
accurately drawn

Figure 1

Figure 1 shows a triangular pyramid $ABCD$.

The base, ABC , of the pyramid is a horizontal isosceles triangle with $AB = AC = 10\text{ cm}$ and $BC = 16\text{ cm}$. The midpoint of BC is M .

The face BCD of the pyramid is an isosceles triangle with $BD = CD = 26\text{ cm}$ and D is vertically above A .

$$\angle BAD = \angle CAD = 90^\circ$$

(a) Calculate the length, in cm, of AM .

(2)

Calculate, in degrees to the nearest degree,

(b) the size of $\angle BCD$,

(3)

(c) the size of the angle between the planes BCA and BCD .

(4)



Question 3 continued



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

(Total for Question 3 is 9 marks)



- 4 The points A , B , C and D are the vertices of a quadrilateral $ABCD$ such that

$$\overrightarrow{AB} = 7\mathbf{i} + p\mathbf{j} \quad \overrightarrow{AC} = 11\mathbf{i} - p\mathbf{j} \quad \overrightarrow{AD} = 4\mathbf{i} - 2p\mathbf{j}$$

- (a) Show that, for all values of p , $ABCD$ is a parallelogram.

(3)

Given that $|\overrightarrow{BD}| = 3\sqrt{10}$

- (b) find the possible values of p .

(3)

Given that $p > 0$

- (c) find a unit vector which is parallel to \overrightarrow{BD} .

(1)



Question 4 continued



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

(Total for Question 4 is 7 marks)



5 Given that α and β are such that $\alpha + \beta = \frac{7}{2}$ and $\alpha\beta = 2$

(a) form a quadratic equation with integer coefficients that has roots α and β ,

(2)

(b) form a quadratic equation with integer coefficients that has roots $\frac{\alpha}{\beta}$ and $\frac{\beta}{\alpha}$.

(6)



Question 5 continued

(Total for Question 5 is 8 marks)



Diagram NOT
accurately drawn




Figure 2

The region R , shown shaded in Figure 2, is bounded by the curve with equation $y = 2x - x^2$ and the line with equation $2y - x = 0$

The curve and the line intersect at the origin O and the point A .

- (a) Show that the point A has coordinates $\left(\frac{3}{2}, \frac{3}{4}\right)$. (2)

The region R is rotated through 360° about the x -axis.

- (b) Use algebraic integration to find, in terms of π , the volume of the solid formed. (6)



Question 6 continued



Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 6 continued

(Total for Question 6 is 8 marks)



7 The 7th term of a geometric series is 192 and the 8th term of this geometric series is 1152

(a) Find, as a fraction in its simplest form, the 4th term of this geometric series.

(3)

A different geometric series G has a common ratio r and n th term t_n

Given that $t_3 = 24$ and $t_2 + t_3 + t_4 = -36$

(b) show that r satisfies the equation

$$2r^2 + 5r + 2 = 0$$

(5)

Given further that G is convergent with sum to infinity S ,

(c) find the value of S .

(4)



Question 7 continued



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 7 continued

(Total for Question 7 is 12 marks)



8 Given that $y = e^{3x} \sin 2x$

show that $\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 13y = 0$

(8)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 8 continued



Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 8 continued

(Total for Question 8 is 8 marks)



P 5 9 9 3 9 A 0 2 7 3 6

- 9 A curve C has equation

$$y = \frac{qx - 2}{x - p} \quad x \neq p$$

The curve crosses the y -axis at the point A .

The line l with equation $y = x + 2$ is the normal to C at A .

(a) (i) Show that $p = 1$


(ii) Find the value of q . (7)

(b) Using the axes on the opposite page, sketch C , showing clearly the asymptotes and the coordinates of the points where C crosses the coordinate axes. (5)

The line l meets C again at the point D .

(c) Find the x coordinate of D . (4)



Question 9 continued

P 5 9 9 3 9 A 0 2 9 3 6

Question 9 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 9 continued

(Total for Question 9 is 16 marks)



P 5 9 9 3 9 A 0 3 1 3 6

10 The volume of a sphere is increasing at a constant rate of $40 \text{ cm}^3/\text{s}$.

Find the rate of increase, in cm^2/s , of the surface area of the sphere at the instant when the radius is 4 cm.

(9)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 10 continued

(Total for Question 10 is 9 marks)



P 5 9 9 3 9 A 0 3 3 3 6

11 (a) Express the equation

$$3 \sin(A - B) = \sin(A + B)$$

in the form $\tan A = k \tan B$, giving the value of the integer k .

(4)

(b) Given that $\theta \neq \frac{(2n+1)\pi}{2}$ where $n \in \mathbb{Z}$,

$$\text{show that } \frac{\cos^4 \theta - \sin^4 \theta}{\cos^2 \theta} = 1 - \tan^2 \theta$$

(3)

(c) Using the exact values of $\sin x^\circ$, $\cos x^\circ$ and $\tan x^\circ$ for $x = 30, 45, 60$

show that

$$(i) \cos 15^\circ = \frac{\sqrt{6} + \sqrt{2}}{4}$$

(2)

$$(ii) \tan 255^\circ = \frac{3 + \sqrt{3}}{3 - \sqrt{3}}$$

(4)



Question 11 continued



P 5 9 9 3 9 A 0 3 5 3 6

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 11 continued

(Total for Question 11 is 13 marks)

TOTAL FOR PAPER IS 100 MARKS

