Please check the examination details below before entering your candidate information

Candidate surname	Other names

Pearson Edexcel International GCSE

Centre Number | | | | | |

Candidate Number | | | | |

Thursday 18 June 2020

Morning (Time: 2 hours) | Paper Reference **4PM1/02**

Further Pure Mathematics

Paper 2

Calculators may be used.

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You must **NOT** write anything on the formulae page.
  Anything you write on the formulae page will gain NO credit.

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶

P62285A

1/1/1/

P 6 2 2 8 5 A 0 1 3 2

Pearson

International GCSE in Further Pure Mathematics Formulae sheet

Mensuration

Surface area of sphere = $4\pi r^2$

Curved surface area of cone = $\pi r \times$ slant height

Volume of sphere = $\frac{4}{3}\pi r^3$

Series

Arithmetic series

Sum to n terms, $S_n = \frac{n}{2}[2a + (n-1)d]$

Geometric series

Sum to n terms, $S_n = \frac{a(1-r^n)}{(1-r)}$

Sum to infinity, $S_\infty = \frac{a}{1-r} \quad |r| < 1$

Binomial series

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \ldots + \frac{n(n-1)\ldots(n-r+1)}{r!}x^r + \ldots \quad \text{for} \quad |x| < 1, n \in \mathbb{Q}$$

Calculus

Quotient rule (differentiation)

$$\frac{\mathrm{d}}{\mathrm{d}x}\left(\frac{\mathrm{f}(x)}{\mathrm{g}(x)}\right) = \frac{\mathrm{f}'(x)\mathrm{g}(x) - \mathrm{f}(x)\mathrm{g}'(x)}{[\mathrm{g}(x)]^2}$$

Trigonometry

Cosine rule

In triangle ABC: $a^2 = b^2 + c^2 - 2bc\cos A$

$$\tan\theta = \frac{\sin\theta}{\cos\theta}$$

$\sin(A+B) = \sin A \cos B + \cos A \sin B$ $\quad$ $\sin(A-B) = \sin A \cos B - \cos A \sin B$

$\cos(A+B) = \cos A \cos B - \sin A \sin B$ $\quad$ $\cos(A-B) = \cos A \cos B + \sin A \sin B$

$$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B} \qquad \tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

Logarithms

$$\log_a x = \frac{\log_b x}{\log_b a}$$

Answer all TEN questions.

Write your answers in the spaces provided.

You must write down all the stages in your working.

1 A particle P is moving in a straight line. At time t seconds, $t \geqslant 0$, the displacement, s metres, of P from a fixed point O of the line is given by

$$s = 3 + 8t + t^2 - \frac{1}{3}t^3$$

Find the distance of P from O when P is instantaneously at rest.

(4)

(Total for Question 1 is 4 marks)

2 The region enclosed by the curve with equation $y = e^{3x}$, the x-axis, the y-axis and the line with equation $x = 3$ is rotated through 360° about the x-axis.

Use algebraic integration to find, in terms of π and e, the volume of the solid generated.

(4)

Question 2 continued

(Total for Question 2 is 4 marks)

3 (a) Expand

$$(1 + px)^{-5} \quad p \neq 0$$

in ascending powers of x, up to and including the term in x^4
Give each term in its simplest form.

(3)

The coefficient of x^r in the expansion is c_r

Given that $c_4 = 2c_3$

(b) find the value of p.

(2)

Question 3 continued

(Total for Question 3 is 5 marks)

4 (i) Solve the equation $16\log_r 4 = \log_4 r$

(2)

(ii) Solve the equation $\log_5 9 + \log_5 12 + \log_5 15 + \log_5 18 = 1 + \log_5 x + \log_5 x^2$

(5)

Question 4 continued

(Total for Question 4 is 7 marks)

5 (a) Show that $\sum_{r=1}^{n}(3r+5) = \frac{1}{2}n(3n+13)$

(3)

(b) Hence evaluate $\sum_{r=35}^{50}(3r+5)$

(2)

Given that $\sum_{r=1}^{n}(3r+5) = 385$

(c) find the value of n.

(3)

Question 5 continued

(Total for Question 5 is 8 marks)

6 $f(x) = 4x^2 - 3x - 5$

The equation $f(x) = 0$ has roots α and β

Without solving the equation $f(x) = 0$

(a) form an equation, with integer coefficients, that has roots $\frac{2\alpha}{\beta}$ and $\frac{2\beta}{\alpha}$

(6)

$g(x) = 4x^2 + px + q$ where p and q are constants

Given that the equation $g(x) = 0$ has roots $3\alpha + \beta$ and $\alpha + 3\beta$

(b) find the value of p and the value of q

(5)

Question 6 continued

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 11 marks)

7 A geometric series has first term $(x - 3)$, second term $(x + 1)$ and third term $(4x - 2)$.

(a) Find the two possible values of x.

(5)

Given that $x < 1$,

(b) show that the series is convergent.

(2)

The sum to infinity of the series is S.

(c) Find the value of S.

(2)

The sum of the first n terms of the series is S_n

Given that $\frac{S}{S_n} = \frac{256}{255}$

(d) find the value of n.

(3)

Question 7 continued

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 7 continued

(Total for Question 7 is 12 marks)

8 The curve C_1 has equation $y = 5e^{-2x} + 4$

The curve C_2 has equation $y = e^{2x}$

The curves C_1 and C_2 intersect at the point A.

(a) Find the exact coordinates of A.

(4)

The tangent at A to C_1 intersects the x-axis at the point B.

(b) Show that the x coordinate of B is $\frac{1}{2}(5 + \ln 5)$

(5)

The tangent at A to C_2 intersects the x-axis at the point D.

(c) Find the area of ΔABD.

(6)

Question 8 continued

Question 8 continued

Question 8 continued

(Total for Question 8 is 15 marks)

9 A curve C has equation

$$y = \frac{2 + 4x - x^2}{2x + 1} \qquad x \neq -\frac{1}{2}$$

(a) Write the equation of C in the form $ax^2 + (by - 4)x + (y - c) = 0$, where a, b and c are integers whose values are to be found.

(3)

(b) Hence show that x is real when $y \leqslant 2$ and when $y \geqslant 3$

(4)

(c) Find the coordinates of the stationary points on C.

(6)

(d) Sketch C showing clearly

(i) the exact coordinates of the points where C crosses the x-axis,

(ii) the asymptote to C that is parallel to the y-axis,

(iii) the coordinates of the stationary points.

(5)

Question 9 continued

Question 9 continued

Question 9 continued

(Total for Question 9 is 18 marks)

10 (a) Show that

$$\cos(A + B) + \cos(A - B) = 2\cos A \cos B$$

(2)

(b) Hence show that

$$\cos P + \cos Q = 2\cos\frac{P + Q}{2}\cos\frac{P - Q}{2}$$

(3)

(c) Solve, for $0 \leqslant \theta \leqslant \frac{\pi}{2}$, the equation

$$\cos 5\theta + \cos 7\theta = 0$$

Give each solution in terms of π

(4)

(d) Show that

$$\cos 8x + 2\cos 6x + \cos 4x = 4\cos 6x \cos^2 x$$

(3)

(e) Use calculus to find the exact value of

$$\int_0^{\frac{\pi}{3}} \cos 6x \cos^2 x \, dx$$

(4)

Question 10 continued

Question 10 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 10 continued

Question 10 continued

(Total for Question 10 is 16 marks)

TOTAL FOR PAPER IS 100 MARKS