Candidate surname

Other names

Pearson Edexcel International GCSE

Centre Number

Candidate Number

Time 2 hours

Paper reference **4PM1/01**

Further Pure Mathematics
PAPER 1

Calculators may be used.

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
  – *there may be more space than you need*.
- You must **NOT** write anything on the formulae page.
  Anything you write on the formulae page will gain NO credit.

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question*.

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.
- Good luck with your examination.

Turn over

Pearson

International GCSE in Further Pure Mathematics Formulae sheet

Mensuration

Surface area of sphere $= 4\pi r^2$

Curved surface area of cone $= \pi r \times$ slant height

Volume of sphere $= \dfrac{4}{3}\pi r^3$

Series

Arithmetic series

Sum to n terms, $S_n = \dfrac{n}{2}\left[2a + (n-1)d\right]$

Geometric series

Sum to n terms, $S_n = \dfrac{a(1 - r^n)}{(1 - r)}$

Sum to infinity, $S_\infty = \dfrac{a}{1 - r}$ $|r| < 1$

Binomial series

$(1 + x)^n = 1 + nx + \dfrac{n(n-1)}{2!}x^2 + \ldots + \dfrac{n(n-1)\ldots(n-r+1)}{r!}x^r + \ldots$ for $|x| < 1, n \in \mathbb{Q}$

Calculus

Quotient rule (differentiation)

$$\dfrac{\mathrm{d}}{\mathrm{d}x}\left(\dfrac{\mathrm{f}(x)}{\mathrm{g}(x)}\right) = \dfrac{\mathrm{f}'(x)\mathrm{g}(x) - \mathrm{f}(x)\mathrm{g}'(x)}{[\mathrm{g}(x)]^2}$$

Trigonometry

Cosine rule
In triangle ABC: $a^2 = b^2 + c^2 - 2bc\cos A$

$\tan\theta = \dfrac{\sin\theta}{\cos\theta}$

$\sin(A + B) = \sin A \cos B + \cos A \sin B$ $\qquad$ $\sin(A - B) = \sin A \cos B - \cos A \sin B$

$\cos(A + B) = \cos A \cos B - \sin A \sin B$ $\qquad$ $\cos(A - B) = \cos A \cos B + \sin A \sin B$

$\tan(A + B) = \dfrac{\tan A + \tan B}{1 - \tan A \tan B}$ $\qquad$ $\tan(A - B) = \dfrac{\tan A - \tan B}{1 + \tan A \tan B}$

Logarithms

$\log_a x = \dfrac{\log_b x}{\log_b a}$

Answer all ELEVEN questions.

Write your answers in the spaces provided.

You must write down all the stages in your working.

1 The quadratic equation

$$3(k + 2)x^2 + (k +5)x + k = 0$$

has real roots.

Find the set of possible values of k.

(6)

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

(Total for Question 1 is 6 marks)

2 Angle α is acute such that $\cos\alpha = \dfrac{3}{5}$

Angle β is obtuse such that $\sin\beta = \dfrac{1}{2}$

(a) Find the exact value of

   (i) $\tan\alpha$

   (ii) $\tan\beta$

(3)

(b) Hence show that

$$\tan(\alpha + \beta) = \frac{m\sqrt{3} - n}{n\sqrt{3} + m}$$

where m and n are positive integers whose values are to be found.

(3)

(Total for Question 2 is 6 marks)

3 A curve C has equation $y = \dfrac{ax - 3}{x + 5}$ where a is a constant and $x \neq -5$

The gradient of C at the point on the curve where $x = 2$ is $\dfrac{18}{49}$

(a) Show that $a = 3$

(3)

Hence

(b) write down an equation of the asymptote to C that is

   (i) parallel to the x-axis,

  (ii) parallel to the y-axis,

(2)

(c) find the coordinates of the point where C crosses

   (i) the x-axis,

  (ii) the y-axis.

(2)

(d) Sketch the curve C, showing clearly its asymptotes and the coordinates of the points where C crosses the coordinate axes.

(3)

P 6 6 0 2 4 A 0 7 3 6

Question 3 continued

(Total for Question 3 is 10 marks)

4 The nth term of an arithmetic series is u_n where

$$u_n = (n+1)\ln 4$$

Given that the sum of the first n terms of the series is S_n

show that $S_n = \ln 2^{(n^2 + an)}$ where a is an integer whose value is to be found.

(5)

Question 4 continued

(Total for Question 4 is 5 marks)

5 (a) Expand $(1 + ax)^n$ in ascending powers of x up to and including the term in x^3

Express each coefficient of x in terms of a and n where a and n are constants and $n > 2$

(2)

The coefficient of x is 15 and the coefficient of x^2 is equal to the coefficient of x^3

(b) Find the value of a and the value of n.

(6)

(c) Find the coefficient of x^3

(2)

(Total for Question 5 is 10 marks)

6 (a) Show that $(\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$

(3)

The quadratic equation $x^2 - 7kx + k^2 = 0$, where k is a positive constant, has roots α and β where $\alpha > \beta$

(b) Show that $\alpha - \beta = 3k\sqrt{5}$

(3)

(c) Hence form a quadratic equation with roots $\alpha + 1$ and $\beta - 1$

Give your equation in the form $x^2 + px + q = 0$ where p and q should be given in terms of k.

(4)

P 6 6 0 2 4 A 0 1 4 3 6

P 6 6 0 2 4 A 0 1 5 3 6

Question 6 continued

(Total for Question 6 is 10 marks)

P 6 6 0 2 4 A 0 1 7 3 6

7 The curve C has equation $y = \dfrac{x}{x^2 + 4}$

(a) Using calculus, find the coordinates of the stationary points on C.

(5)

(b) Show that $\dfrac{d^2 y}{dx^2} = \dfrac{2x(x^2 - 12)}{(x^2 + 4)^3}$

(4)

(c) Hence, or otherwise, determine the nature of each of these stationary points.

(2)

P 6 6 0 2 4 A 0 1 9 3 6

(Total for Question 7 is 11 marks)

8 Given that n satisfies the equation

$$\log_a n = \log_a 3 + \log_a(2n - 1)$$

(a) find the value of n.

(3)

Given that $\log_p x = 3$ and $\log_p y - 3 \log_p 2 = 4$

(b) (i) express x in terms of p,

(1)

(ii) express xy in terms of p.

(4)

P 6 6 0 2 4 A 0 2 3 3 6

Question 8 continued

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

(Total for Question 8 is 8 marks)

P 6 6 0 2 4 A 0 2 5 3 6

9 Find an equation of the normal to the curve with equation

$$y = (x^3 - 2x)e^{(1-x)}$$

at the point on the curve with coordinates $(1, -1)$

(5)

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

(Total for Question 9 is 5 marks)

Diagram **NOT** accurately drawn

Figure 1

Figure 1 shows triangle *OAB* and triangle *OCD*.

$$\overrightarrow{OA} = 5\mathbf{p} \quad \overrightarrow{AB} = 3\mathbf{q} \quad \overrightarrow{OC} = \frac{3}{2}\overrightarrow{OB} \quad \overrightarrow{OD} = \frac{3}{5}\overrightarrow{OA}$$

(a) Find $\overrightarrow{DC}$ as a simplified expression in terms of **p** and **q**.

(3)

The line *DC* meets the line *AB* at *F*.

(b) Using a vector method, find $\overrightarrow{OF}$ as a simplified expression in terms of **p** and **q**.

(7)

The point *G* lies on *OB* such that *FG* is parallel to *AO*.

(c) Using a vector method, find $\overrightarrow{OG}$ as a simplified expression in terms of **p** and **q**.

(4)

Question 10 continued

P 6 6 0 2 4 A 0 3 0 3 6

Question 10 continued

(Total for Question 10 is 14 marks)

11 (a) Using a formula from page 2, show that $\cos 2x = 1 - 2\sin^2 x$

(3)

Diagram **NOT** accurately drawn

$y = \sin x + 2$

R_2

A

R_1

B

$y = \cos 2x + 2$

C

O

x

Figure 2

Figure 2 shows a sketch of part of the curves with equations $y = \sin x + 2$ and $y = \cos 2x + 2$

The points A, B and C, shown in Figure 2, are three points that are common to both curves.

(b) Find the coordinates of each of these points.

(4)

R_1 and R_2, shown shaded in Figure 2, are two regions enclosed by the two curves.

(c) Use calculus to find, in its simplest form, the ratio

$$\text{area of } R_1 : \text{area of } R_2$$

(8)

P 6 6 0 2 4 A 0 3 3 3 6

Question 11 continued

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

(Total for Question 11 is 15 marks)

TOTAL FOR PAPER IS 100 MARKS