

Mark Scheme (Results)

Summer 2019

Pearson Edexcel International GCSE In Mathematics A (4MA1) Paper 1HR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2019
Publications Code 4MA1_1HR_1906_MS
All the material in this publication is copyright
© Pearson Education Ltd 2019

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.
 - Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Types of mark

- o M marks: method marks
- o A marks: accuracy marks
- o B marks: unconditional accuracy marks (independent of M marks)

Abbreviations

- o cao correct answer only
- o ft follow through
- isw ignore subsequent working
- SC special case
- o oe or equivalent (and appropriate)
- o dep dependent
- o indep independent
- o awrt answer which rounds to
- o eeoo each error or omission

No working

If no working is shown then correct answers normally score full marks If no working is shown then incorrect (even though nearly correct) answers score no marks.

With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

If a candidate misreads a number from the question. Eg. Uses 252 instead of 255; method marks may be awarded provided the question has not been simplified. Examiners should send any instance of a suspected misread to review. If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

• Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded to another.

International GCSE Maths

Apart from questions 5a, 10, 18a, 19, 22, 23, 24 (where the mark scheme states otherwise) the correct answer, unless clearly obtained from an incorrect method, should be taken to imply a correct method.

| Question | Working | Answer | Mark | | Notes |
|----------|---|--------|------|----|-------------------------------|
| 1 | e.g. $\pi \times 8.2^2$ (= 211.24, $\frac{1681}{25}\pi$) | | 3 | M1 | for a correct first step |
| | or 1.5 × 1000 (= 1500) | | | | |
| | or $\pi \times 8.2^2 \times 10$ (= 2112.4, $\frac{3362}{5}\pi$) | | | | |
| | e.g. (1.5 × 1000) ÷ (π × 8.2²) (= 7.1009) | | | M1 | for a complete method to find |
| | or (1.5 × 1000) ÷ "2112.4" × 10 oe (= 7.1009) | | | | the depth of the water or an |
| | or 10 – (("2112.4" – 1.5 × 1000) ÷ (π × 8.22)) (= 7.1009) | | | | answer of 2.89 – 2.91 |
| | | 7.1 | | A1 | accept 7.09 – 7.11 |
| | | | | | Total 3 marks |

| 2 | Ext \angle = 180 - 162 (= 18) oe or $\frac{(n-2)180}{n}$ = 162 oe | | 3 | M1 |
|---|--|----|---|---------------|
| | 360 ÷ "18" oe or 18 <i>n</i> = 360 | | | M1 |
| | | 20 | | A1 |
| | | | | Total 3 marks |

| 3 (i) | | 12, 18 | 1 | B1 |
|--------------|---|---------------------------|---|---|
| (ii) | | 12, 14, 15, 16, 18, 20 | 1 | B1 |
| (iii) | | 11, 13, 15, 17, 19 | 1 | B1 |
| | | | | Total 3 marks |
| | • | • | | |
| 4 | e.g. $4x - 8x = 17 + 13$ oe | | 2 | M1 For collecting terms in <i>x</i> and number terms on either side of a correct equation |
| | | -7.5 | | A1 oe e.g. $-\frac{30}{4}$ |
| | | | | Total 2 marks |
| 5 (a) | e.g. 720 = 2 × 360 = 2 × 2 × 180 or 720 = 3 × 240 = 3 × 3 × 80 etc | | 3 | M1 At least 2 correct stages in prime factorisation |
| | 2, 2, 2, 2, 3, 3, 5 | | | M1 condone inclusion of 1 (may be a fully correct factor tree or ladder) |
| | | 2 × 2 × 2 × 2 × 3 × 3 × 5 | | A1 dep on M2, accept $2^4 \times 3^2 \times 5$ |

1

B1

Total 4 marks

(b)

| 6 | (a) | 4.25 × 0.08 (= 0.34) oe | | 3 | M1 | M2 for 4.25 × 1.08 oe |
|---|-----|--------------------------|------|---|--------------------|--------------------------|
| | | 4.25 + "0.34" | | | M1 | 4.23 ** 1.00 00 |
| | | | 4.59 | | A1 SC: B1 for 4.25 | × 0.92 (= 3.91) oe |
| | (b) | 9.45 ÷ 108 (= 0.0875) oe | | 3 | M1 | M2 for — 9.45 ÷ 1.08 |
| | | 9.45 ÷ 108 × 100 oe | | | M1 | 9.43 · 1.08 |
| | | | 8.75 | | A1 | |
| | | | | | | Total 6 marks |

| 7 | 7.5 ² - 6 ² (= 20.25) | | 4 | M1 | OR for a correct trig statement involving one of the angles |
|---|---|----|---|----|--|
| | , ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,, | | | | e.g. $\cos BAM = \frac{6}{7.5}$ or $\sin ABC = \frac{6}{7.5}$ |
| | | | | | where <i>M</i> is the midpoint of <i>BC</i> |
| | $\sqrt{7.5^2-6^2}$ (= 4.5) | | | M1 | OR for a method to find one of the angles in the triangle e.g. |
| | \(\frac{7.5}{-0} \) (-4.5) | | | | $BAM = \cos^{-1}\left(\frac{6}{7.5}\right) (=36.8) \text{ or } ABC = \sin^{-1}\left(\frac{6}{7.5}\right) (=53.1)$ |
| | "4.5"× 6 oe | | | M1 | for a complete method to find the area of triangle ABC e.g. |
| | 4.5 ^ 0 00 | | | | $2 \times \frac{1}{2} \times 7.5 \times 6 \times \sin("36.8")$ oe or $2 \times \frac{1}{2} \times 7.5 \times \sqrt{7.5^2 - 6^2} \times \sin("53.1")$ oe |
| | | 27 | | A1 | cao |
| | | | | | Total 4 marks |

| | | | | | | ີ າ | N 4 1 | |
|----|-----|--|--------|------------------------|---|------|--|--------------------|
| 8 | | 10 × 79.2 (= 792) or 3 × 68 (= 204) | | | | 3 | M1 | |
| | | $(10 \times 79.2 - 3 \times 68) \div 7$ | | | | | M1 | |
| | | | 84 | | | | A1 | |
| | | | | | | | | Total 3 marks |
| | | | | | | | | |
| 9 | (a) | | | <i>t</i> ⁶ | 1 | B1 | | |
| | (b) | | ı | № ¹² | 1 | B1 | | |
| | (c) | | 12 | $5x^3y^6$ | 2 | B2 | | |
| | . , | | | | | (B1 |) for 2 correct ter | rms as part of a |
| | | | | | | | | Total 4 marks |
| | | | | | | | | |
| 10 | | 22 × 60 × 60 (= 79 200) oe or 22 ÷ 1000 (= 0.02 | 22) oe | | 3 | m | or converting from n/s to m/h or from n to km | M2 for 22 × 3.6 oe |
| | | 22 × 60 × 60 ÷ 1000 oe | | | | | or a complete nethod | |
| | | | | 79.2 | | A1 o | e, dep on at least M | 1 |
| | | | | | | | | Total 3 marks |

| 11 | 15 - 3 : x - 3 = 2 : 7 or (15 - 3) ÷ 2 (= 6) | $(n =) (15 - 3) \div \frac{2}{2+7}$ (= 54) where n is the total age 3 years ago | | 3 | M1 | M2 for $\frac{(15-3)\times7}{2}$ (= 42) |
|----|--|--|----|---|----|---|
| | $\frac{x-3}{15-3} = \frac{7}{2}$ oe or $7 \times \text{"6"} (= 42)$ | "54" × $\frac{7}{2+7}$ (= 42) | | | M1 | |
| | | | 45 | | A1 | |
| | | | | | | Total 3 marks |

| 12 | 105 ÷ (5 × 4) (=5.25) oe or 105 ÷ (4 × 3) (=8.75) oe or 105 ÷ (3 × 5) (=7) | | 3 | M1 |
|----|--|-----|---|--|
| | "8.75" – "5.25" | | | M1 dep on previous M1. If M1 gained and they have worked out 3 pressures, award M1 for their highest minus their lowest. |
| | | 3.5 | | A1 oe |
| | | | | Total 3 marks |

| 13 (a) | A B | 1 | B1 | Professional judgment required, eg allow double shading if meaning clear. |
|---------------|----------|---|----|---|
| (b) | (D∪E) ∩F | 1 | B1 | oe eg (<i>D∩F</i>)∪(<i>E∩F</i>) |
| | | | | Total 2 marks |

| 14 | (a) | | 0.7, 0.2, 0.8, 0.1, 0.9 | 2 | B2 | oe, all correct |
|----|-----|---------------------------------------|-------------------------|---|------|---|
| | | | | | (B1) | 2 or 3 or 4 correct probabilities |
| | (b) | 0.3 × "0.2" (= 0.06) | | 4 | M1 | ft from (a) dep on probabilities being between 0 and 1, OR $0.3 \times "0.8"$ (= 0.24) |
| | | or 0.7 × "0.1" (= 0.07) oe | | | | or 0.7 × "0.9" (= 0.63) oe |
| | | 0.3 × "0.2" + 0.7 × "0.1" (= 0.13) oe | | | M1 | ft from (a), OR 0.3 × "0.8" + 0.7 × "0.9" (= 0.87) oe |
| | | or "0.06" × 200 (= 12) | | | | or "0.24" × 200 (= 48) |
| | | or "0.07" × 200 (= 14) | | | | or "0.63" × 200 (= 126) |
| | | "0.13" × 200 oe or "12" + "14" | | | M1 | ft from (a), 200 – "0.87" × 200 oe or (1 – "0.87") × 200 or 200 – "48" – "126" |
| | | | 26 | | A1 | cao |
| | | | | | | Total 6 marks |

| 15 | (Gradient of L ₁ =) 6 ÷ 2 (=3) | | 4 | M1 | could be seen as part of an equation. Ignore constant term if candidate rearranges L ₁ |
|----|---|------------------------|---|----|--|
| | $m \times "3" = -1 \text{ or } m = -\frac{1}{"3"}$ | | | M1 | for use of $m_1m_2 = -1$ could be seen as part of an equation |
| | $-1 = "-\frac{1}{3}" \times 9 + c$ or $y1 = "-\frac{1}{3}"(x - 9)$ or $c = 2$ | | | M1 | |
| | | $y + \frac{1}{3}x = 2$ | | A1 | oe in required form eg $3y + x = 6$, 6y + 2x = 12 etc |
| | | | | | Total 4 marks |

| 16 (a) | $3\times 4t^2 - 2\times 6t + 5$ | | 2 | M1 | For 2 terms correct |
|---------------|---------------------------------|-------------------|---|----|--|
| | | $12t^2 - 12t + 5$ | | A1 | Fully correct |
| (b) | 24t-12 | | 3 | M1 | Method to differentiate their <i>v</i> , ft a 3 term quadratic expression from (a) |
| | 24t - 12 = 6 | | | M1 | ft if previous M1 awarded |
| | | 0.75 | | A1 | oe |
| | | | | | Total 5 marks |

| 17 (a) | e.g. one correct value on the vertical scale e.g. 1 at 1 cm high or 1 cm ² = 5 passengers or 5 small squares = 1 passenger or (FD =) 24 ÷ 20 (= 1.2) | | 3 | M1 | For a correct scale on the vertical axis or a 1 cm × 1 cm square = 5 passengers or other correct scale or one correct product or frequency (other than the 24) or (FD =) 24 ÷ 20 (= 1.2) |
|---------------|--|-------|---|------|--|
| | 10×0.4 (= 4) 10×1.8 (= 18) 5×6.4 (= 32) 15×2 (= 30) 20×0.8 (= 16) | | | M1 | At least 3 correct products or frequencies (other than the 24) stated (could be seen on diagram) |
| | | 124 | | A1 | |
| (b) | e.g. 0.25 × 24 + 20 × 0.8 (= 22) or "1.2" × 5 + 20 × 0.8 (= 22) | | 2 | M1 | ft from (a) |
| | | "22" | | A1ft | oe (0.17(741)) |
| | | "124" | | | Total 5 marks |
| | | | | | |

| 18 (a) | $(x+2)(2x+3) = 2x^2 + 3x + 4x + 6$ $(2x+3)(x-7) = 2x^2 - 14x + 3x - 21$ $(x+2)(x-7) = x^2 - 7x + 2x - 14$ | | 3 | M1 | For multiplying a pair of brackets and getting 3 out of 4 terms correct. |
|---------------|---|----------------------------------|---|-------|---|
| | $(2x^2+7x+6)(x-7) = 2x^3-14x^2+7x^2-49x+6x-42$ $(2x^2-11x-21)(x+2) = 2x^3+4x^2-11x^2-22x-21x-42$ $(x^2-5x-14)(2x+3) = 2x^3+3x^2-10x^2-15x-28x-42$ | | | M1dep | For multiplying the product of the first 2 brackets (ft from the 1 st stage) by the 3 rd bracket, and getting at least 3 out of 6 or 4 out of 8 terms correct |
| | | $2x^3 - 7x^2 - 43x - 42$ | | A1 | Fully correct. isw extra work as long as correct e.g. $x(2x^2 - 7x - 43) - 42$ |
| | Alternative (all in one method) | | | | |
| | (x+2)(2x+3)(x-7) = | | | M2 | For at least 6 out of 8 correct terms |
| | $2x^3 - 14x^2 + 3x^2 - 21x + 4x^2 - 28x + 6x - 42$ | | | (M1) | for 4 or 5 out of 8 correct terms |
| | | $2x^3 - 7x^2 - 43x - 42$ | | A1 | |
| (b) | $p^{2}(2m-y) = x+m$ $2p^{2}m-p^{2}y = x+m$ | | 3 | M1 | Multiplying by denominator and expanding bracket |
| | e.g. $2p^2m - m = x + p^2y$ $m(2p^2 - 1) = x + p^2y$ | | | M1 | Collect terms in <i>m</i> and factorise in a correct equation |
| | | $m = \frac{x + p^2 y}{2p^2 - 1}$ | | A1 | oe eg $m = \frac{-x - p^2 y}{1 - 2p^2}$ must have $m =$ |

| | | | | | | Total 6 marks |
|----|---|------|-------------------|----------------|--------------------|---|
| | | 1 | | T | ' | |
| 19 | a + 24d = 44.5 | | 5 | M1 | oe | |
| | $\frac{30}{2}(2a+(30-1)d) = 765 \text{ oe,}$ | | | M1 | oe (n | nay be simplified) |
| | eg (15(2a + 29d) = 765), (2a + 29d = 51), etc | | | | | |
| | e.g. 2(44.5 – 24d) + 29d = 51 oe or | | | M1 | - | on M2, a complete method to eliminate one ble, allow one arithmetic error |
| | _2a + 48d = 89 | | | | | |
| | <u>2a + 29d = 51</u> oe | | | | | |
| | | 26.5 | | A2 | dep | on M2, oe |
| | | | | (A1) | dep or d | on M2. If not A2, award A1 for $a = -3.5$ |
| | | | | | | Total 5 marks |
| | | 1 | | | | |
| 20 | 125 or 10 ²¹ⁿ | | | | 3 | M1 |
| | 125×10^{21n} | | | | | M1 |
| | | 1.25 | × 10 ² | 1 <i>n</i> + 2 | | A1 |
| | | | | | | Total 3 marks |

| 21 (a)(i) | (9, 3) | 1 | B1 | |
|------------------|-----------------------------|---|------|---|
| (a)(ii) | (4, 9) | 1 | B1 | |
| (b) | <i>a</i> = −2, <i>b</i> = 3 | 2 | B2 | or $a = 2$, $b = -3$ |
| | | | (B1) | for $a = -2$ or $a = 2$ or $b = 3$ or $b = -3$ |
| | | | | Total 4 marks |

| 22 | $2x^2 + 3(2x + 1)^2 = 5$ | | 5 | M1 | $2\left(\frac{y-1}{2}\right)^2 + 3y^2 = 5$ |
|----|--|---|---|----|---|
| | eg $14x^2 + 12x - 2 = 0$ or if completing the square, allow $14x^2 + 12x = 2$ oe | | | A1 | $7y^2 - 2y - 9 = 0$ or if completing the square, allow $7y^2 - 2y = 9$ oe |
| | eg $(7x - 1)(x + 1)$ or $(7x - 1)(2x + 2)$ eg $\frac{-12 \pm \sqrt{12^2 - 4 \times 14 \times -2}}{2 \times 14}$ oe eg $7\left(\left(x + \frac{3}{7}\right)^2 - \frac{9}{49}\right) = 2$ oe | | | M1 | ft as long as M1 awarded and 3 term quadratic eg $(7y - 9)(y + 1)$ eg $\frac{2 \pm \sqrt{(-2)^2 - 4 \times 7 \times -9}}{2 \times 7}$ oe eg $7\left(\left(y - \frac{1}{7}\right)^2 - \frac{1}{49}\right) = 9$ oe |
| | $x = \frac{1}{7}, x = -1 $ (need both) | | | A1 | $y = \frac{9}{7}, y = -1 $ (need both) |
| | | $x = \frac{1}{7}$, $y = \frac{9}{7}$ x = -1, $y = -1$ | | A1 | Dep on M1 Must be paired correctly Must be 3 sf or better (0.142857) (1.28571) |
| | | | | | Total 5 marks |

| 23 | <i>ABF</i> = 180 – <i>x</i> or <i>CDF</i> = 180 – <i>x</i> | | 4 | M1 | for finding an expression for <i>ABF</i> or <i>CDF</i> . May be seen on diagram. |
|----|---|-----|---|----|---|
| | FDE = $180 - (180 - x) (= x)$ AFB or $ACE = 180 - (180 - x) - 54 (= x - 54)$ DFE or $ACE = 180 - x - 32 (= 148 - x)$ e.g. $54 + y + 180 - x = 180$ where $AFB = y$ 32 + y + (180 - (180 - x)) = 180 where $DFE = y$ | | | M1 | method to find <i>FDE</i> and <i>AFB</i> or method to find <i>FDE</i> and <i>DFE</i> or method to find <i>ACE</i> or method to find <i>FDE</i> and an equation for <i>AFB</i> e.g. $54 + y + 180 - x = 180$ where $AFB = y$ or method to find <i>FDE</i> and an equation for <i>DFE</i> e.g. $32 + y + (180 - (180 - x)) = 180$ where $DFE = y$ May be seen on diagram. |
| | e.g. $32 + x + x - 54 = 180$ or $54 + 180 - x + 148 - x = 180$ or $x - 54 = 148 - x$ oe or $54 + y + 180 - x = 180$ and $32 + y + (180 - (180 - x)) = 180$ where $AFB = DFE = y$ | | | M1 | for setting up an equation or a pair of correct simultaneous equations to solve for x |
| | | 101 | | A1 | dep on at least M1 |
| | | | | | Total 4 marks |

| 24 | $\overrightarrow{AP} = \frac{3}{4} \times 2\mathbf{c} \ (=\frac{3}{2}\mathbf{c}) \text{ oe}$ | 5 | For $\overrightarrow{AP} = \frac{3}{2} \mathbf{c}$ oe, eg could be part of $\overrightarrow{OP} = \mathbf{a} + \frac{3}{2} \mathbf{c}$ oe or on diagram |
|----|--|-----|---|
| | \overrightarrow{AC} = c - a oe or \overrightarrow{CA} = a - c oe | | M1 |
| | \overrightarrow{OQ} = c + n (a - c) or \overrightarrow{OQ} = a + n (c - a) | | M1 |
| | or $\overrightarrow{QP} = n(\mathbf{a} - \mathbf{c}) + \frac{3}{2}\mathbf{c}$ | | |
| | $\frac{n}{1-n} = \frac{2}{3} \Rightarrow n = \frac{2}{5} \text{oe or}$ | | M1 |
| | $\frac{1-n}{n} = \frac{2}{3} \Rightarrow n = \frac{3}{5} \text{oe or}$ | | |
| | $\frac{n}{\frac{3}{2}-n} = \frac{2}{3} \implies n = \frac{3}{5} \text{ oe}$ | | |
| | | 3:2 | A1 oe, dep on M3 |
| | | | Total 5 marks |

| 25 | e.g. (220 – 180) + (360 – 280) (= 120) | | 5 | M1 | for a method to find angle <i>XYZ</i> . Could be seen on a diagram |
|----|--|-------|---|----|---|
| | $XZ = \sqrt{3.5^2 + 6^2 - 2 \times 3.5 \times 6 \times \cos("120")}$ (=8.3 or $\frac{\sqrt{277}}{2}$) | | | M1 | |
| | $\frac{\sin YXZ}{6} = \frac{\sin("120")}{"8.32"}$ | | | M1 | or $6^2 = 3.5^2 + "8.32"^2 - 2 \times 3.5 \times "8.32" \times \cos YXZ$ |
| | $YXZ = \sin^{-1}\left(\frac{6\sin("120")}{"8.32"}\right)$ (=38.6) | | | M1 | or $YXZ = \cos^{-1}\left(\frac{3.5^2 + "8.32"^2 - 6^2}{2 \times 3.5 \times "8.32"}\right) (= 38.6)$ |
| | | 241.4 | | A1 | accept 241.2 – 241.4 |
| | | | | | Total 5 marks |

