

Mark Scheme (Results)

January 2017

Pearson Edexcel International GCSE Mathematics B (4MB0)

Paper 02R



Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2017
Publications Code 4MB0_02R_1701_MS
All the material in this publication is copyright
© Pearson Education Ltd 2017

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Types of mark

- M marks: method marks
- A marks: accuracy marks
- B marks: unconditional accuracy marks (independent of M marks)

Abbreviations

- o cao correct answer only
- ft follow through
- o isw ignore subsequent working
- o SC special case
- o oe or equivalent (and appropriate)
- o dep dependent
- o indep independent
- o eeoo each error or omission

No working

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another


January 2017 – Paper 2R Mark Scheme

1.	2(-2) + x = 4		M1		
	x = 8		A1		
	(3+y)(-2)+(-3)"8"=-16 (subst)		M1 dep		
	y = -7		A1	4 4	ļ
	NB: 1 st M is for a correct equation				
	SC: $\binom{(3+y)(-2)+(-3)x}{2(-2)+x} = \binom{-16}{4}$ scores 1 st M1 (see	eeing one correct equation)			
2.	$2 \times (\text{base area}) = (400\sqrt{2})^2 \text{ (oe)}$	$2 \times (\text{side})^2 = \left(400\sqrt{2}\right)^2 \text{ (oe)}$	M1		
	base area = 160000 cm ² (oe cao) Allow ISW	side = 400 (cm) (oe cao) Allow ISW	A1		
	$\frac{1}{3} \times "\left(\frac{400\sqrt{2} \times 400\sqrt{2}}{2} \times \frac{1}{10^4}\right) " \times \frac{150}{10^2} \text{ (oe)}$	$\frac{1}{3} \times " \left(\frac{400 \times 400}{10^2 \times 10^2} \right) " \times \frac{150}{10^2} $ (o.e.)	M1 dep		
	8 m ³ NB: (1) The 1 st A is for a correct side length (cm or m) or c (2) The 2 nd M is for a correct volume statement using their of 150 cm to m,		A1	4 4	ļ

3.	(a)	$\frac{65}{100} \times 360$ (oe)	M1		
		234	A1	2	
	(b)	$\frac{5}{1+3+5}$ ×"234"	M1	2	
		130	A1	2	
	(c)	$\frac{3}{1+3+5}$ ×"234" (78)	M1		
		"78" : "130"-4	M1 dep		
		13:21	A1	3	7

4.	(a)	55 (m)	B1	1	
	(b)	one term correctly differentiated (ie $3t^2$ or -27)	M1		
		$3t^2 - 27$	A1	2	
	(c)	$3t^2 - 27'' = 0$	M1		
		t = 3 (cao)	A1	2	
	(d)	1 (m)	B1 ft	1	
	(e) NB: T	$(5)^3 - 27 \times 5 + 55$ (45) The "45" might be seen in a table	M1		
		("55"-"1")+("45"-"1")	M1 dep		
		98 (m)	A1	3	9

5. (a)



5,4 2,7,8 x,3x

(b) correctly adding all their terms and equating to 50

6

(c) (i) (3×"6"+8+5+7+4 = OR 50 - "6" - "2" =) 42 (ii) ("2" + "5" + "8" =) 15 B1, B1 4 B1 B1 M1 A1 2

> B1 ft, B1 ft

2

(d) $\frac{"6"}{"20"}$ (o.e.) ft numerator, ft denominator	B1 ft, B1 ft	2	10
"20" SC: Case where their Venn Diagram has 7 in place of 2, 12 in place of 7 and 13 in place of 8:			
(a) scores B1 B1 B0 B1			
(b) should be $4x + 41 = 50$ (M1), $x = 2.25$ (A0)			
(c) (i) B0, (ii) 25 B1ft (d) 3/35 B1 B0			

6.	$\frac{-5-20}{5} < x \text{ (o.e.)} \text{OR} x < \frac{(13-20)}{5} \text{ (oe)}$	M1		
	$\frac{-5-20}{5} < x$ AND $x < \frac{(13-20)}{5}$	M1 dep		
	-4, -3 , $-2NB: (1) Use of ≤ or = correctly gains only the M marks$	A2 (-1 eeoo)	4 4	

7.	(a)	(i) 12	(ii) 1.5 (o.e.)	B1, B1	2
	(b)	$\frac{1}{3x}$		B1	1
	(c)	(i)	$+\frac{5}{4}$, $-\frac{5}{4}$ (o.e.)	B1, B1	
		(ii)	$2(9x^2 - 3x - 3x + 1) = x$	M1	
			$18x^2 - 13x + 2 \ (=0)$	A1	
			attempt to solve a trinomial quadratic	M1	
			$\frac{2}{9}$, awrt 0.222	A1	
			$\frac{1}{2}$, 0.5	A1	7 10

	1			Т	1
8.	(a)	(i)	$\frac{1}{3}$, awrt 0.333	B1	1
		(ii)	$\frac{1}{3} \times \frac{1}{2} (A \to B \to A) \text{or} \frac{1}{3} \times \frac{1}{3} (A \to D \to A)$	M1	
			$\frac{1}{3} \times \frac{1}{2} + \frac{1}{3} \times \frac{1}{3}$	M1 dep	
			$\frac{5}{18} \left(\frac{15}{54} , \text{ awrt } 0.278 \right)$	A1	3
		(iii)	$\frac{1}{3} \times \frac{1}{2} \left(A \to B \to C \right) + \frac{1}{3} \times \frac{2}{3} \left(A \to D \to C \right)$	M1	
			$\frac{7}{18}$ (awrt 0.389)	A1	
			ith reference to part (ii)) so $\frac{7}{18} > \frac{5}{18}$ t eg P(aii) > P(aiii) (cso)	A1	3

$\frac{1}{3} \times \frac{1}{2} \times \frac{1}{3} + \frac{2}{3} \times \frac{1}{3} \times \frac{1}{3} \times \left(\frac{7}{54}\right)$	M1 dep		
$(C \rightarrow B \rightarrow A \rightarrow \text{Exit}) \qquad (C \rightarrow D \rightarrow A \rightarrow \text{Exit})$ $\frac{7}{54} \text{ or } 0.129 \text{ (o.e.) seen + conclusion}$ $NB: A sufficient conclusion would be "7/54 = 0.13"$	A1	3	10

	T	1
9. (a) (i) $6c-2a$ (ii) $5c$	B1, B1	
(iii) $\overrightarrow{AN} = -2\mathbf{a} + "5\mathbf{c}"$ $\overrightarrow{OM} = \frac{1}{2} (2\mathbf{a} + "5\mathbf{c}") \text{ (gains M2)}$	M1	
$\overrightarrow{OM} = \begin{cases} \overrightarrow{OA} + \overrightarrow{AM} = 2\mathbf{a} + \frac{1}{2}"(-2\mathbf{a} + 5\mathbf{c})" \\ \overrightarrow{ON} + \overrightarrow{NM} = "5\mathbf{c}" - \frac{1}{2}"(-2\mathbf{a} + 5\mathbf{c})" \end{cases}$	M1 dep	
$\boxed{\overrightarrow{ON} + \overrightarrow{NM} = "5\mathbf{c}" - \frac{1}{2}"(-2\mathbf{a} + 5\mathbf{c})"}$		
$\mathbf{a} + \frac{5}{2}\mathbf{c}$	A1	5
NB: $\frac{1}{2}(2\mathbf{a} + 5\mathbf{c})$ earns all 3 marks		
(b) $\overrightarrow{PM} = \begin{cases} \overrightarrow{PO} + \overrightarrow{OM} = -\mathbf{a} + "\left(\mathbf{a} + \frac{5}{2}\mathbf{c}\right)" \\ \overrightarrow{PA} + \overrightarrow{AM} = \mathbf{a} + \frac{1}{2}"\left(-2\mathbf{a} + 5\mathbf{c}\right)" \end{cases}$	M1	
correct conclusion	A1	2
NB: Must be a conclusion based on the directions of \overrightarrow{OC} and \overrightarrow{PM} and not just on their ratio.		

(c) area of triangle $OAN = \frac{5}{6} \times 30$ (25)	M1		
area of triangle $APM = \frac{1}{2} \times \frac{1}{2} \times$ " area of triangle OAN "	M1 dep		
6.25 , $\frac{25}{4}$ (square units)	A1		
NB: Ignore vector division in (c), eg $\frac{\mathbf{a}}{2\mathbf{a}} = \frac{1}{2}$		3	10

10.	Penalise incorrect rounding (i.e. not giving answers to 3 significant figures) ONCE only in the question, the first time it		
	occurs		
	$\frac{80}{100} = \frac{110}{100}$	M1	
	(a) $\frac{\cos \angle ACB}{\sin \angle ACB} = \frac{110}{\sin 75}$		
		M1 dep	
	$\sin\angle ACB = 80 \times \frac{\sin 75}{110}$	Wii dep	
	110	A1	3
	$\angle ACB = 44.6 \ (44.6272)$		
	$\angle ACD = 44.0 \ (44.0272)$		
		B1	
	(b) $\angle ABC = 180 - (75 + 44.6)$ (60.4, 60.3727)		
	Cosine Rule:		
	Cosine Ruie.	M1	
	$(AC^2 =)$ $80^2 + 110^2 - 2 \times 80 \times 110 \times \cos 60.4$	M1 dep	
	40.700 0.500 0.7	Wii dep	4
	= 18500-8693.37	A1	
	AC = 99.0 m (98.9916)		

Sine Rule:		
$\frac{AC}{\sin"60.4"} = \frac{110}{\sin 75} \text{ (oe, } = \frac{80}{\sin"44.6"} \text{)} (M1)$		
$AC = \frac{110}{\sin 75} \times \sin'' 60.4''$ (oe) (M1 dep)		
AC = 99.0 (98.9916), 99.1 (A1) NB: Accept 99 for A1		
$80^2 + 55^2 - 2 \times 80 \times 55 \times \cos 60.4$		
(c) $ (AM^{2} =) \begin{cases} 80^{2} + 55^{2} - 2 \times 80 \times 55 \times \cos 60.4 \\ "99"^{2} + 55^{2} - 2 \times "99" \times 55 \times \cos 44.6 \end{cases} $	M1	
9425-4346.69		
$(AM^2 =) \begin{cases} 9425 - 4346.69 \\ 12826 - 7753.96 \end{cases}$	M1 dep	
(71.3 <i>m</i> (71.262)		
$AM = \begin{cases} 71.3 \ m \ (71.262) \\ 71.2 \ m \ (71.218) \end{cases}$	A1	3
$(d) \qquad \frac{PA}{80} = \tan 41$		
80	M1	
69.5 m (69.5429)	A1	2

(e)	$QM^{2} = \begin{cases} "71.3"^{2} + ("69.543"/2)^{2} \\ "71.2"^{2} + ("69.5"/2)^{2} \end{cases}$	M1	
	$QM = \begin{cases} 79.3 & (79.327) \\ 79.2 & (79.227) \end{cases}$	A1	2
(f)	$\tan QMA = \frac{"69.5"/2}{"71.3"} \text{(oe)}$	M1	
NB: A	26.0° (26.0175) Accept 26 for A1	A1	2 16

11.	(a) $3(-3)^3 + k(-3)^2 - 27 \times -3 + 36 = 0$		
	-81+9k+81+36=0	M1	
	correct conclusion	A1	2
	Algebraic division Method: $3x^2 - 13x + 12$ (M1)		
	$\begin{array}{ccc} 3x & -13x + 12 & \text{(M1)} \\ \text{Statement of zero denominator} & \text{(A1)} \end{array}$		
	(b) $3x-4-\frac{27}{x}+\frac{36}{x^2}$ (dividing by x^2 , no slips)		
	$ \begin{array}{cccc} & & & & & & \\ & & & & & \\ & & & & & \\ & & & & $	M1	
	Multiply $\frac{27}{x} - \frac{36}{x^2} = px + q$ by $x^2 \implies 27x - 36 = px^3 + qx^2 \implies px^3 + qx^2 - 27x + 36$		
	and comparing coefficients		
	p=3 , $q=-4$	A1, A1	
	OR 27 36		
	$\frac{27}{x} - \frac{36}{x^2} = 3x - 4$		2
	So A1 for 3 <i>x</i> , A1 for -4		3
	(c) -1.4, 4.5, 5	D1 D1 D1	2
		B1, B1, B1	3
	Note: Accept -1.44 without penalty		

(d)	graph penalties (-1)	straight line segments	В3	
	each point missed ($\pm \frac{1}{2}$ small sq.)		(-1 eeoo)	3
	each missed segment			
	each point not plotted			
	each point incorrectly plotted ($\pm \frac{1}{2}$ small sq.)			
	tramlines			
	very poor curve i.e. line too thick			
(e)	straight line, gradient = 3 AND intersecting their curve TWICE		M1	
	intercept on y-axis "-4"			
~			A1 ft	2
OR			71110	_
	y = p''x + q'' going through two points on their line Going through "one	e point" AND intersecting their	71111	2
Their	y = "p"x + "q" going through two points on their line Going through "one TWICE" (M1)	e point" AND intersecting their	AT It	2
Their		e point" AND intersecting their	AT II	2
Their	TWICE (M1)	e point" AND intersecting their	AT II	2
Their	TWICE (M1)	e point" AND intersecting their		

Pearson Education Limited. Registered company number 872828 with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom