

Mark Scheme (Results)

Summer 2013

International GCSE
Physics (4PH0) Paper 1 PR

Science Double Award (4SC0)
Paper 1 PR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.

Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at www.edexcel.com/ask. You will need an Edexcel username and password to access this service




Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2013

Publications Code UG036657

All the material in this publication is copyright
© Pearson Education Ltd

Question number		Answer	Notes	Marks
1 (a)		total internal reflection	Accept TIR	1
1 (b) (i)		prism drawn in correct orientation (by eye) 	Accept a freehand sketch of the triangular prism Size of prism unimportant, e.g. can fill the entire square, but horizontal and vertical edges must be drawn	1
1 (b) (ii)		correct reflection of rays (by eye): 	Accept freehand sketch Accept correct external reflection e.g. reflection as shown below gets 1 mark for 1(b)(ii) despite the error in the 1(b)(i) response 	1

Question number	Answer	Notes	Marks
2 (a) (i)	B- 2 cm		1
(ii)	C- 8 cm		1
(b)	<p>Idea that in a transverse wave the direction of vibration is perpendicular to the direction of the wave; (May be shown with labels on the diagram)</p> <p>Idea that longitudinal wave the direction of vibration is parallel to the direction of the wave; (May be shown with labels on the diagram)</p> <p>A named freehand sketch of either wave indicating the two directions; e.g.</p> <div data-bbox="465 884 808 983" style="border: 1px solid red; padding: 5px;"> </div> <p>transverse</p> <div data-bbox="465 1023 907 1137" style="border: 1px solid red; padding: 5px;"> </div> <p>Longitudinal</p>	<p>Allow (for vibration) oscillation / displacement / disturbance (for direction of wave) direction of travel / energy / transfer (for perpendicular) at right angles, is \perp to (for parallel) the same as, //</p> <p>the minimum labeling is to name of the type of wave they have drawn.</p> <p>Allow sine waves with appropriate arrows</p> <p>Allow diagrams indicating compression and rarefaction e.g. in a spring</p> <p>Allow for 1 mark (but only if other mark is scored) a comparison of the directions of vibration of both waves without relating them to the direction of the wave</p> <p>e.g. transverse vibrates up and down but longitudinal vibrates back and forward</p>	3
(c)	any two of		2

		<p>MP1 can travel through vacuum OR needs no medium;</p> <p>MP2 speed (in a vacuum) OR speed = 3×10^8 (m/s);</p> <p>MP3 obeys laws of reflection / refraction;</p> <p>MP4 obeys wave equation OR speed = frequency \times wavelength;</p> <p>MP5 carries energy/ information;</p> <p>MP6 they are transverse</p>	<p>“speed in a vacuum” where seen, scores 2 marks (MP1 and MP2)</p> <p>Accept reflect, refract, diffract</p>	
(d)	i	D - X-rays		1
	ii	A – absorbed by the bone		1
	iii	<p>X-rays OR gamma rays</p> <p>idea of causing damage to cancer cells e.g. cells killed/mutated/ionised/destroys;</p>	<p>allow symbol γ do not allow UV</p> <p>Independent mark</p>	2

Question number		Answer	Notes	Marks									
3 (a)		<table border="1" data-bbox="566 268 1142 539"><tr><td data-bbox="566 268 784 411">temperature</td><td data-bbox="784 268 967 411">boiling point of nitrogen</td><td data-bbox="967 268 1142 411">boiling point of water</td></tr><tr><td data-bbox="566 411 784 475">in °C</td><td data-bbox="784 411 967 475">-196</td><td data-bbox="967 411 1142 475"></td></tr><tr><td data-bbox="566 475 784 539">in Kelvin</td><td data-bbox="784 475 967 539"></td><td data-bbox="967 475 1142 539">373</td></tr></table> <p data-bbox="465 619 801 651">one mark for each correct;;</p>	temperature	boiling point of nitrogen	boiling point of water	in °C	-196		in Kelvin		373	ignore -273	2
temperature	boiling point of nitrogen	boiling point of water											
in °C	-196												
in Kelvin		373											

Question number		Answer	Notes	Marks
4 (a)		any three of MP1 idea that there is current (in the wire/coil); MP2 idea that (the coil has) a magnetic field; MP3 idea that coil's magnetic field interacts with field of permanent magnet; MP4 idea that there is a force on the coil/wire; MP5 Idea that current or force reverses every half turn;	Allow ideas of electromagnetic field, electromagnet Allow - 'magnetic fields touch / overlap' Ignore - 'cutting of magnetic fields' Allow ideas of LHM rule, Fleming's LHR, catapult field, attraction, repulsion Allow action of a commutator described	3


<p>(b) (i)</p>		<p>any two of</p> <p>MP1 increase magnetic field(e.g. stronger magnets or magnets closer or magnets curved round coil);</p> <p>MP2 increase current OR voltage Or more cells;</p> <p>MP3 increase number of turns (on coil);</p> <p>MP4 a sensible alternative suggestion e.g. use two or more sets of coils at angles, lubricate axle;</p>	<p>Allow “use thicker wire”</p> <p>Ignore “stronger battery”</p> <p>Allow idea of 3 phase supply, iron stator</p>	<p>2</p>
	<p>(ii)</p>	<p>Suggestion that clearly results in reversal of the current OR the cell connections OR the magnet’s field;</p>		<p>1</p>
<p>(c)</p>		<p>any two of</p> <p>MP1 Idea that force is increased (by stronger field);</p> <p>MP2 Idea of radial magnetic field (rather than a uniform one);</p> <p>MP3 Coil remains in the field for a longer time;</p>	<p>Allow idea that iron is magnetised</p> <p>Allow idea that magnetic field acts “all the way around”</p> <p>Allow idea that force acts over a larger part of a cycle</p>	<p>2</p>

Question number	Answer	Notes	Marks
5 (a) B D	constant velocity of <u>5 m/s</u> Idea that velocity/speed = 0	Allow speed is <u>5 m/s</u> Allow “stops”, “stationary”, “at rest”	2
(b)	Idea of greater slope (for stage E); e.g. the gradient is steeper	Allow reverse argument, provided stage A is identified e.g. “stage A has a shallower slope” Allow attempts to demonstrate through - calculation of both gradients - qualitative comparison of data	1
(c)	distance = speed × time OR distance = area under graph; attempt to find any area; attempt to total correct areas (or use trapezium method); evaluation; e.g. distance = area under graph 7×7 or $\frac{1}{2} \times 7 \times 3$ $(7 \times 7) + (\frac{1}{2} \times 7 \times 3) = 49 + 10.5$ 59.5 (m)	The correct relationship can be implicit in the working 59.5 (m) with no working = full marks Allow the trapezium method - e.g. $7 \times ((7+10) \div 2) = 7 \times 8.5$ = 59.5 (m)	4
(d)	Correct equation shown ; e.g. (average speed) = distance (moved) / time (taken) Substitution of correct distance and suitable time; Correct evaluation ; e.g. $106.5/27$ 3.94 (m/s)	Allow d/t Allow (ecf) max 2 4.26 (m/s) (use of time = 25 s) 3.55 (m/s) (use of time = 30 s) Allow reverse argument max 2 e.g. $106.5 \div 4 = 26.6$ (s)	3

Question number		Answer	Notes	Marks
6 (a)	(i)	any three of Idea of collisions / impact (with walls); Continuous bombardment; force produced; Pressure = force ÷ area;	Ignore collisions between particles Allow idea of momentum changing	3
	(ii)	Idea that the student is right OR the pressure decreases; AND any two of The number(or mass) of molecules stays the same; The gas volume increases; Pressure is inversely proportional to volume; Particles collide with the wall less frequently;	Both marks depend on previous correct response (e.g. pressure decreases) Allow idea that area of can in contact with gas increases OR gas particles have more space Allow mention of $p_1V_1 = p_2V_2$ in this context Allow “longer time between collisions”	3
(b)		(Average speed) increases;		1

Question number		Answer	Notes	Marks
7	(a) (i)	pressure difference = height (or depth) x density x g ;	Allow $h \times \rho \times g$ (and rearrangements) Reject “gravity” for g in 7(a)(i)	1
	(ii)	substitution into correct equation; evaluation; e.g. $1028 \times 10 \times 700$ $7\,196\,000$ (Pa)	Allow standard form Allow use of $g = 9.8(1) \rightarrow 7\,059\,276$ or $7\,052\,080$	2
	(iii)	(total pressure =) $72 \times 10^5 + 1 \times 10^5$ (Pa);	Allow $7\,296\,000$ (Pa) OR answer to 7(a)(ii) + $100\,000$	1
(b)	(i)	pressure = force/area	Allow $p = F/A$	1
	(ii)	Substitution into correct equation; Transformation; Evaluation; e.g. $41 \times 10^5 = F/3.1$ $F = 41 \times 10^5 \times 3.1$ 1.271×10^7 (N)	Substitution and transposition either order $12\,710\,000$, 127.1×10^5 , 1.3×10^7	3
(c)		because fresh water has a lower density than sea water OR reverse argument;		1
(d)		any five of MP1 suitable measuring instruments mentioned; e.g. measuring cylinder and (electronic) balance MP2 method of obtaining correct mass; e.g. subtract mass of container, use of tare MP3 detail to ensure accuracy of liquid volume; e.g. burette, pipette, density bottle, account taken of meniscus MP4 equation stated - density = mass \div volume; MP5 suitable units used, e.g. g for mass and cm^3 for volume MP6 Idea of appropriate repeating or averaging at any stage	Allow scales Ignore newtonmeter, weighing machine Ignore weight Allow keep temperature constant Allow $\rho = m/V$ Allow ml, l Allow “discard anomalous results”	5

Question number	Answer	Notes	Marks																		
8 (a)	<p>all 3 for both marks;;</p> <p>any two for 1 mark ;</p> <table border="1" data-bbox="385 368 1016 874"><thead><tr><th data-bbox="385 368 752 437">item</th><th data-bbox="752 368 1016 437">Tick if needed</th></tr></thead><tbody><tr><td data-bbox="385 437 752 491">ammeter</td><td data-bbox="752 437 1016 491"></td></tr><tr><td data-bbox="385 491 752 545">steel spring</td><td data-bbox="752 491 1016 545"></td></tr><tr><td data-bbox="385 545 752 600">retort stand and clamp</td><td data-bbox="752 545 1016 600">✓</td></tr><tr><td data-bbox="385 600 752 654">rubber band</td><td data-bbox="752 600 1016 654">given ✓</td></tr><tr><td data-bbox="385 654 752 708">ruler</td><td data-bbox="752 654 1016 708">✓</td></tr><tr><td data-bbox="385 708 752 762">thermometer</td><td data-bbox="752 708 1016 762"></td></tr><tr><td data-bbox="385 762 752 817">mass hanger</td><td data-bbox="752 762 1016 817">✓</td></tr><tr><td data-bbox="385 817 752 871">mass</td><td data-bbox="752 817 1016 871">given ✓</td></tr></tbody></table>	item	Tick if needed	ammeter		steel spring		retort stand and clamp	✓	rubber band	given ✓	ruler	✓	thermometer		mass hanger	✓	mass	given ✓	each incorrect tick = -1	2
item	Tick if needed																				
ammeter																					
steel spring																					
retort stand and clamp	✓																				
rubber band	given ✓																				
ruler	✓																				
thermometer																					
mass hanger	✓																				
mass	given ✓																				

(b) i	5.5 (in the table)		1																					
ii	<p>suitable scale for axes; axes labelled with units; points plotted to nearest mm square (minus one for each plotting, up to max 2 marks);; Line (curve) of best fit acceptable;</p> 	<p>-1 for each incorrect plot Allow (ecf) a balanced straight line of best fit that takes account of any plotting errors and indicated anomalies</p> <table border="1" data-bbox="1187 502 1803 853"> <thead> <tr> <th>Mass in g</th> <th>Force in N</th> <th>Extension in cm</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>0</td> <td>0.0</td> </tr> <tr> <td>150</td> <td>1.5</td> <td>2.4</td> </tr> <tr> <td>350</td> <td>3.5</td> <td>6.3</td> </tr> <tr> <td>550</td> <td></td> <td>12.8</td> </tr> <tr> <td>750</td> <td>7.5</td> <td>18.6</td> </tr> <tr> <td>1050</td> <td>10.5</td> <td>24.0</td> </tr> </tbody> </table>	Mass in g	Force in N	Extension in cm	0	0	0.0	150	1.5	2.4	350	3.5	6.3	550		12.8	750	7.5	18.6	1050	10.5	24.0	5
Mass in g	Force in N	Extension in cm																						
0	0	0.0																						
150	1.5	2.4																						
350	3.5	6.3																						
550		12.8																						
750	7.5	18.6																						
1050	10.5	24.0																						
iii	<p>No / yes (no mark)</p> <p>Idea that Hooke's law should show (direct) proportionality;</p> <p>Use of data (from the table or graph) to explain that the results do not show this; e.g. 'line is a curve', '(table shows) rubber band extends unevenly'</p>	<p>Allow (ecf) - converse from <u>straight</u> drawn line, using data from their graph (not the table) e.g. 'Yes' AND 'line is a straight'</p>	2																					
		Total	10																					

Question number	Answer	Notes	Marks	
9 (a)	i	0.45;	no unit penalty	1
	ii	Power = current × voltage;	Allow $P = I \times V$ and rearrangements	1
	iii	Substitution; Evaluation; e.g. $1.5 = I \times 0.45$ $I = 3.3$ (A) (answer to at least 2 s.f.)	Allow reverse argument yielding <u>1.35</u> (W) for 1 mark	2
(b)	i	conversion of time to seconds; substitution into correct equation ($E = I \times V \times t$); evaluation; e.g. time = $7 \times 5 \times 60 \times 60$ (= 126 000) $E = 3.3 \times 9 \times 7 \times 5 \times 60 \times 60$ 3 742 000 (J)	Allow solution in stages i.e. from $P=IV$ and $P =E/t$ Allow for full marks 3 402 000 (J) (from use of 3 A given above) 3 780 000 (J) (from $1.5 \times 20 \times 7 \times 5 \times 60 \times 60$) Allow max of 1 if time not in seconds, e.g. 1040 (J) (from $3.3 \times 9 \times 7 \times 5$, time in hours) 62400 (J) (from $3.3 \times 9 \times 7 \times 5 \times 60$, time in minutes)	3
	ii	A description to include electrical; to light (and heat);	Reject “electricity” for the first mark Allow chemical to electrical to light for 1 mark only	2
		Total	9	

Question number	Answer	Notes	Marks
1 (a)	<p>i MP1 Any circuit including correct circuit symbols for</p> <ul style="list-style-type: none"> • battery /cell / d.c. power supply • ammeter • voltmeter ; <p>ii MP2 ammeter clearly measures current through the wire; MP3 voltmeter clearly across wire;</p> <p>Idea of measuring current through the wire;</p> <p>Idea of measuring voltage across the wire;</p> <p>Idea of a range of values (of I and V); e.g. alter variable resistor OR repeat for different voltages</p>	<p>ignore other components for MP1</p> <p>allow even if voltmeter in series with ammeter allow circuit line drawn through meter allow voltmeter across a section of the test wire</p>	<p>3</p> <p>3</p>
(b)	<p>i any one of resistance changes (with temperature) ;</p> <p>ii wire gets hot and melts/burns/catches fire/dangerous;</p> <p>V proportional to I only at constant temperature;</p> <p>(c) Ohms Law is only true if temperature constant;</p> <p>i any one of putting the wire in a water bath ; taking the reading quickly; switching off between readings; using only small currents; voltage = current × resistance ;</p> <p>ii horizontal line above axis;</p>	<p>Reject incorrect relationship between R and θ</p> <p>Ignore damage to wire</p> <p>Reject insulating the wire</p> <p>Allow to return to room temperature</p> <p>Allow $V = I \times R$ and rearrangements</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p>

Question number	Answer	Notes	Marks
11 (a) i	GPE = mass \times g \times height ;	Allow GPE = $m \times g \times h$ and rearrangements	1
ii	Substitution into correct equation; Evaluation; e.g. $0.25 \times 10 \times 1.75$ 4.375 (J)	Reject "gravity" for g in 11(a)(i) 4.4, 4.38 Allow use of 9.81 (or 9.8) \rightarrow 4.29 for full marks	2
(b)	Value given in 11(a)(ii);		1
(c) i	KE = $\frac{1}{2} \times$ mass \times speed ² ;	Allow KE = $\frac{1}{2} \times m \times v^2$ and rearrangements	1
ii	Substitution into correct equation; Transformation; Evaluation; e.g. $3.1 = \frac{1}{2} \times 0.25 \times v^2$ $v^2 = 3.1 \div \frac{1}{2} \times 0.25$ $v = 4.98$ (m/s)	Substitution and transposition either order Accept 5.0, 5 and allow truncation e.g. 4.97 m/s	3
		Total	11

Question number		Answer	Notes	Marks
12 (a)		<p>A description to include any 5 of MP1 nucleus absorbs neutron OR nucleus hit by neutron; MP2 splits into (two) fragments/parts OR daughter atoms OR daughter nuclei; MP3 extra neutrons released; MP4 (kinetic) energy released; MP5 released neutrons hit further nuclei OR uranium nuclei; MP6 moderator slows down the neutrons/ makes it more likely for a neutron to be absorbed; MP7 control rods absorb extra neutrons; MP8 idea that control rods help prevent a “runaway” chain reaction;</p>	<p>Correct process using consistently incorrect particle instead of neutron (e.g. electron) = max 4 NB uranium, U-235 or nucleus must be mentioned Reject cells, molecules, more uranium Ignore heat allow atoms OR uranium atoms</p>	5
(b)		kinetic/movement energy;		1
(c)		Idea that the shielding absorbs radiation / particles / energy;	<p>Allow “stops radiation / particles from escaping” Ignore “radioactivity” escaping</p>	1
			Total	12

Question number	Answer	Notes	Marks
13 (a) i	there is a voltage; And one of (because there is a) change of flux OR field (lines) are cut; (which is) an induced voltage / emf;	Allow induced current	2
ii	greater deflection/voltage; Idea that rate of change of flux (linkage) is greater; eg more magnetic field lines cutting coil (per second)	ignore speed of magnet	2
(b) i	Idea that deflection is smaller;		1
ii	Idea that deflection is greater;		1
iii	Idea that deflection is in opposite direction;		1
		Total	7

Question number	Answer	Notes	Marks	
14 (a)	i	(Nuclei / atoms) with same number of protons OR same atomic number; different number of neutrons OR different mass number;	Ignore electrons Allow “(nuclei) of the same element” Allow different number of nucleons	2
	ii	(stable isotopes) do not emit (ionising) radiation OR (stable isotopes) do not emit alpha, beta and gamma radiation ;	Ignore “radioactive”, “decay” ignore idea of remaining the same element for ever	1
(b)	i	210 – 84 OR 126		1
	ii	ideas that proton number increases by 1; neutron number decreases by 1;	allow a calculation / nuclear equation Ignore discussion of “number of nucleons”	2
	iii	beta decay	allow β or β^- or β^+	1
(c)	Any two of idea that gamma is not a particle; e.g. gamma rays have no (rest) mass gamma rays do not have a proton number gamma rays do not contain any protons or neutrons gamma rays are electromagnetic radiation OR energy; no particles are lost (from the nucleus) when a gamma ray is emitted;	Allow photons	2	
Total			9	

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publication.orders@edexcel.com

Order Code UG036657 Summer 2013

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
.....



Llywodraeth Cynulliad Cymru
Welsh Assembly Government

