



22127102



**FURTHER MATHEMATICS
STANDARD LEVEL
PAPER 2**

Monday 7 May 2012 (morning)

2 hours

INSTRUCTIONS TO CANDIDATES

- Do not open this examination paper until instructed to do so.
- Answer all questions.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A graphic display calculator is required for this paper.
- A clean copy of the **Mathematics HL and Further Mathematics SL information booklet** is required for this paper.
- The maximum mark for this examination paper is [120 marks].

Please start each question on a new page. Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, e.g. if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

1. [Total mark: 23]

Part A [Maximum mark: 11]

The area of an equilateral triangle is 1 cm^2 . Determine the area of

- (a) the circumscribed circle; [8 marks]
- (b) the inscribed circle. [3 marks]

Part B [Maximum mark: 12]

The points **A**, **B** have coordinates $(1, 0)$, $(0, 1)$ respectively. The point $P(x, y)$ moves in such a way that $AP = kBP$ where $k \in \mathbb{R}^+$.

- (a) When $k = 1$, show that the locus of P is a straight line. [3 marks]
- (b) When $k \neq 1$, the locus of P is a circle.
 - (i) Find, in terms of k , the coordinates of C , the centre of this circle.
 - (ii) Find the equation of the locus of C as k varies. [9 marks]

2. [Total mark: 25]

Part A [Maximum mark: 16]

The graph H has the following adjacency matrix.

$$\begin{array}{c}
 \begin{array}{ccccccc}
 & \text{A} & \text{B} & \text{C} & \text{D} & \text{E} & \text{F} & \text{G} \\
 \text{A} & \left(\begin{array}{ccccccc}
 0 & 1 & 0 & 0 & 0 & 0 & 1 \\
 1 & 0 & 1 & 1 & 0 & 1 & 0 \\
 0 & 1 & 0 & 0 & 1 & 0 & 0 \\
 0 & 1 & 0 & 0 & 1 & 0 & 0 \\
 0 & 0 & 1 & 1 & 0 & 0 & 0 \\
 0 & 1 & 0 & 0 & 0 & 0 & 1 \\
 1 & 0 & 0 & 0 & 0 & 1 & 0
 \end{array} \right)
 \end{array}
 \end{array}$$

- (a) (i) Show that H is bipartite.
- (ii) Draw H as a planar graph. [3 marks]
- (b) (i) Explain what feature of H guarantees that it has an Eulerian circuit.
- (ii) Write down an Eulerian circuit in H . [3 marks]
- (c) (i) Find the number of different walks of length five joining A to B.
- (ii) Determine how many of these walks go through vertex F after passing along two edges. [6 marks]
- (d) Find the maximum number of extra edges that can be added to H while keeping it simple, planar and bipartite. [4 marks]

Part B [Maximum mark: 9]

- (a) Find the smallest positive integer m such that $3^m \equiv 1 \pmod{22}$. [2 marks]
- (b) Given that $3^{49} \equiv n \pmod{22}$ where $0 \leq n \leq 21$, find the value of n . [4 marks]
- (c) Solve the equation $3^x \equiv 5 \pmod{22}$. [3 marks]

3. [Maximum mark: 29]

(a) (i) Show that $\frac{d}{d\theta}(\sec\theta \tan\theta + \ln(\sec\theta + \tan\theta)) = 2\sec^3\theta$.

(ii) Hence write down $\int \sec^3\theta d\theta$. [5 marks]

(b) Consider the differential equation $(1+x^2)\frac{dy}{dx} + xy = 1+x^2$ given that $y=1$ when $x=0$.

(i) Use Euler's method with a step length of 0.1 to find an approximate value for y when $x=0.3$.

(ii) Find an integrating factor for determining the exact solution of the differential equation.

(iii) Find the solution of the equation in the form $y = f(x)$.

(iv) To how many significant figures does the approximation found in part (i) agree with the exact value of y when $x=0.3$? [24 marks]

4. [Total mark: 25]

Part A [Maximum mark: 12]

The function $f: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ is defined by $X \mapsto AX$, where $X = \begin{bmatrix} x \\ y \end{bmatrix}$ and

$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ where a, b, c, d are all non-zero.

(a) Show that f is a bijection if A is non-singular. [7 marks]

(b) Suppose now that A is singular.

(i) Write down the relationship between a, b, c, d .

(ii) Deduce that the second row of A is a multiple of the first row of A .

(iii) Hence show that f is not a bijection. [5 marks]

(This question continues on the following page)

(Question 4 continued)

Part B [Maximum mark: 13]

Consider the group $\{S, +_m\}$ where $S = \{0, 1, 2, \dots, m-1\}$, $m \in \mathbb{N}$, $m \geq 3$ and $+_m$ denotes addition modulo m .

- (a) Show that $\{S, +_m\}$ is cyclic for all m . [3 marks]
- (b) Given that m is prime,
- (i) explain why all elements except the identity are generators of $\{S, +_m\}$;
 - (ii) find the inverse of x , where x is any element of $\{S, +_m\}$ apart from the identity;
 - (iii) determine the number of sets of two distinct elements where each element is the inverse of the other. [7 marks]
- (c) Suppose now that $m = ab$ where a, b are unequal prime numbers. Show that $\{S, +_m\}$ has two proper subgroups and identify them. [3 marks]

5. [Maximum mark: 18]

- (a) The continuous random variable X takes values only in the interval $[a, b]$ and F denotes its cumulative distribution function. Using integration by parts, show that:

$$E(X) = b - \int_a^b F(x) dx. \quad [4 \text{ marks}]$$

- (b) The continuous random variable Y has probability density function f given by:

$$f(y) = \cos y, \quad 0 \leq y \leq \frac{\pi}{2}$$

$$f(y) = 0, \quad \text{elsewhere.}$$

- (i) Obtain an expression for the cumulative distribution function of Y , valid for $0 \leq y \leq \frac{\pi}{2}$. Use the result in (a) to determine $E(Y)$.
- (ii) The random variable U is defined by $U = Y^n$, where $n \in \mathbb{Z}^+$. Obtain an expression for the cumulative distribution function of U valid for $0 \leq u \leq \left(\frac{\pi}{2}\right)^n$.
- (iii) The medians of U and Y are denoted respectively by m_u and m_y . Show that $m_u = m_y^n$.

[14 marks]