



# **MARKSCHEME**

**May 2014**

**BIOLOGY**

**Higher Level**

**Paper 3**

11 pages

**Option D — Evolution**

1. (a) both show a (similar) range of life spans;  
both have a peak at 12–14 hours;  
SFS flies have higher mean life span than NFS flies / vice versa;  
(some) SFS flies live longer than 16 hours, NFS flies do not;  
(some) NFS flies live less than 10 hours, SFS flies do not; *[3 max]*
- (b) SFS has the drier climate as SFS flies live longer (when subjected to drought stress);  
both SFS and NFS have similar climates as both SFS and NFS flies have peak at 12–14 hours; *[1 max]*
- (c) NFS and SFS flies have different gene pools;  
abiotic differences between the two slopes create different selection pressures;  
genes/alleles for advantageous characteristics passed on from one generation to the next/become genetically different over time;  
may not mate / reproductive barriers/isolation leads to speciation;  
abiotic/geographical differences may not be sufficient to cause reproductive barrier;  
the gene pools may not be separated if flies can fly between sides of canyon/mix at base of canyon; *[3 max]*

2. (a) (i) the time taken for radioactivity (of a radioisotope) to fall to half of its original level / for half of the atoms of the isotope to decay [1]

(ii)  $^{40}\text{K}$  decays into  $^{40}\text{Ar}$ ;  
 ratio/proportion of  $^{40}\text{K}$  to  $^{40}\text{Ar}$  indicates the age of the rock/fossil;  
 half-life of  $^{40}\text{K}$  is 1250 million/1.25 billion years;  
 $^{40}\text{K}$  can (only) be used to date very old samples/older than 100 000 years; [2 max]  
*Do not accept if make reference to age less than 100 000 years.*

(b) (i)  $200 - 168 = 32$  flies had short wings  
 $q^2 = 32 / 200 = 0.16$   
 $q = 0.4$   
 $p = 0.6$   
 $2pq = 0.48$ ; } *Award [1] for correct working.  
 (Looking for the logic of the working)*

percentage of heterozygotes = 48 %; [2]  
*Award [1] for correct answer.*

(ii) large population/random mating/no selection/no migration/no mutation [1]  
*Award [1] for any **two** correct answers. Mark the first two answers given.*

(c) a group of organisms consisting of all descendants from a common ancestor [1]

3. eukaryotic cells contain mitochondria/chloroplasts that are not found in prokaryotic cells;  
 organelles evolved from (independent/free living) prokaryotes that were taken into/engulfed by larger cells by endocytosis;  
 these cells were not digested/survived in mutualistic/symbiotic relationship;  
 they continued to carry out respiration/photosynthesis;  
 mitochondria/chloroplasts grow and divide like cells;  
 mitochondria/chloroplasts have a loop of/naked DNA like prokaryotes;  
 mitochondria/chloroplasts have 70S ribosomes like prokaryotes to synthesise proteins;  
 mitochondria/chloroplasts have double membranes expected when taken into a vesicle by endocytosis;  
 thylakoids (in chloroplasts) similar to structures containing chlorophyll in prokaryotes; [6 max]

**Option E — Neurobiology and behaviour**

4. (a)  $0.16 \text{ mg min}^{-1}$  (units needed) [1]  
Allow answers in the range 0.15–0.17.

(b) *hypothesis supported:*  
general increase in food collection over the twelve day period;  
large increases day 1–2/day 6–7/day 10–11;

*hypothesis not supported:*  
most food collected on day 7;  
on some days there are declines / days 3–5/8–10 no increase overall / large decline day 7–8;  
large error bars so data very variable/much overlap between data points;  
only 38 bees in study/sample size too small for drawing valid conclusions; [3 max]

(c) (weather) variations in temperature/some days warm, some days cold/rainfall/wet days, dry days / in wind speed;  
flowers produce more nectar on some days than on others / different flowers open on different days / more flowers open on some days / flowers different distances away so time spent travelling differs;  
bees needed for other duties – defending colony/swarming/ventilating colony;  
harvesting behaviour develops after day 1 as foragers find flowers/communicate; [2 max]

5. (a) sound waves/vibrations in air cause ear drum/tympanic membrane to vibrate;  
vibrations amplified by middle ear bones/ossicles/malleus, incus, stapes;  
causes oval window/fluid in cochlea to vibrate;  
stimulates mechanoreceptors/hair cells;  
auditory nerve passes nerve impulse to brain; [3 max]

(b) (i)

Rods	Cones
function well in dim light / more sensitive to low light	function well in bright light;
absorb all wavelengths of visible light / not responsible for colour vision	sensitive to red, green or blue wavelengths / responsible for colour vision;
poor visual acuity / impulses from several rods pass to a single neuron in the optic nerve	good visual acuity / impulses from a single cone pass to a single neuron in the optic nerve;

Do not accept “rods detect black and white images”.

(ii) (bright) light detected by photoreceptors/rods/cones in retina of eye;  
(sensory) impulses/messages sent to brain along optic nerve;  
medulla oblongata/brain stem processes impulses;  
(motor) impulses/messages sent to (circular) muscles of iris (which contract); pupils constrict; [3 max]

6. development of bird song has both innate and learned components;  
when birds hatch they may possess a basic song (crude template);  
species specific (so inherited/innate/genetic);  
after hatching young birds hear songs from adults of their species;  
(so learn as) mimic/memorise/modify the basic template;  
song does not develop properly if the young bird does not hear other members of its  
species singing;  
some birds have a sensitive period of development when they learn;  
later young birds practise what they have heard;  
song becomes modified/improved to form mature adult song/ song perfected when birds  
become sexually mature;

**[6 max]**

*Marks can be awarded if specific named example is used containing the points above.*

**Option F — Microbes and biotechnology**

7. (a) *cases of malaria: 150 (accept answers in the range 145–155)*  
*monthly rainfall: 165 mm (accept answers in the range 160–170)* **[1]**  
*Both needed for [1].*
- (b) both graphs show fluctuating patterns;  
 malaria rises/falls later than/after rainfall;  
 malaria peaks in July, rainfall peaks in May/Apr–May;  
 June to November – little variation in rainfall, decline in cases of malaria;  
 both decline August to December / during Mar–Apr malaria declines while rain  
 fall increases;  
 both show lowest value in December; **[2 max]**
- (c) *hypothesis supported:*  
 increased rainfall is followed by increased malaria / strong positive correlation  
 between rainfall and number of (future) cases of malaria;  
 rainfall provides pools for mosquito larvae to survive (when adults emerge later  
 they spread the disease);
- hypothesis not supported:*  
 very large values of standard deviation suggests data is unreliable / malaria data  
 less reliable than rainfall;  
 correlation between malaria cases and rainfall does not prove causation; **[2 max]**
- (d) temperature fluctuations;  
 cycles in breeding and populations of mosquitos;  
 cycles in populations of malaria parasites in human hosts;  
 malaria only identified when symptoms show;  
 immigration of infected workers to tea plantations;  
 seasonal spraying of mosquitos/insecticide (causes population decline); **[2 max]**

8. (a) (i) symptom;  
transmission;  
treatment;

[3]

*eg Salmonella food poisoning*

*symptoms: diarrhoea/fever/abdominal cramps/Reiter's syndrome.*

*transmission: by contact after poor personal hygiene/eating contaminated food (not properly cooked)/transferred from faeces of pets/from reptiles/poor kitchen hygiene/raw eggs/unpasteurised milk.*

*treatment: rehydration/lots of drinking/intravenous fluids/antibiotics.*

*Award [1 max] for each symptom, transmission and treatment.*

*Award [2 max] if candidate does not mention the type of food poisoning.*

*"Bacterial food poisoning" is too vague.*

*Example given must be verifiable.*

- (ii) *acids:*  
low pH/pickling/preservation in vinegar restricts growth of microbes/denatures enzymes;  
*sugar:*  
sugar (added to food) dehydrates microbes so it restricts growth / microbes lose water by osmosis;

[2]

(b)

<b>Intracellular (<i>Chlamydia</i>)</b>	<b>Extracellular (<i>Streptococcus</i>)</b>
lives inside cells of host	lives outside host cells;
does not produce toxins/no toxins to irritate tissues/does not damage cells	produces toxins/damages cells;
host may not be aware of infection/asymptomatic	produces symptoms (sore throat) so host aware of infection;
not targeted by immune system / (usually) long-term infection	targeted by immune system / (usually) short-term infection;
sexually transmitted disease / infectious conjunctivitis	respiratory infection / pneumonia / skin infections;

[2 max]

*Answers do not need to be in table format.*

9. prion hypothesis has protein as infecting agent;  
prions are abnormal/misfolded forms of proteins (already present);  
name of prion protein is called major prion protein/protease-resistant protein/PrP<sup>SC</sup>/CD230;  
the (infecting) protein is found in the brain;  
prions can cause normal proteins to change (to abnormal/misfolded) shape;  
misfolded proteins form chains/agglutinate/accumulate;  
cause chain reaction/positive feedback creating more and more abnormal proteins;  
chains interfere with normal cellular functions/cause disease symptoms;  
prion hypothesis supported as no foreign/viral/bacterial DNA/RNA found;  
named example of prion disease *eg* scrapie/CJD/BSE/kuru;

[6 max]

**Option G — Ecology and conservation**

10. (a) 2006 [1]
- (b) increases steadily from 1998 to 2002 and plateaus between 2002 and 2006;  
overall increasing trend / lowest percentage in 1998 and highest in 2006; [1 max]
- (c) fledging success is always greater than breeding success;  
show opposite trends before 2002; (*accept a description*)  
follow (closely) similar trends after 2002; (*accept a description*)  
maximum difference (in percentage) in 1998;  
difference remains smallest between 2002 and 2006; [2 max]
- (d) many of the eggs laid do not hatch but those that do hatch fledge successfully [1]
- (e) eggs may have been laid late in the breeding season so warmer temperatures /  
shorter time for parental care (leading to low fledging success);  
predation of parents/chicks;  
weather conditions at time of fledging may have been unusually harsh;  
food sources may have been reduced; [2 max]
11. (a) (i) temperature;  
water;  
breeding sites;  
food supply;  
territory;  
predators / parasites / pathogens; [2 max]
- (ii) only one species can occupy a niche indefinitely;  
more than one species results in competition for breeding sites/food/other  
resource;  
one species will disappear from the ecosystem/be excluded; [2 max]
- (b) sufficiently large sample size/area of study for both capture and recapture  
samples;  
appropriate time interval between capture and recapture;  
marking techniques must be suitable for the animal species being studied /  
marking must not directly harm/increase chance of predation/bias results;  
 $\frac{n_1 \times n_2}{n_3}$  gives estimate of population size;
- |  |   |   |
|--|---|---|
| $n_1$ = number captured in first sample<br>$n_2$ = number captured in second sample<br>$n_3$ = number in second sample that are marked | } | <i>All three needed for<br/>the mark;</i> |
|--|---|---|
- [3 max]**



12. alien species is one introduced (by humans) to an area in which it does not naturally occur;  
inter-specific competition – competition between different species for resources/ food/light/space/other valid answers;  
valid example stated *eg* in the UK the larger grey squirrel is better adapted to utilize food resources than the native red squirrel / *eg* *Salvinia molesta*/floating fern grows very rapidly over the surface of tropical lakes eliminating native plant species;  
**[2 max]**

predation – one species feeding on another;  
valid example stated *eg* lampreys in St. Lawrence Seaway depleting stocks of lake trout and whitefish / *eg* rats (*Rattus rattus*) introduced onto islands in New Zealand fed on eggs/young birds/adults of native species (which were not behaviourally able to resist them); **[2 max]**

species extinction – one species causing another to become extinct;  
valid example stated *eg* many species of cichlids in Lake Victoria extinct after introduction of Nile perch (introduced to increase fish population) ; **[2 max]**

biological control of pest species – species introduced deliberately to control a pest;  
valid example stated *eg* purple loosestrife in USA and Canada invades wetlands and displaces native species / *eg* *Salvinia* weevil introduced to feed on *Salvinia*/floating fern that has damaged many lakes in the (sub) tropics; **[2 max]**

**[6 max]**

**Option H — Further human physiology**

13. (a)  $510 - 90 = 420$  (beats  $\text{min}^{-1}$ ) (*accept 420/-420/decrease of 420*) [1]  
*Accept answers in the range 400–440.*
- (b) during diving heart rate decreases while arterial blood pressure increases;  
 swimming causes little/no change from control in both heart rate and arterial blood pressure;  
 diving produces greater change than swimming in heart rate and arterial blood pressure;  
 little/no differences between rats diving voluntarily or submerged involuntarily; [2 max]  
*Award [1] if candidate describes swimming and diving for heart rate and then swimming and diving for arterial bp correctly.*
- (c) diving rats hold their breath while swimming rats do not;  
 so heart rate decreases/peripheral blood vessels constrict in diving/submerged rats;  
 swimming rats have no need for the diving response so little/no change from control;  
 constriction of blood vessels in diving rats raises arterial blood pressure;  
 diving response conserves oxygen (for essential functions); [2 max]
- (d) heat loss (from skin) is greater in cold water therefore vasoconstriction is greater / vice versa;  
 diving response slows heart rate so less blood flows to skin so less heat lost in cold water;  
 increased vasoconstriction increases arterial blood pressure;  
 vasoconstriction/increased blood pressure helps to maintain core temperature in cold water;  
 decrease in body temperature causes heart rate to slow; [2 max]
14. (a) (i) *steroid hormone:* oestrogen/progesterone/testosterone;  
*protein hormone:* ADH (vasopressin)/insulin/TRH;  
 other valid examples; [2 max]
- (ii) presence of food/stretching of stomach stimulates endocrine glands (in stomach wall);  
 gastrin secreted;  
 gastrin/hormone stimulates increased secretion of HCl; [2 max]
- (b) increased  $\text{CO}_2$  in blood / lower blood pH;  
 detected by chemosensors/chemoreceptors/receptors in aortic/carotid arteries;  
 nerve impulses/messages sent to breathing centre of medulla/brain;  
 breathing centres/medulla also monitor blood pH/ $\text{CO}_2$ ;  
 (more) nerve impulses/messages sent to diaphragm and intercostal muscles (causing increased muscle contraction); [3 max]

15. erythrocytes have a lifespan of approximately 120 days;  
erythrocytes rupture releasing hemoglobin into bloodstream;  
phagocytosis removes hemoglobin/erythrocytes;  
by Kupffer cells/macrophages;  
hemoglobin split into heme and globin;  
globins hydrolysed into amino acids;  
iron removed from heme;  
(heme) converted into (biliverdin then) bilirubin (bile pigment);  
iron stored/released into blood;

*[6 max]*