



# **MARKSCHEME**

**November 2014**

**MATHEMATICS  
DISCRETE MATHEMATICS**

**Higher Level**

**Paper 3**

11 pages

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Assessment Centre.*

## Instructions to Examiners

### Abbreviations

- M** Marks awarded for attempting to use a correct **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

### Using the markscheme

#### 1 General

Mark according to RM™ Assessor instructions and the document “**Mathematics HL: Guidance for e-marking November 2014**”. It is essential that you read this document before you start marking. In particular, please note the following.

Marks must be recorded using the annotation stamps. Please check that you are entering marks for the right question.

- If a part is **completely correct**, (and gains all the “must be seen” marks), use the ticks with numbers to stamp full marks.
- If a part is completely wrong, stamp **A0** by the final answer.
- If a part gains anything else, it **must** be recorded using **all** the annotations.
- All the marks will be added and recorded by RM™ Assessor.

#### 2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **AI**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, for example, **MIAI**, this usually means **MI** for an **attempt** to use an appropriate method (for example, substitution into a formula) and **AI** for using the **correct** values.
- Where the markscheme specifies **(M2)**, **N3**, etc, do **not** split the marks.
- Once a correct answer to a question or part-question is seen, ignore further working.

### 3 ***N* marks**

Award *N* marks for **correct** answers where there is **no** working.

- Do **not** award a mixture of *N* and other marks.
- There may be fewer *N* marks available than the total of *M*, *A* and *R* marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

### 4 **Implied marks**

Implied marks appear in **brackets, for example, (M1)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

### 5 **Follow through marks**

Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (for example,  $\sin \theta = 1.5$ ), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

### 6 **Mis-read**

If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). A candidate should be penalized only once for a particular mis-read. Use the **MR** stamp to indicate that this has been a mis-read. Then deduct the first of the marks to be awarded, even if this is an **M** mark, but award all others so that the candidate only loses one mark.

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (for example,  $\sin \theta = 1.5$ ), do not award the mark(s) for the final answer(s).

### 7 **Discretionary marks (d)**

An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. In such cases the annotation **DM** should be used and a brief **note** written next to the mark explaining this decision.

## 8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

## 9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

**Example:** for differentiating  $f(x) = 2 \sin(5x - 3)$ , the markscheme gives:

$$f'(x) = (2 \cos(5x - 3)) 5 \quad (= 10 \cos(5x - 3)) \quad \mathbf{A1}$$

Award **A1** for  $(2 \cos(5x - 3)) 5$ , even if  $10 \cos(5x - 3)$  is not seen.

## 10 Accuracy of Answers

Candidates should **NO LONGER** be penalized for an accuracy error (**AP**).

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy. When this is not specified in the question, all numerical answers should be given exactly or correct to three significant figures. Please check work carefully for **FT**.

## 11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

**12 Calculators**

*A GDC is required for paper 3, but calculators with symbolic manipulation features (for example, TI-89) are not allowed.*

**Calculator notation**

The Mathematics HL guide says:

*Students must always use correct mathematical notation, not calculator notation.*

Do **not** accept final answers written using calculator notation. However, do not penalize the use of calculator notation in the working.

**13 More than one solution**

*Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise.*

1. (a) 240, 1020, 3120 A2

**Note:** Award **A2** for three correct answers, **A1** for two correct answers.

*[2 marks]*

(b) (i)  $1020 = 240 \times 4 + 60$  (M1)

$240 = 60 \times 4$

$\text{gcd}(1020, 240) = 60$  A1

(ii)  $3120 = 1020 \times 3 + 60$  (M1)

$1020 = 60 \times 17$

$\text{gcd}(1020, 3120) = 60$  A1

*[4 marks]*

**Note:** Must be done by Euclid's algorithm.

(c) by Fermat's little theorem with  $p = 5$  R1

$n^5 \equiv n \pmod{5}$

so 5 divides  $f(n)$

*[1 mark]*

(d)  $f(n) = n(n^2 - 1)(n^2 + 1) = n(n - 1)(n + 1)(n^2 + 1)$  (A1)A1

$n - 1, n, n + 1$  are consecutive integers and so contain a multiple of 2 and 3

R1R1

**Note:** Award **R1** for justification of 2 and **R1** for justification of 3.

And therefore  $f(n)$  is a multiple of 6. AG

*[4 marks]*

(e) from (c) and (d)  $f(n)$  is always divisible by 30 R1

considering the factorization, it is divisible by 60 when  $n$  is an odd number A1

or when  $n$  is a multiple of 4 A1

**Note:** Accept answers such as when  $n$  is not congruent to 2 (mod 4).

*[3 marks]*

**Total [14 marks]**

2. (a) let there be  $v$  vertices in the graph; because the graph is simple the degree of each vertex is  $\leq v - 1$  *AI*  
the degree of each vertex is  $\geq 1$  *AI*  
there are therefore  $v - 1$  possible values for the degree of each vertex *AI*  
given there are  $v$  vertices by the pigeon-hole principle there must be at least two with the same degree *RI*  
*[4 marks]*
- (b) consider a graph in which the people at the meeting are represented by the vertices and two vertices are connected if the two people shake hands *MI*  
the graph is simple as no-one shakes hands with the same person more than once (nor can someone shake hands with themselves) *AI*  
  
every vertex is connected to at least one other vertex as everyone shakes at least one hand *AI*  
  
the degree of each vertex is the number of handshakes so by the proof above there must be at least two who shake the same number of hands *RI*  
*[4 marks]*
- Note:** Accept answers starting afresh rather than quoting part (a).
- (c) (the handshaking lemma tells us that) the sum of the degrees of the vertices must be an even number *AI*  
the degree of each vertex would be 9 and  $9 \times 17$  is an odd number (giving a contradiction) *AI*  
*[2 marks]*
- Total [10 marks]**



3. (a)

|   | A | B   | C  | D   | E  | F   | G  | H  | I  | J  |
|---|---|-----|----|-----|----|-----|----|----|----|----|
| A | 0 | ←10 | 11 | 18  |    |     |    |    |    |    |
| B |   | 10  | ←  | ←17 | 21 | 23  |    |    |    |    |
| C |   |     | 11 | ↑   |    |     |    |    |    |    |
| D |   |     |    | 17  | ←  | ←24 |    | 22 |    |    |
| E |   |     |    |     | 21 |     |    | ↑  | 30 |    |
| H |   |     |    |     |    |     |    | 22 | ←  | 27 |
| F |   |     |    |     |    | 23  |    |    |    | 29 |
| G |   |     |    |     |    |     | 24 |    |    | 28 |
| J |   |     |    |     |    |     |    |    |    | 27 |

*MIAIAIAI*

(*MI* for an attempt at Dijkstra's)  
 (*AI* for value of D = 17)  
 (*AI* for value of H = 22)  
 (*AI* for value of G = 24)  
 route is ABDHJ  
 cost is \$27

(*MI*)*AI*  
*AI*

[7 marks]

**Note:** Accept other layouts.

(b) there are 4 odd vertices A, D, F and J  
 these can be joined up in 3 ways with the following extra costs  
 AD and FJ  $17 + 13 = 30$   
 AF and DJ  $23 + 10 = 33$   
 AJ and DF  $27 + 12 = 39$

*AI*

*MIAIAI*

**Note:** Award *MI* for an attempt to find different routes.  
 Award *AI* for correct values for all three costs *AI* for one correct.

need to repeat AB, BD, FG and GJ  
 total cost is  $139 + 30 = \$169$

*AI*  
*AI*

[6 marks]

Total [13 marks]

4. (a) **METHOD 1**

listing 9, 20, 31, ... and 1, 6, 11, 16, 21, 26, 31, ...

*MI*

one solution is 31

*(AI)*

by the Chinese remainder theorem the full solution is

$$x \equiv 31 \pmod{55}$$

*AI*

*N2*

**METHOD 2**

$$x \equiv 9 \pmod{11} \Rightarrow x = 9 + 11t$$

*MI*

$$\Rightarrow 9 + 11t \equiv 1 \pmod{5}$$

$$\Rightarrow t \equiv 2 \pmod{5}$$

*AI*

$$\Rightarrow t = 2 + 5s$$

$$\Rightarrow x = 9 + 11(2 + 5s)$$

$$\Rightarrow x = 31 + 55s \left( \Rightarrow x \equiv 31 \pmod{55} \right)$$

*AI*

**Note:** Accept other methods *eg* formula, Diophantine equation.

**Note:** Accept other equivalent answers e.g.  $-79 \pmod{55}$ .

*[3 marks]*

(b)  $41^{82} \equiv 8^{82} \pmod{11}$

by Fermat's little theorem  $8^{10} \equiv 1 \pmod{11}$  (or  $41^{10} \equiv 1 \pmod{11}$ )

*MI*

$$8^{82} \equiv 8^2 \pmod{11}$$

*MI*

$$\equiv 9 \pmod{11}$$

*(AI)*

remainder is 9

*AI*

*[4 marks]*

**Note:** Accept simplifications done without Fermat.

(c)  $41^{82} \equiv 1^{82} \equiv 1 \pmod{5}$

*AI*

so  $41^{82}$  has a remainder 1 when divided by 5 and a remainder 9 when divided by 11

*RI*

hence by part (a) the remainder is 31

*AI*

*[3 marks]*

**Total [10 marks]**

5. (a)  $\frac{1}{2}$  *AI*  
*[1 mark]*

(b) Andy could win the  $n^{\text{th}}$  game by winning the  $n-1^{\text{th}}$  and then winning the  $n^{\text{th}}$  game or by losing the  $n-1^{\text{th}}$  and then winning the  $n^{\text{th}}$  *(M1)*  

$$u_n = \frac{1}{2}u_{n-1} + \frac{1}{4}(1-u_{n-1})$$
 *A1A1M1*

**Note:** Award *AI* for each term and *M1* for addition of two probabilities.

$$u_n = \frac{1}{4}u_{n-1} + \frac{1}{4}$$
 *AG*  
*[4 marks]*

(c) general solution is  $u_n = A\left(\frac{1}{4}\right)^n + p(n)$  *(M1)*

for a particular solution try  $p(n) = b$  *(M1)*

$$b = \frac{1}{4}b + \frac{1}{4}$$
 *(A1)*

$$b = \frac{1}{3}$$

hence  $u_n = A\left(\frac{1}{4}\right)^n + \frac{1}{3}$  *(A1)*

using  $u_1 = \frac{1}{2}$  *M1*

$$\frac{1}{2} = A\left(\frac{1}{4}\right) + \frac{1}{3} \Rightarrow A = \frac{2}{3}$$

hence  $u_n = \frac{2}{3}\left(\frac{1}{4}\right)^n + \frac{1}{3}$  *A1*

**Note:** Accept other valid methods.

*[6 marks]*

(d) for large  $n$   $u_n \approx \frac{1}{3}$  *(M1)A1*

*[2 marks]*

**Total [13 marks]**