

**Further mathematics**  
**Higher level**  
**Paper 2**

Friday 20 May 2016 (morning)

2 hours 30 minutes

---


**Instructions to candidates**

- Do not open this examination paper until instructed to do so.
- Answer all questions.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A graphic display calculator is required for this paper.
- A clean copy of the **mathematics HL and further mathematics HL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[150 marks]**.

Please start each question on a new page. Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working. For example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

1. [Maximum mark: 14]

Consider the following weighted graph.



- (a) Determine whether or not the graph is Eulerian. [2]
- (b) Determine whether or not the graph is Hamiltonian. [2]
- (c) Use Kruskal's algorithm to find a minimum weight spanning tree and state its weight. [6]
- (d) Deduce an upper bound for the total weight of a closed walk of minimum weight which visits every vertex. [2]
- (e) Explain how the result in part (b) can be used to find a different upper bound and state its value. [2]

2. [Maximum mark: 17]

- (a) Use l'Hôpital's rule to show that  $\lim_{x \rightarrow \infty} \frac{x^3}{e^x} = 0$ . [3]

The random variable  $X$  has probability density function given by

$$f(x) = \begin{cases} xe^{-x}, & \text{for } x \geq 0, \\ 0, & \text{otherwise} \end{cases}.$$

- (b) (i) Find  $E(X^2)$ .  
(ii) Show that  $\text{Var}(X) = 2$ . [10]
- (c) State the central limit theorem. [2]

A sample of size 50 is taken from the distribution of  $X$ .

- (d) Find the probability that the sample mean is less than 2.3. [2]

3. [Maximum mark: 15]

A circle  $C$  passes through the point  $(1, 2)$  and has the line  $3x - y = 5$  as the tangent at the point  $(3, 4)$ .

- (a) Find the coordinates of the centre of  $C$  and its radius. [9]
- (b) Write down the equation of  $C$ . [1]
- (c) Find the coordinates of the second point on  $C$  on the chord through  $(1, 2)$  parallel to the tangent at  $(3, 4)$ . [5]

Turn over

## 4. [Maximum mark: 23]

Consider the differential equation  $\frac{dy}{dx} = \frac{x}{y}$ , where  $y \neq 0$ .

- (a) Find the general solution of the differential equation, expressing your answer in the form  $f(x, y) = c$ , where  $c$  is a constant. [3]
- (b) (i) Hence find the particular solution passing through the points  $(1, \pm \sqrt{2})$ .  
(ii) Sketch the graph of your solution and name the type of curve represented. [5]
- (c) (i) Write down the particular solution passing through the points  $(1, \pm 1)$ .  
(ii) Give a geometrical interpretation of this solution in relation to part (b). [3]
- (d) (i) Find the general solution of the differential equation  $\frac{dy}{dx} = \frac{x}{y} + \frac{y}{x}$ , where  $xy \neq 0$ .  
(ii) Find the particular solution passing through the point  $(1, \sqrt{2})$ .  
(iii) Sketch the particular solution.  
(iv) The graph of the solution only contains points with  $|x| > a$ . Find the exact value of  $a$ ,  $a > 0$ . [12]

## 5. [Maximum mark: 19]

(a) The sequence  $\{u_n : n \in \mathbb{Z}^+\}$  satisfies the recurrence relation  $2u_{n+2} - 3u_{n+1} + u_n = 0$ , where  $u_1 = 1$ ,  $u_2 = 2$ .

(i) Find an expression for  $u_n$  in terms of  $n$ .

(ii) Show that the sequence converges, stating the limiting value. [9]

(b) The sequence  $\{v_n : n \in \mathbb{Z}^+\}$  satisfies the recurrence relation  $2v_{n+2} - 3v_{n+1} + v_n = 1$ , where  $v_1 = 1$ ,  $v_2 = 2$ .

Without solving the recurrence relation prove that the sequence diverges. [3]

(c) The sequence  $\{w_n : n \in \mathbb{N}\}$  satisfies the recurrence relation  $w_{n+2} - 2w_{n+1} + 4w_n = 0$ , where  $w_0 = 0$ ,  $w_1 = 2$ .

(i) Find an expression for  $w_n$  in terms of  $n$ .

(ii) Show that  $w_{3n} = 0$  for all  $n \in \mathbb{N}$ . [7]

Turn over

6. [Maximum mark: 19]

Consider the set  $J = \{a + b\sqrt{2} : a, b \in \mathbb{Z}\}$  under the binary operation multiplication.

- (a) Show that  $J$  is closed. [2]
- (b) State the identity in  $J$ . [1]
- (c) Show that
  - (i)  $1 - \sqrt{2}$  has an inverse in  $J$ ;
  - (ii)  $2 + 4\sqrt{2}$  has no inverse in  $J$ . [5]
- (d) Show that the subset,  $G$ , of elements of  $J$  which have inverses, forms a group of infinite order. [7]
- (e) Consider  $a + b\sqrt{2} \in G$ , where  $\gcd(a, b) = 1$ ,
  - (i) Find the inverse of  $a + b\sqrt{2}$ .
  - (ii) Hence show that  $a^2 - 2b^2$  divides exactly into  $a$  and  $b$ .
  - (iii) Deduce that  $a^2 - 2b^2 = \pm 1$ . [4]

7. [Maximum mark: 19]

Consider the functions  $f_n(x) = \sec^n(x)$ ,  $|x| < \frac{\pi}{2}$  and  $g_n(x) = f_n(x)\tan x$ .

(a) Show that

(i) 
$$\frac{df_n(x)}{dx} = ng_n(x);$$

(ii) 
$$\frac{dg_n(x)}{dx} = (n + 1)f_{n+2}(x) - nf_n(x).$$
 [5]

(b) (i) Use these results to show that the Maclaurin series for the function  $f_5(x)$  up to and including the term in  $x^4$  is  $1 + \frac{5}{2}x^2 + \frac{85}{24}x^4$ .

(ii) By considering the general form of its higher derivatives explain briefly why all coefficients in the Maclaurin series for the function  $f_5(x)$  are either positive or zero.

(iii) Hence show that  $\sec^5(0.1) > 1.02535$ . [14]

Turn over

8. [Maximum mark: 24]

The set of all permutations of the list of the integers  $1, 2, 3 \dots n$  is a group,  $S_n$ , under the operation of composition of permutations.

- (a) (i) Show that the order of  $S_n$  is  $n!$ ;
- (ii) List the 6 elements of  $S_3$  in cycle form;
- (iii) Show that  $S_3$  is not Abelian;
- (iv) Deduce that  $S_n$  is not Abelian for  $n \geq 3$ . [9]

(b) Each element of  $S_4$  can be represented by a  $4 \times 4$  matrix. For example, the cycle  $(1\ 2\ 3\ 4)$  is represented by the matrix

$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \text{ acting on the column vector } \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}.$$

- (i) Write down the matrices  $M_1, M_2$  representing the permutations  $(1\ 2), (2\ 3)$ , respectively;
  - (ii) Find  $M_1 M_2$  and state the permutation represented by this matrix;
  - (iii) Find  $\det(M_1), \det(M_2)$  and deduce the value of  $\det(M_1 M_2)$ . [7]
- (c) (i) Use mathematical induction to prove that  $(1\ n)(1\ n-1)(1\ n-2)\dots(1\ 2) = (1\ 2\ 3\dots n)$   $n \in \mathbb{Z}^+, n > 1$ .
- (ii) Deduce that every permutation can be written as a product of cycles of length 2. [8]