

# Markscheme

November 2016

Chemistry

Higher level

Paper 2

20 pages

This markscheme is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Assessment Centre.

Question			Answers	Notes	Total
1.	a	i	$\llcorner K_c = \frac{[\text{HOCH}_2\text{CH}_2\text{OH}]}{[\text{CO}]^2 \times [\text{H}_2]^3} \checkmark$		1
1.	a	ii	<i>Position of equilibrium:</i> moves to right <b>OR</b> favours product $\checkmark$  <i>K<sub>c</sub>:</i> no change <b>OR</b> is a constant at constant temperature $\checkmark$		2
1.	a	iii	<i>Bonds broken:</i> $2\text{C}\equiv\text{O} + 3(\text{H}-\text{H}) / 2(1077 \text{ kJ mol}^{-1}) + 3(436 \text{ kJ mol}^{-1}) / 3462 \llcorner \text{kJ} \checkmark$  <i>Bonds formed:</i> $2(\text{C}-\text{O}) + 2(\text{O}-\text{H}) + 4(\text{C}-\text{H}) + (\text{C}-\text{C}) / 2(358 \text{ kJ mol}^{-1}) + 2(463 \text{ kJ mol}^{-1}) + 4(414 \text{ kJ mol}^{-1}) + 346 \text{ kJ mol}^{-1} / 3644 \llcorner \text{kJ} \checkmark$  $\llcorner \text{Enthalpy change} = \text{bonds broken} - \text{bonds formed} = 3462 \text{ kJ} - 3644 \text{ kJ} = \llcorner -182 \llcorner \text{kJ} \checkmark$	<i>Award [3] for correct final answer.            Award [2 max] for <math>\llcorner + \llcorner 182 \llcorner \text{kJ}</math>.</i>	3
1.	b	i	$\llcorner \Delta H = \sum \Delta H_f \text{ products} - \sum \Delta H_f \text{ reactants} = -454.8 \text{ kJ mol}^{-1} - 2(-110.5 \text{ kJ mol}^{-1}) = \llcorner -233.8 \llcorner \text{kJ} \checkmark$		1
1.	b	ii	in (a)(iii) gas is formed and in (b)(i) liquid is formed <b>OR</b> products are in different states <b>OR</b> conversion of gas to liquid is exothermic <b>OR</b> conversion of liquid to gas is endothermic <b>OR</b> enthalpy of vapourisation needs to be taken into account $\checkmark$	<i>Accept product is <math>\llcorner \text{now} \llcorner</math> a liquid.             Accept answers referring to bond enthalpies being means/averages.</i>	1

(continued)

(Question 1 continued)

Question			Answers	Notes	Total
1.	b	iii	« $\Delta S$ is negative because five mols of» gases becomes «one mol of» liquid <b>OR</b> increase in complexity of product «compared to reactants» <b>OR</b> product more ordered «than reactants» ✓	Accept “fewer moles of <u>gas</u> ” but not “fewer molecules”.	1
1.	b	iv	$\Delta S = \left(\frac{-620.1}{1000}\right) \text{ «kJ K}^{-1}\text{» } \checkmark$ $\Delta G = -233.8 \text{ kJ} - (298 \text{ K} \left(\frac{-620.1}{1000}\right) \text{ kJ K}^{-1}) = -49.0 \text{ «kJ» } \checkmark$	Award [2] for correct final answer. Award [1 max] for «+» $185 \times 10^3$ . If $-244.0 \text{ kJ}$ used, answer is: $\Delta G = -244.0 \text{ kJ} - (298 \text{ K} \left(\frac{-620.1}{1000}\right) \text{ kJ K}^{-1}) =$ $-59.2 \text{ «kJ»}$ Award [2] for correct final answer.	2
1.	b	v	increasing T makes $\Delta G$ larger/more positive/less negative <b>OR</b> $-T\Delta S$ will increase ✓		1
1.	c		Ethene: $-2 \checkmark$  Ethane-1,2-diol: $-1 \checkmark$	Do not accept 2-, 1- respectively.	2

(continued)

(Question 1 continued)

Question		Answers	Notes	Total									
1.	d	ethane-1,2-diol can hydrogen bond to other molecules «and ethene cannot» <b>OR</b> ethane-1,2-diol has «significantly» greater van der Waals forces ✓  hydrogen bonding is «significantly» stronger than other intermolecular forces ✓	Accept converse arguments. Award [0] if answer implies covalent bonds are broken.	2									
1.	e	acidified «potassium» dichromate«(VI)» / $H^+$ <b>AND</b> $K_2Cr_2O_7$ / $H^+$ <b>AND</b> $Cr_2O_7^{2-}$ <b>OR</b> «acidified potassium» manganate(VII) / « $H^+$ » $KMnO_4$ / « $H^+$ » $MnO_4^-$ ✓	Accept $H_2SO_4$ or $H_3PO_4$ for $H^+$ . Accept “permanganate” for “manganate(VII)”.	1									
1.	f	<table border="1"> <thead> <tr> <th></th> <th>Number of signals</th> <th>Splitting pattern</th> </tr> </thead> <tbody> <tr> <td>Ethanedioic acid:</td> <td>1</td> <td><b>AND</b> singlet ✓</td> </tr> <tr> <td>Ethane-1,2-diol:</td> <td>2 ✓</td> <td>Not required</td> </tr> </tbody> </table>		Number of signals	Splitting pattern	Ethanedioic acid:	1	<b>AND</b> singlet ✓	Ethane-1,2-diol:	2 ✓	Not required	Accept “none/no splitting” for singlet.	2
	Number of signals	Splitting pattern											
Ethanedioic acid:	1	<b>AND</b> singlet ✓											
Ethane-1,2-diol:	2 ✓	Not required											
2.	a	<i>Weak acid:</i> partially dissociated/ionized «in solution/water» <b>AND</b> <i>Strong acid:</i> «assumed to be almost» completely/100% dissociated/ionized «in solution/water» ✓	Accept answers relating to pH, conductivity, reactivity if solutions of equal concentrations stated.	1									

(continued)

(Question 2 continued)

Question			Answers	Notes	Total
2.	b		«log scale» reduces a wide range of numbers to a small range <b>OR</b> simple/easy to use <b>OR</b> converts exponential expressions into a linear scale/simple numbers ✓	<i>Do not accept "easy for calculations".</i>	1
2.	c	i	phenolphthalein <b>OR</b> phenol red ✓		1
2.	c	ii	« $n(\text{NaOH}) = \left(\frac{14.0}{1000}\right) \text{dm}^3 \times 0.100 \text{mol dm}^{-3} \Rightarrow 1.40 \times 10^{-3}$ «mol» ✓		1
2.	c	iii	« $\frac{1}{2} \times 1.40 \times 10^{-3} \Rightarrow 7.00 \times 10^{-4}$ «mol» ✓		1


(continued)

(Question 2 continued)

Question			Answers	Notes	Total
2.	c	iv	<p><b>ALTERNATIVE 1:</b>            «mass of pure hydrated ethanedioic acid in each titration = <math>7.00 \times 10^{-4} \text{ mol} \times 126.08 \text{ g mol}^{-1} \Rightarrow 0.0883 / 8.83 \times 10^{-2} \text{ «g»} \checkmark</math>            mass of sample in each titration = « <math>\frac{25}{1000} \times 5.00 \text{ g} \Rightarrow 0.125 \text{ «g»} \checkmark</math>            «% purity = <math>\frac{0.0883 \text{ g}}{0.125 \text{ g}} \times 100 \Rightarrow 70.6 \text{ «%»} \checkmark</math></p> <p><b>ALTERNATIVE 2:</b>            «mol of pure hydrated ethanedioic acid in <math>1 \text{ dm}^3</math> solution = <math>7.00 \times 10^{-4} \times \frac{1000}{25} \Rightarrow 2.80 \times 10^{-2} \text{ «mol»} \checkmark</math>            «mass of pure hydrated ethanedioic acid in sample = <math>2.80 \times 10^{-2} \text{ mol} \times 126.08 \text{ g mol}^{-1} \Rightarrow 3.53 \text{ «g»} \checkmark</math>            «% purity = <math>\frac{3.53 \text{ g}}{5.00 \text{ g}} \times 100 \Rightarrow 70.6 \text{ «%»} \checkmark</math></p> <p><b>ALTERNATIVE 3:</b>            mol of hydrated ethanedioic acid (assuming sample to be pure) =  <math>\frac{5.00 \text{ g}}{126.08 \text{ g mol}^{-1}} = 0.03966 \text{ «mol»} \checkmark</math>            actual amount of hydrated ethanedioic acid =            «<math>7.00 \times 10^{-4} \times \frac{1000}{25} \Rightarrow 2.80 \times 10^{-2} \text{ «mol»} \checkmark</math>            «% purity = <math>\frac{2.80 \times 10^{-2}}{0.03966} \times 100 \Rightarrow 70.6 \text{ «%»} \checkmark</math></p>	<p><i>Award suitable part marks for alternative methods.</i></p> <p><i>Award [3] for correct final answer.</i></p> <p><i>Award [2 max] for 50.4% if anhydrous ethanedioic acid assumed.</i></p>	<b>3</b>

(continued)

(Question 2 continued)

Question		Answers	Notes	Total
2.	d		<p>Accept single negative charges on two O atoms singly bonded to C. Do not accept resonance structures. Allow any combination of dots/crosses or lines to represent electron pairs.</p>	1
2.	e	<p>electrons delocalized «across the O–C–O system» <b>OR</b> resonance occurs ✓</p> <p>122 «pm» &lt; C–O &lt; 143 «pm» ✓</p>	<p>Accept delocalized <math>\pi</math>-bond(s). No ECF from (d). Accept any answer in range 123 «pm» to 142 «pm». Accept “bond intermediate between single and double bond” or “bond order 1.5”.</p>	2
2.	f	<p>coordinate/dative/covalent bond from O to «transition» metal «ion» <b>OR</b> acts as a Lewis base/nucleophile ✓</p> <p>can occupy two positions <b>OR</b> provide two electron pairs from different «O» atoms <b>OR</b> form two «coordinate/dative/covalent» bonds «with the metal ion» <b>OR</b> chelate «metal/ion» ✓</p>		2

(continued)



Question		Answers	Notes	Total
3.	a	H <sub>2</sub> O <b>AND</b> (l) ✓	Do <b>not</b> accept H <sub>2</sub> O(aq).	1
3.	b	SO <sub>2</sub> (g) is an irritant/causes breathing problems <b>OR</b> SO <sub>2</sub> (g) is poisonous/toxic ✓	Accept SO <sub>2</sub> (g) is acidic but do not accept "causes acid rain". Accept SO <sub>2</sub> (g) is harmful. Accept SO <sub>2</sub> (g) has a foul/pungent smell.	1
3.	c	$n(\text{HCl}) = \llcorner \frac{10.0}{1000} \text{ dm}^3 \times 2.00 \text{ mol dm}^{-3} \Rightarrow 0.0200 / 2.00 \times 10^{-2} \llcorner \llcorner \text{mol} \llcorner \llcorner$ <b>AND</b> $n(\text{Na}_2\text{S}_2\text{O}_3) = \llcorner \frac{50}{1000} \text{ dm}^3 \times 0.150 \text{ mol} \times \text{dm}^{-3} \Rightarrow 0.00750 / 7.50 \times 10^{-3} \llcorner \llcorner \text{mol} \llcorner \llcorner \checkmark$ $0.0200 \llcorner \llcorner \text{mol} \llcorner \llcorner > 0.0150 \llcorner \llcorner \text{mol} \llcorner \llcorner$ <b>OR</b> $2.00 \times 10^{-2} \llcorner \llcorner \text{mol} \llcorner \llcorner > 2 \times 7.50 \times 10^{-3} \llcorner \llcorner \text{mol} \llcorner \llcorner$ <b>OR</b> $\frac{1}{2} \times 2.00 \times 10^{-2} \llcorner \llcorner \text{mol} \llcorner \llcorner > 7.50 \times 10^{-3} \llcorner \llcorner \text{mol} \llcorner \llcorner \checkmark$	Accept answers based on volume of solutions required for complete reaction.  Award <b>[2]</b> for second marking point.  Do <b>not</b> award M2 unless factor of 2 (or half) is used.	2

(continued)

(Question 3 continued)

Question		Answers	Notes	Total
3.	d	<p>five points plotted correctly ✓ best fit line drawn with ruler, going through the origin ✓</p>		2


(continued)

(Question 3 continued)

Question			Answers	Notes	Total
3.	e	i	first order ✓ «because» $[\text{Na}_2\text{S}_2\text{O}_3]$ is «directly» proportional to rate of reaction « $\frac{1}{t}$ » ✓	Do not accept “linear” for M2.	2
3.	e	ii	rate = $k [\text{Na}_2\text{S}_2\text{O}_3][\text{HCl}]$ ✓		1
3.	f		<p> <math>22.5 \times 10^{-3} \text{ «s}^{-1}\text{»} \checkmark</math>  <math>\text{«Time} = \frac{1}{22.5 \times 10^{-3}} \Rightarrow 44.4 \text{ «s»} \checkmark</math> </p>	Award [2] for correct final answer. Accept value based on candidate's graph.  Award M2 as ECF from M1.  Award [1 max] for methods involving taking mean of appropriate pairs of $\frac{1}{t}$ values.  Award [0] for taking mean of pairs of time values.  Award [2] for answers between 42.4 and 46.4 «s».	2

(continued)

(Question 3 continued)

Question			Answers	Notes	Total
3.	g	i	 <p>correctly labelled axes ✓ peak of <math>T_2</math> curve lower <b>AND</b> to the right of <math>T_1</math> curve ✓</p>	<p>Accept "probability «density» / number of particles / N / fraction" on y-axis. Accept "kinetic E/KE/<math>E_k</math>" but not <b>just</b> "Energy/E" on x-axis.</p>	2

(continued)

(Question 3 continued)

Question			Answers	Notes	Total
3.	g	ii	greater proportion of molecules have $E \geq E_a$ or $E > E_a$ <b>OR</b> greater area under curve to the right of the $E_a$ ✓  greater frequency of collisions «between molecules» <b>OR</b> more collisions per unit time/second ✓	Accept more molecules have energy greater than $E_a$ . Do <b>not</b> accept just “particles have greater kinetic energy”.  Accept “rate/chance/probability/likelihood” instead of “frequency”. Accept suitably shaded/annotated diagram.  Do <b>not</b> accept just “more collisions”.	2
3.	h		shorter reaction time so larger «%» error in timing/seeing when mark disappears ✓	Accept cooling of reaction mixture during course of reaction.	1

(continued)



Question		Answers	Notes	Total	
4.	a	${}^{26}_{12}\text{Mg}$ ✓		1	
4.	b	« $A_r$ =>» $\frac{24 \times 78.60 + 25 \times 10.11 + 26 \times 11.29}{100}$ ✓ «= 24.3269 =>» 24.33 ✓	Award [2] for correct final answer. Do <b>not</b> accept data booklet value (24.31).	2	
4.	c	contamination with sodium/other «compounds» ✓		1	
4.	d	i	energy levels are closer together <u>at high energy / high frequency / short wavelength</u> ✓	1	
4.	d	ii	ionisation energy ✓	1	
4.	e	$\text{MgO(s)} + \text{H}_2\text{O(l)} \rightarrow \text{Mg(OH)}_2\text{(s)}$ <b>OR</b> $\text{MgO(s)} + \text{H}_2\text{O(l)} \rightarrow \text{Mg}^{2+}\text{(aq)} + 2\text{OH}^-\text{(aq)}$ ✓	Accept $\rightleftharpoons$ .	1	
4.	f	from basic to acidic ✓ through amphoteric ✓	Accept “alkali/alkaline” for “basic”. Accept “oxides of Na and Mg: basic <b>AND</b> oxide of Al: amphoteric” for M1. Accept “oxides of non-metals/Si to Cl acidic” for M2. Do <b>not</b> accept just “become more acidic”.	2	
4.	g	$\text{Mg}_3\text{N}_2$ ✓	Accept $\text{MgO}_2$ , $\text{Mg(OH)}_2$ , $\text{Mg(NO}_x)_2$ , $\text{MgCO}_3$ .	1	
4.	h	«3-D/giant» regularly repeating arrangement «of ions» <b>OR</b> lattice «of ions» ✓  electrostatic attraction between oppositely charged ions <b>OR</b> electrostatic attraction between $\text{Mg}^{2+}$ and $\text{O}^{2-}$ ions ✓	Accept “giant” for M1 unless “giant covalent” stated.  Do <b>not</b> accept “ionic” without description.	2	
4.	i	i	Anode (positive electrode): $2\text{Cl}^- \rightarrow \text{Cl}_2\text{(g)} + 2\text{e}^-$ ✓  Cathode (negative electrode): $\text{Mg}^{2+} + 2\text{e}^- \rightarrow \text{Mg(l)}$ ✓	Penalize missing/incorrect state symbols at $\text{Cl}_2$ and Mg once only.  Award [1 max] if equations are at wrong electrodes. Accept Mg (g).	2

(continued)

(Question 4 continued)

Question			Answers	Notes	Total
4.	i	ii	reduction ✓		1
4.	i	iii	<p>Anode (positive electrode): oxygen/O<sub>2</sub> <b>OR</b> hydrogen ion/proton/H<sup>+</sup> <b>AND</b> oxygen/O<sub>2</sub> ✓</p> <p>Cathode (negative electrode): hydrogen/H<sub>2</sub> <b>OR</b> hydroxide «ion»/OH<sup>-</sup> <b>AND</b> hydrogen/H<sub>2</sub> ✓</p>	Award [1 max] if correct products given at wrong electrodes.	2
4.	j		<p>Any two of: «inert» Pt electrode <b>OR</b> platinum black conductor ✓ 1 mol dm<sup>-3</sup> H<sup>+</sup>(aq) ✓ H<sub>2</sub>(g) at 100 kPa ✓</p>	<p>Accept 1 atm H<sub>2</sub>(g). Accept 1 bar H<sub>2</sub>(g) Accept a labelled diagram. Ignore temperature if it is specified.</p>	2 max
4.	k	i	Mg(s) + Cu <sup>2+</sup> (aq) → Mg <sup>2+</sup> (aq) + Cu(s) ✓		1
4.	k	ii	«+0.34 V - (-2.37 V) = +»2.71 «V» ✓		1
4.	k	iii	<p>cell potential increases ✓</p> <p>reaction «in Q4(k)(i)» moves to the right <b>OR</b> potential of the copper half-cell increases/becomes more positive ✓</p>	Accept correct answers based on the Nernst equation.	2


(continued)

Question			Answers	Notes	Total									
5.	a		<p><i>Propane:</i></p> <pre>       H   H   H                 H — C — C — C — H                       H   H   H           </pre> <p><b>AND</b></p> <p><i>Propene:</i></p> <pre>       H       H        \     /         C = C — C — H ✓        /     \           H       H   H           </pre>		1									
5.	b	i	<p><i>Sigma (<math>\sigma</math>):</i></p>  <p><i>Pi (<math>\pi</math>):</i></p> 		2									
5.	b	ii	<table border="1"> <thead> <tr> <th></th> <th>Number of sigma (<math>\sigma</math>) bonds</th> <th>Number of pi (<math>\pi</math>) bonds</th> </tr> </thead> <tbody> <tr> <td>Propane</td> <td>10</td> <td>0</td> </tr> <tr> <td>Propene</td> <td>8</td> <td>1</td> </tr> </tbody> </table> <p style="text-align: right;">✓✓</p>		Number of sigma ( $\sigma$ ) bonds	Number of pi ( $\pi$ ) bonds	Propane	10	0	Propene	8	1	<p>Award [1] for two or three correct answers. Award [2] for all four correct.</p>	2
	Number of sigma ( $\sigma$ ) bonds	Number of pi ( $\pi$ ) bonds												
Propane	10	0												
Propene	8	1												

(continued)



(Question 5 continued)

Question			Answers	Notes	Total
5.	c	i	$C_3H_8 + Br_2 \rightarrow C_3H_7Br + HBr$ ✓ «sun»light/UV/hν <b>OR</b> high temperature ✓	Do not accept "reflux" for M2.	2
5.	c	ii	$C_3H_6 + Br_2 \rightarrow C_3H_6Br_2$ ✓		1
5.	c	iii	Propane: «free radical» substitution / S <sub>R</sub> <b>AND</b> Propene: «electrophilic» addition / A <sub>E</sub> ✓		1
5.	d		 <p>curly arrow going from C=C to H of HBr <b>and</b> curly arrow showing Br leaving ✓                      representation of carbocation ✓                      curly arrow going from lone pair/negative charge on Br<sup>-</sup> to C<sup>+</sup> ✓</p>	Award [2 max] for formation of 1-bromopropane.	3

(continued)


Question		Answers	Notes	Total
6.	a	<p>correct isomer ✓ mirror image shown clearly ✓</p>		2
6.	b	<p>S<sub>N</sub>2 would give inversion of configuration «almost 100%» <b>OR</b> S<sub>N</sub>1 would give «approximately» 50% of each ✓ so mechanism is a mixture of both mechanisms ✓</p>		2
6.	c	<p>C–I bond «longer, so» weaker «than C–Br bond» <b>OR</b> I<sup>-</sup> is a better leaving group than Br<sup>-</sup> ✓</p>		1

(continued)

Question		Answers	Notes	Total
7.	a	<p><i>Calculation:</i>  <b>ALTERNATIVE 1:</b>  <math>[H^+] = (K_a \times [HA])^{1/2} / (1.6 \times 10^{-4} \times 0.0100)^{1/2} / 1.3 \times 10^{-3} \text{ «mol dm}^{-3}\text{»} \checkmark</math>  <math>pH = \text{«}-\log_{10}[H^+] \approx \text{»} 2.9 \checkmark</math>  <b>ALTERNATIVE 2:</b>  <math>pH = 0.5(pK_a - \log_{10}[HA]) \checkmark</math>  <math>pH = 2.9 \checkmark</math></p> <p><i>Assumption:</i>                      ionisation is <math>\ll 0.0100</math> so <math>0.0100 - [A^-] \approx 0.0100</math>  <b>OR</b>  <math>[HA]_{eqm} = [HA]_{initial}</math>  <b>OR</b>                      all <math>H^+</math> ions in the solution come from the acid «and not from the self-ionisation of water»  <b>OR</b>  <math>[H^+] = [HCOO^-] \checkmark</math></p>	<p><i>Award [2] for correct final answer.</i></p> <p><i>Do not accept partial dissociation.</i></p>	3

(continued)

(Question 7 continued)

Question		Answers	Notes	Total
b	i	 <p>correct shape of graph ✓ pH at half neutralization/equivalence ✓</p>	<p>M1: must show buffer region at pH &lt; 7 and equivalence at pH &gt; 7.</p> <p>Accept graph starting from where two axes meet as pH scale is not specified.</p>	2
b	ii	<p><b>ALTERNATIVE 1:</b>  <math>\text{HCOOH} \rightleftharpoons \text{HCOO}^- + \text{H}^+</math> ✓  <math>\text{H}^+</math> ions consumed in reaction with <math>\text{OH}^-</math> are produced again as equilibrium moves to the right «so <math>[\text{H}^+]</math> remains almost unchanged» ✓</p> <p><b>ALTERNATIVE 2:</b>  <math>\text{HCOOH} + \text{OH}^- \rightleftharpoons \text{HCOO}^- + \text{H}_2\text{O}</math> ✓                      added <math>\text{OH}^-</math> are neutralized by <math>\text{HCOOH}</math></p> <p><b>OR</b>                      strong base replaced by weak base ✓</p>	<p>Accept HA or any other weak acid in equations.                      Equilibrium sign must be included in equation for M1.</p>	2