

# Markscheme

May 2017

Mathematical studies

Standard level

Paper 1

23 pages

This markscheme is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Global Centre, Cardiff.

**Paper 1 Markscheme  
Instructions to Examiners**

**Notes:** If in doubt about these instructions or any other marking issues, contact your team leader for clarification.

The number of marks for each question is 6.

**1 Abbreviations**

The markscheme may make use of the following abbreviations:

- M** Marks awarded for **Method**
- A** Marks awarded for an **Answer** or for **Accuracy**
- C** Marks awarded for **Correct** answers (irrespective of working shown)
- R** Marks awarded for clear **Reasoning**
- ft** Marks that can be awarded as **follow through** from previous results in the question

**2 Method of Marking**

- (a) All marking must be done in RM Assessor using the mathematical studies annotations and in accordance with the current document for guidance in e-marking Mathematical Studies SL. It is essential that you read this document before you start marking.
- (b) If the candidate has full marks on a question use the **C6** annotation, if the candidate has made an attempt but scores zero marks use **C0**. If there is no attempt use the No response button. If a candidate does not score full or zero marks then full annotations **MUST** be shown.
- (c) In this paper, if the **correct answer is seen on the answer line the** maximum mark is awarded. **There is no need to check the working!** Award **C** marks and move on.
- (d) If the answer does not appear on the answer line, but the correct answer is seen in the working box with no subsequent working, award the maximum mark.
- (e) If the **answer is wrong**, marks should be awarded for the working according to the markscheme.
- (f) Working crossed out by the candidate should not be awarded any marks. Where candidates have written two solutions to a question, only the first solution should be marked.
- (g) A correct answer in the working box transcribed inaccurately to the answer line can receive full marks.
- (h) If correct working results in a correct answer **in the working box** but then further working is developed, indicating a lack of mathematical understanding full marks should **not** be awarded. In most such cases it will be a single final answer mark that is lost, however, a statement on the answer line should always be taken as the candidate's final decision on the answer **as long as it is unambiguous**. An exception to this may be in numerical answers, where a correct exact value is followed by an incorrect decimal.

Example:

	Correct answer seen	Further working seen	Action
1.	$8\sqrt{2}$	5.65685... (incorrect decimal value)	Award the final ( <b>A1</b> ) (ignore the further working)
2.	$(x-6)(x+1)$	$x = 6$ and $-1$	Do <b>not</b> award the final ( <b>A1</b> ) (see next example)

**Example:** Factorise  $x^2 - 5x - 6$

Markscheme	Candidates' Scripts	Marking
$(x-6)(x+1)$ <b>(A1)(A1)</b>	(i) Answer line: $(x+6)(x+1)$	<b>(A0)(A1)</b>
	(ii) Working box: $(x-6)(x+1)$	<b>(A1)</b>
	followed by $x=6$ and $-1$ , or just $6, -1$ in either working box or on answer line.	<b>(A0)</b>

### 3 Follow through (ft) Marks

Errors made at any step of a solution affect all working that follows. To limit the severity of the penalty, **follow through (ft)** marks can be awarded. Markschemes will indicate where it is appropriate to apply follow through in a question with '(ft)'.

- (a) Follow through applies only from one part of a question to a subsequent part of the question. Follow through does not apply within the same part.
- (b) If an answer resulting from follow through is extremely unrealistic (eg, negative distances or incorrect by large order of magnitude) then the final **A** mark should not be awarded.
- (c) If a question is transformed by an error into a **different, much simpler question** then follow through may not apply.
- (d) To award follow through marks for a question part, **there must be working present for that part**. An isolated follow through answer, without working is regarded as incorrect and receives no marks **even if it is approximately correct**.
- (e) The exception to the above would be in a question which is testing the candidate's use of the GDC, where working will not be expected. **The markscheme will clearly indicate where this applies**.
- (f) Inadvertent use of radians will be penalised the first time it occurs. The markscheme will give clear instructions to ensure that only one mark per paper can be lost for the use of radians.

**Example:** Finding angles and lengths using trigonometry

Markscheme	Candidates' Scripts	Marking
(a) $\frac{\sin A}{3} = \frac{\sin 30}{4}$ <b>(M1)(A1)</b>  $A = 22.0^\circ$ (22.0243...) <b>(A1)</b>	(a) $\frac{\sin A}{4} = \frac{\sin 30}{3}$	<b>(M1)(A0)</b>  <i>(use of sine rule but with wrong values)</i>
	$A = 41.8^\circ$ <i>(Note: the 2<sup>nd</sup> (A1) here was not marked (ft) and cannot be awarded because there was an earlier error in the same question part.)</i>	<b>(A0)</b>
(b) $x = 7 \tan (22.0243\dots^\circ)$ <b>(M1)</b> $= 2.83$ (2.83163...) <b>(A1)(ft)</b>	(b) case (i) $x = 7 \tan 41.8^\circ$ $= 6.26$	<b>(M1)</b> <b>(A1)(ft)</b>
	<b>but</b> case (ii) $6.26$	<b>(C0)</b> <i>since no working shown</i>

#### 4 Using the Markscheme

(a) **A** marks are **dependent** on the preceding **M** mark being awarded, it is **not** possible to award **(M0)(A1)**. Once an **(M0)** has been awarded, all subsequent **A** marks are lost in that part of the question, even if calculations are performed correctly, until the next **M** mark.  
The only exception will be for an answer where the accuracy is specified in the question – see section 5.

(b) **A** marks are **dependent** on the **R** mark being awarded, it is **not** possible to award **(A1)(R0)**. Hence the **(A1)** is not awarded for a correct answer if no reason or the wrong reason is given.

(c) **Alternative methods** may not always be included. Thus, if an answer is wrong then the working must be carefully analysed in order that marks are awarded for a different method consistent with the markscheme.  
Where alternative methods for complete questions are included in the markscheme, they are indicated by '**OR**' etc.

(d) Unless the question specifies otherwise, accept **equivalent forms**. For example:  $\frac{\sin \theta}{\cos \theta}$  for  $\tan \theta$ .

On the markscheme, these equivalent numerical or algebraic forms will sometimes be written in brackets after the required answer.

Where numerical answers are required as the final answer to a part of a question in the markscheme, the scheme will show, in order:

the 3 significant figure answer worked through from full calculator display;

the exact value  $\left( \text{for example } \frac{2}{3} \text{ if applicable} \right)$ ;

the full calculator display in the form 2.83163... as in the example above.

Where answers are given to 3 significant figures and are then used in subsequent parts of the question leading to a **different** 3 significant figure answer, these solutions will also be given.

(e) As this is an international examination, all valid **alternative forms of notation** should be accepted. Some examples of these are:

Decimal points: 1.7; 1'7; 1·7; 1,7 .

Decimal numbers less than 1 may be written with or without a leading zero: 0.49 or .49 .

Different descriptions of an interval:  $3 < x < 5$ ; (3, 5); ] 3, 5 [ .

Different forms of notation for set properties (e.g. complement):  $A'$ ;  $\bar{A}$ ;  $A^c$ ;  $U - A$ ;  $(A; U \setminus A$ .

Different forms of logic notation:  $\neg p$ ;  $p'$ ;  $\tilde{p}$ ;  $\bar{p}$ ;  $\sim p$ .

$p \Rightarrow q$ ;  $p \rightarrow q$ ;  $q \Leftarrow p$  .

Significance level may be written as  $\alpha$  .

(f) Discretionary marks: There will be very rare occasions where the markscheme does not cover the work seen. In such cases the annotation DM should be used to indicate where an examiner has used discretion. Discretion should be used sparingly and if there is doubt an exception should be raised through RM Assessor to the team leader.

As with previous sessions there will be no whole paper penalty marks for accuracy AP, financial accuracy FP and units UP. Instead these skills will be assessed in particular questions and the marks applied according to the rules given in sections 5, 6 and 7 below.

### 5 Accuracy of Answers

**Incorrect accuracy should be penalized once only in each question according to the rules below.**

Unless otherwise stated in the question, all numerical answers should be given exactly or correct to 3 significant figures.

1. If the **candidate's answer** is seen to 4 sf or greater **and** would round to the required 3 sf answer, then award **(A1)** and ignore subsequent rounding.

**Note:** The unrounded answer may appear in either the working box or on the final answer line.

2. If the candidate's unrounded answer is **not** seen then award **(A1)** if the answer given is **correctly** rounded to 2 or more significant figures, otherwise **(A0)**.

**Note:** *If the candidate's unrounded answer is not seen and the answer is given correct to 1 sf (correct or not), the answer will be considered wrong and will not count as incorrect accuracy.* If this answer is used in subsequent parts, then working must be shown for further marks to be awarded.

3. If a correct 2 sf answer is used in subsequent parts, then working **must** be shown for further marks to be awarded. (This treatment is the same as for following through from an incorrect answer.)

These 3 points (see numbers in superscript) have been summarized in the table below and illustrated in the examples which follow.

	If candidates final answer is given ...					
	Exact or to 4 or more sf (and would <b>round to the correct 3 sf</b> )	<b>Correct to 3 sf</b>	<b>Incorrect to 3 sf</b>	Correct to 2 sf <sup>3</sup>	Incorrect to 2 sf	Correct or incorrect to 1 sf
Unrounded answer seen <sup>1</sup>	Award the final <b>(A1)</b> irrespective of correct or incorrect rounding					
Unrounded answer not seen <sup>2</sup>	<b>(A1)</b>	<b>(A1)</b>	<b>(A0)</b>	<b>(A1)</b>	<b>(A0)</b>	<b>(A0)</b>
Treatment of subsequent parts	As per MS		Treat as follow through, only if working is seen. <sup>3</sup>			

Examples:

Markscheme	Candidates' Scripts	Marking
9.43 (9.43398...) <b>(A1)</b>	(i) 9.43398... is seen in the working box followed by 9; 9.4; 9.43; 9.434 etc. (correctly rounded)	<b>(A1)</b>
	(ii) 9.43398... is seen in the working box followed by 9.433; 9.44 etc. (incorrectly rounded)	<b>(A1)</b>
	(iii) 9.4	<b>(A1)</b>
	(iv) 9	<b>(A0)</b> <i>(correct to 1 sf)</i>
	(v) 9.3	<b>(A0)</b> <i>(incorrectly rounded to 2 sf)</i>
	(vi) 9.44	<b>(A0)</b> <i>(incorrectly rounded to 3 sf)</i>

Markscheme	Candidates' Scripts	Marking
7.44 (7.43798...) <b>(A1)</b>	(i) 7.43798... is seen in the working box followed by 7; 7.4; 7.44; 7.438 etc. (correctly rounded)	<b>(A1)</b>
	(ii) 7.43798... is seen in the working box followed by 7.437; 7.43 etc. (incorrectly rounded)	<b>(A1)</b>
	(iii) 7.4	<b>(A1)</b>
	(iv) 7	<b>(A0)</b> <i>(correct to 1 sf)</i>
	(v) 7.5	<b>(A0)</b> <i>(incorrectly rounded to 2 sf)</i>
	(vi) 7.43	<b>(A0)</b> <i>(incorrectly rounded to 3 sf)</i>

**Example:** ABC is a right angled triangle with angle  $ABC = 90^\circ$ ,  $AC = 32$  cm and  $AB = 30$  cm. Find (a) the length of BC, (b) The area of triangle ABC.

Markscheme	Candidates' Scripts	Marking
<p>(a) <math>BC = \sqrt{32^2 - 30^2}</math> <b>(M1)</b> Award <b>(M1)</b> for correct substitution in Pythagoras' formula</p> <p><math>= 11.1 (\sqrt{124}, 11.1355\dots)</math>(cm) <b>(A1)</b></p> <p>(b) <math>\text{Area} = \frac{1}{2} \times 30 \times 11.1355\dots</math> <b>(M1)</b> Award <b>(M1)</b> for correct substitution in area of triangle formula</p> <p><math>= 167(167.032\dots)</math>(cm<sup>2</sup>) <b>(A1)(ft)</b></p>	<p>(a) <math>BC = \sqrt{32^2 - 30^2}</math></p> <p>11 (cm) <b>(A1)</b> <i>(2 sf answer only seen, but correct)</i></p> <p>(b) case (i) <math>\text{Area} = \frac{1}{2} \times 30 \times 11</math> <b>(M1)</b> <i>(working shown)</i></p> <p><math>= 165</math> (cm<sup>2</sup>) <b>(A1)(ft)</b></p> <p>case (ii) <math>= 165</math> (cm<sup>2</sup>) <b>(M0)(A0)(ft)</b> <i>(No working shown, the answer 11 is treated as a ft, so no marks awarded here)</i></p>	

Rounding of an exact answer to 3 significant figures **should be accepted if performed correctly**.

Exact answers such as  $\frac{1}{4}$  can be written as decimals to fewer than 3 significant figures if the result is still exact. Reduction of a fraction to its lowest terms is **not** essential, however where an answer simplifies to an integer this is expected. Fractions that include a decimal in the numerator and/or the denominator are acceptable for showing correct substitution, but not as a final answer.

Ratios of  $\pi$  and answers taking the form of square roots of integers or any rational power of an integer (e.g.  $\sqrt{13}, 2^{\frac{2}{3}}, \sqrt[4]{5}$ ,) may be accepted as exact answers. All other powers (eg, of non-integers) and values of transcendental functions such as sine and cosine must be evaluated.

**If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy.** In **all** such cases the final mark is not awarded if the rounding does not follow the instructions given in the question. A mark for specified accuracy can be regarded as a **(ft)** mark regardless of an immediately preceding **(M0)**.



Certain answers obtained from the GDC are worth 2 marks and working will not be seen. In these cases only one mark should be lost for accuracy.

eg, Chi-squared, correlation coefficient, mean

Markscheme	Candidates' Scripts	Marking
Chi-squared	(a) 7.7	<b>(A2)</b>
7.68 (7.67543...) <b>(A2)</b>	(b) 7.67	<b>(A1)</b>
	(c) 7.6	<b>(A1)</b>
	(d) 8	<b>(A0)</b>
	(e) 7	<b>(A0)</b>
	(e) 7.66	<b>(A0)</b>

Regression line

Markscheme	Candidates' Scripts	Marking
$y = 0.888x + 13.5$ <b>(A2)</b> ( $y = 0.887686\dots x + 13.4895\dots$ )	(a) $y = 0.89x + 13$	<b>(A2)</b> <i>(both accepted)</i>
If an answer is not in the form of an equation award at most <b>(A1)(A0)</b> .	(b) $y = 0.88x + 13$	<b>(A1)</b> <i>(one rounding error)</i>
	(c) $y = 0.88x + 14$	<b>(A1)</b> <i>(rounding error repeated)</i>
	(d) (i) $y = 0.9x + 13$	<b>(A1)</b> <i>(1 sf not accepted)</i>
	(ii) $y = 0.8x + 13$	
	(e) $0.88x + 14$	<b>(A0)</b> <i>(two rounding errors and not an equation)</i>

Maximum/minimum/points of intersection

Markscheme	Candidates' Scripts	Marking
(2.06, 4.49) <b>(A1)(A1)</b> (2.06020..., 4.49253...)	(a) (2.1, 4.5)	<b>(A1)(A1)</b> <i>(both accepted)</i>
	(b) (2.0, 4.4)	<b>(A1)</b> <i>(same rounding error twice)</i>
	(c) (2.06, 4.4)	<b>(A1)</b> <i>(one rounding error)</i>
	(d) (2, 4.4)	<b>(A0)</b> <i>(1sf not accepted, one rounding error)</i>

### 6 Level of accuracy in finance questions

The accuracy level required for answers will be specified in all questions involving money. This will usually be either whole units or two decimal places. The first answer not given to the specified level of accuracy will not be awarded the final **A** mark. The markscheme will give clear instructions to ensure that only one mark per paper can be lost for incorrect accuracy in a financial question.

**Example:** A financial question demands accuracy correct to 2 dp.

Markscheme	Candidates' Scripts	Marking
\$231.62 (231.6189) <b>(A1)</b>	(i) 231.6	<b>(A0)</b>
	(ii) 232	<b>(A0)</b> <i>(Correct rounding to incorrect level)</i>
	(iii) 231.61	<b>(A0)</b>
	(iv) 232.00	<b>(A0)</b> <i>(Parts (iii) and (iv) are both incorrect rounding to correct level)</i>

### 7 Units in answers

There will be specific questions for which the units are required and this will be indicated clearly in the markscheme. The first correct answer with no units or incorrect units will not be awarded the final **A** mark. The markscheme will give clear instructions to ensure that only one or two marks per paper can be lost for lack of units or incorrect units.

The units are considered only when the numerical answer is awarded **(A1)** under the accuracy rules given in Section 5.

Markscheme	Candidates' Scripts	Marking
(a) 37 000 m <sup>2</sup> <b>(A1)</b>	(a) 36 000 m <sup>2</sup>	<b>(A0)</b> <i>(Incorrect answer so units not considered)</i>
(b) 3200 m <sup>3</sup> <b>(A1)</b>	(b) 3200 m <sup>2</sup>	<b>(A0)</b> <i>(Incorrect units)</i>

If no method is shown and the answer is correct but with incorrect or missing units award the C marks with a one mark penalty.

### 8 Graphic Display Calculators

Candidates will often obtain solutions directly from their calculators. They must use mathematical notation, not calculator notation. No method marks can be awarded for incorrect answers supported only by calculator notation. The comment "I used my GDC" cannot receive a method mark.

1. (a)  $\sqrt[3]{\frac{2.78 \times 10^{11}}{3.12 \times 10^{-3}}}$  OR  $\sqrt[3]{8.91025... \times 10^{13}}$  (M1)

**Note:** Award (M1) for correct substitution into given expression.

44664.59503 (A1) (C2)

**Note:** Award (A1) for a correct answer with at least 8 digits.  
Accept 44664.5950301.

[2 marks]

(b) (i) 44664.60 (A1)(ft) (C1)

**Note:** For a follow through mark, the answer to part (a) must be to at least 3 decimal places.

(ii) 44700 (A1)(ft) (C1)

**Notes:** Answer to part (a) must be to at least 4 significant figures.  
Accept any equivalent notation which is correct to 3 significant figures.  
For example  $447 \times 10^2$  or  $44.7 \times 10^3$ .  
Follow through from part (a).

[2 marks]

(c)  $4.47 \times 10^4$  (A1)(ft)(A1)(ft) (C2)


**Notes:** Award (A1)(ft) for 4.47 and (A1)(ft) for  $10^4$ .  
Award (A0)(A0) for answers such as  $44.7 \times 10^3$ .  
Follow through from part (b)(ii) **only**.

[2 marks]

**Total [6 marks]**

2. (a)  $\frac{1}{2}x$  (A1) (C1)  
 [1 mark]

(b)



(A1)(A1)(ft) (C2)

**Notes:** Award (A1) for 15 placed in the correct position, award (A1)(ft) for  $x$  and their  $\frac{1}{2}x$  placed in the correct positions of diagram. Do not penalize the absence of 0 inside the rectangle and award at most (A1)(A0) if any value other than 0 is seen outside the circles. Award at most (A1)(A0) if 35 and 70 are seen instead of  $x$  and their  $\frac{1}{2}x$ .

[2 marks]

- (c)  $x + \frac{1}{2}x + 15 = 120$  or equivalent (M1)

**Note:** Award (M1) for adding the values in their Venn and equating to 120 (or equivalent).

(x =) 70 (A1)(ft) (C2)

**Note:** Follow through from their Venn diagram, but only if the answer is a positive integer and  $x$  is seen in their Venn diagram.

[2 marks]

- (d) 85 (A1)(ft) (C1)

**Note:** Follow through from their Venn diagram and their answer to part (c), but only if the answer is a positive integer and less than 120.

[1 mark]

Total [6 marks]

3. (a)  $\cos 60^\circ = \frac{20}{b}$  **OR**  $b = \frac{20}{\cos 60^\circ}$  **(M1)**

**Note:** Award **(M1)** for correct substitution into a correct trig. ratio.

$(b =) 40$  (cm) **(A1)** **(C2)**  
**[2 marks]**

(b)  $4 \times 40 + 2\pi(20)$  **(M1)(M1)**

**Note:** Award **(M1)** for correct substitution in the circumference of the circle formula, **(M1)** for adding 4 times their answer to part (a) to their circumference of the circle.

285.6637... **(A1)(ft)**

**Note:** Follow through from part (a). This **(A1)** may be implied by a correct rounded answer.

285.7 (cm) **(A1)(ft)** **(C4)**

**Notes:** Award **(A1)(ft)** for rounding their answer (consistent with their method) to the nearest millimetre, irrespective of unrounded answer seen.  
The final **(A1)(ft)** is not dependent on any of the previous **M** marks. It is for rounding their unrounded answer correctly.

**[4 marks]**

**Total [6 marks]**

4. (a)  $-3$  (A1)(A1) (C2)

**Notes:** Award (A1) for 3 and (A1) for a negative value.  
Award (A1)(A0) for either  $3x$  or  $-3x$ .

[2 marks]

(b)  $6 = -3(2) + c$  OR  $(y - 6) = -3(x - 2)$  (M1)

**Note:** Award (M1) for substitution of their gradient from part (a) into a correct equation with the coordinates (2, 6) correctly substituted.

$y = -3x + 12$  (A1)(ft) (C2)

**Notes:** Award (A1)(ft) for their correct equation. Follow through from part (a).  
If no method seen, award (A1)(A0) for  $y = -3x$ .  
Award (A1)(A0) for  $-3x + 12$ .

[2 marks]

(c)  $0 = -3x + 12$  (M1)

**Note:** Award (M1) for substitution of  $y = 0$  in their equation from part (b).

$(x =) 4$  (A1)(ft) (C2)

**Notes:** Follow through from their equation from part (b). Do not follow through if no method seen. Do not award the final (A1) if the value of  $x$  is negative or zero.

[2 marks]

**Total [6 marks]**

5. (a)  $4 + 3(n - 1) = 52$  **(M1)(A1)**

**Note:** Award **(M1)** for substitution into the formula of the  $n$ th term of an arithmetic sequence, **(A1)** for correct substitution.

$n = 17$  **(A1)** **(C3)**  
[3 marks]

(b)  $\frac{24}{2}(2 \times 4 + 23 \times 3)$  **OR**  $\frac{24}{2}(4 + 73)$  **(M1)(A1)(ft)**

**Notes:** Award **(M1)** for substitution into the sum of the first  $n$  terms of an arithmetic sequence formula, **(A1)(ft)** for their correct substitution, consistent with part (a).

924 **(A1)(ft)** **(C3)**

**Note:** Follow through from part (a).

[3 marks]

**Total [6 marks]**

6. (a) 9 (cm) **(A1)** **(C1)**  
[1 mark]

(b) 40 (leaves) **(A1)** **(C1)**  
[1 mark]

(c) (i)  $(200 \times 0.90 =) 180$  or equivalent **(M1)**

**Note:** Award **(M1)** for a horizontal line drawn through the cumulative frequency value of 180 and meeting the curve (or the corresponding vertical line from 10.5 cm).

$(k =) 10.5$  (cm) **(A1)** **(C2)**

**Note:** Accept an error of  $\pm 0.1$ .

(ii)  $\left| \frac{9.5 - 10.5}{10.5} \right| \times 100\%$  **(M1)**

**Notes:** Award **(M1)** for their correct substitution into the percentage error formula.

9.52 (%) (9.52380... (%)) **(A1)(ft)** **(C2)**

**Notes:** Follow through from their answer to part (c)(i).  
Award **(A1)(A0)** for an answer of  $-9.52$  with or without working.

[4 marks]

**Total [6 marks]**

7. (a)  $18 + x + y + 22 = 100$  or equivalent (A1) (C1)  
[1 mark]

(b)  $\frac{18 + 2x + 3y + 88}{100} = 2.71$  or equivalent (M1)(A1) (C2)

**Note:** Award (M1) for a sum including  $x$  and  $y$ , divided by 100 and equated to 2.71, (A1) for a correct equation.

[2 marks]

(c)  $x + y = 60$  and  $2x + 3y = 165$  (M1)

**Note:** Award (M1) for obtaining a correct linear equation in one variable from their (a) and their (b). This may be implied if seen in part (a) or part (b).

$x = 15; y = 45$  (A1)(ft)(A1)(ft) (C3)

**Notes:** Follow through from parts (a) and (b), irrespective of working seen provided the answers are positive integers.

[3 marks]

**Total [6 marks]**



8. (a)  $8000 \times 0.09819 \times 0.98$  (M1)(M1)

**Note:** Award (M1) for multiplying 8000 by 0.09819, (M1) for multiplying by 0.98 (or equivalent).

769.81 (EUR) (A1) (C3)  
[3 marks]

- (b)  $r\% \times \frac{85}{0.08753} = 14.57$  (M1)(M1)

**Note:** Award (M1) for dividing 85 by 0.08753, and (M1) for multiplying their  $\frac{85}{0.08753}$  by  $r\%$  and equating to 14.57.

OR

$$\frac{85}{0.08753} = 971.095\dots \quad (M1)$$

**Note:** Award (M1) for dividing 85 by 0.08753.

$$\frac{14.57}{9.71095\dots} \quad \text{OR} \quad \frac{14.57}{971.095\dots} \times 100 \quad (M1)$$

**Note:** Award (M1) for dividing 14.57 by 9.71095... or equivalent.

$r = 1.50$  (1.50036...) (A1) (C3)  
[3 marks]

Total [6 marks]

9. (a)  $\frac{1}{2}$  (0.5) (A1) (C1)  
[1 mark]

(b)  $18 \times \left(\frac{1}{2}\right)^4$  (M1)

**Note:** Award **(M1)** for their correct substitution into the geometric sequence formula. Accept a list of their five correct terms.

$1.125 \left(1.13, \frac{9}{8}\right)$  (A1)(ft) (C2)

**Note:** Follow through from their common ratio from part (a).

[2 marks]

(c)  $18 \times \left(\frac{1}{2}\right)^{n-1} < 10^{-3}$  (M1)(M1)

**Notes:** Award **(M1)** for their correct substitution into the geometric sequence formula with a variable in the exponent, **(M1)** for comparing their expression with  $10^{-3} \left(\frac{1}{1000}\right)$ . Accept an equation.

$n = 16$  (A1)(ft) (C3)

**Note:** Follow through from their common ratio from part (a). “n” must be a positive integer for the **(A1)** to be awarded.

[3 marks]

**Total [6 marks]**

10. (a)  $0.93$  (93%) (A1) (C1)  
[1 mark]

(b) (i)  $0.93 \times 0.93$  (M1)

**Note:** Award (M1) for squaring their answer to part (a).

$0.865$  (0.8649; 86.5%) (A1)(ft) (C2)

**Notes:** Follow through from part (a).  
Accept  $0.86$  (unless it follows  $\frac{93}{100} \times \frac{92}{99}$ ).

(ii)  $1 - 0.8649$  (M1)

**Note:** Follow through from their answer to part (b)(i).

**OR**

$0.07 \times 0.07 + 2 \times (0.07 \times 0.93)$  (M1)

**Note:** Follow through from part (a).

$0.135$  (0.1351; 13.5%) (A1)(ft) (C2)  
[4 marks]

(c)  $1 - a^3$  (A1) (C1)

**Note:** Accept  $3a^2(1-a) + 3a(1-a)^2 + (1-a)^3$  or equivalent.

[1 mark]

**Total [6 marks]**

11. (a)  $0.5 \left( 50\%, \frac{1}{2} \right)$  (A1) (C1)

[1 mark]

(b)  $P(X > a) = 0.25$  OR  $P(X < a) = 0.75$  (M1)

**Note:** Award (M1) for a sketch of approximate normal curve with a vertical line drawn to the right of the mean with the area to the right of this line shaded.

$a = 434$  (g) (433.724... (g)) (A1) (C2)  
[2 marks]

(c)  $33.7244... \times 2$  (A1)(ft)(M1)


**Note:** Award (A1)(ft) for 33.7244... (or 433.7244... - 400) seen, award (M1) for multiplying their 33.7244... by 2. Follow through from their answer to part (b).

OR

$434 - 366.275...$  (A1)(ft)(M1)

**Note:** Award (A1)(ft) for their 366.275... (366) seen, (M1) for difference between their answer to (b) and their 366.

OR



(A1)(ft)(M1)

**Note:** Award (A1)(ft) for their 366.275... (366) seen. Award (M1) for correct symmetrical region indicated on labelled normal curve.

67.4 (g) (A1)(ft) (C3)

**Note:** Accept an answer of 68 from use of rounded values. Follow through from part (b).

[3 marks]

Total [6 marks]

12. (a)  $\pi \times 8^2 \times 12$  (M1)

**Note:** Award (M1) for correct substitution into the volume of a cylinder formula.

$2410 \text{ cm}^3$  ( $2412.74 \dots \text{ cm}^3$ ,  $768\pi \text{ cm}^3$ ) (A1) (C2)  
[2 marks]

(b)  $\frac{4}{3}\pi \times 2.9^3 + 768\pi = \pi \times 8^2 h$  (M1)(M1)(M1)

**Note:** Award (M1) for correct substitution into the volume of a sphere formula (this may be implied by seeing 102.160...), (M1) for adding their volume of the ball to their part (a), (M1) for equating a volume to the volume of a cylinder with a height of  $h$ .

OR

$\frac{4}{3}\pi \times 2.9^3 = \pi \times 8^2 (h - 12)$  (M1)(M1)(M1)

**Note:** Award (M1) for correct substitution into the volume of a sphere formula (this may be implied by seeing 102.160...), (M1) for equating to the volume of a cylinder, (M1) for the height of the water level increase,  $h - 12$ . Accept  $h$  for  $h - 12$  if adding 12 is implied by their answer.

$(h =) 12.5 \text{ (cm)}$  ( $12.5081 \dots \text{ (cm)}$ ) (A1)(ft) (C4)

**Note:** If 3 sf answer used, answer is 12.5 (12.4944...). Follow through from part (a) if first method is used.

[4 marks]

Total [6 marks]

13. (a) 3

(A1) (C1)

Notes: Accept  $y = 3$

[1 mark]

(b)  $3 = 0.5(1) + c$  OR  $y - 3 = 0.5(x - 1)$

(A1)(A1)

Note: Award (A1) for correct gradient, (A1) for correct substitution of A(1, 3) in the equation of line.

$x - 2y + 5 = 0$  or any integer multiple


(A1)(ft)

(C3)

Note: Award (A1)(ft) for their equation correctly rearranged in the indicated form. The candidate's answer **must** be an equation for this mark.

[3 marks]

(c)



(M1)(A1)(ft)

(C2)

Note: Award M1) for a straight line, with positive gradient, passing through (1, 3), (A1)(ft) for line (or extension of their line) passing approximately through 2.5 or their intercept with the y-axis..

[2 marks]

Total [6 marks]

14. (a) 400 (USD) (A1) (C1)  
[1 mark]

(b)  $8500(0.95)^t = 400 \times t + 2000$  (M1)

**Note:** Award (M1) for equating  $8500(0.95)^t$  to  $400 \times t + 2000$  or for comparing the difference between the two expressions to zero or for showing a sketch of both functions.

$(t =) 8.64$  (months)  $(8.6414\dots(\text{months}))$  (A1) (C2)

**Note:** Accept 9 months.

[2 marks]

(c)  $8500(0.95)^2 - (400 \times 2 + 2000)$  (M1)(M1)

**Note:** Award (M1) for correct substitution of  $t = 2$  into equation for  $P$ , (M1) for finding the difference between a value/expression for  $P$  and a value/expression for  $S$ . The first (M1) is implied if 7671.25 seen.

4870 (USD)  $(4871.25)$  (A1) (C3)

**Note:** Accept 4871.3.

[3 marks]

**Total [6 marks]**

15.

Conditions	Graph Number
$a > 0, b < 0, c > 0$	2
$a < 0, b = 0, c > 0$	6
$a < 0, b > 0, c < 0$	3
$a > 0, b = 0, c = 0$	5
$a > 0, b > 0, c < 0$	4
$a < 0, b < 0, c = 0$	1

(A1)(A1)(A1)(A1)(A1)(A1) (C6)

**Note:** Award (A1) for each correct entry.

[6 marks]