

Mathematics
Higher level
Paper 3 – calculus

Thursday 16 November 2017 (afternoon)

1 hour

Instructions to candidates

- Do not open this examination paper until instructed to do so.
- Answer all the questions.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A graphic display calculator is required for this paper.
- A clean copy of the **mathematics HL and further mathematics HL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[50 marks]**.

Please start each question on a new page. Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working. For example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

1. [Maximum mark: 5]

The function f is defined by

$$f(x) = \begin{cases} x^2 - 2, & x < 1 \\ ax + b, & x \geq 1 \end{cases}$$

where a and b are real constants.

Given that both f and its derivative are continuous at $x = 1$, find the value of a and the value of b .

2. [Maximum mark: 10]

Consider the differential equation $\frac{dy}{dx} + \frac{x}{x^2 + 1}y = x$ where $y = 1$ when $x = 0$.

(a) Show that $\sqrt{x^2 + 1}$ is an integrating factor for this differential equation. [4]

(b) Solve the differential equation giving your answer in the form $y = f(x)$. [6]

3. [Maximum mark: 12]

(a) Use the limit comparison test to show that the series $\sum_{n=1}^{\infty} \frac{1}{n^2 + 2}$ is convergent. [3]

Let $S = \sum_{n=1}^{\infty} \frac{(x-3)^n}{n^2 + 2}$.

(b) Find the interval of convergence for S . [9]

4. [Maximum mark: 10]

The mean value theorem states that if f is a continuous function on $[a, b]$ and differentiable on $]a, b[$ then $f'(c) = \frac{f(b) - f(a)}{b - a}$ for some $c \in]a, b[$.

The function g , defined by $g(x) = x \cos(\sqrt{x})$, satisfies the conditions of the mean value theorem on the interval $[0, 5\pi]$.

- (a) For $a = 0$ and $b = 5\pi$, use the mean value theorem to find all possible values of c for the function g . [6]
- (b) Sketch the graph of $y = g(x)$ on the interval $[0, 5\pi]$ and hence illustrate the mean value theorem for the function g . [4]

5. [Maximum mark: 13]

Consider the function $f(x) = \sin(p \arcsin x)$, $-1 < x < 1$ and $p \in \mathbb{R}$.

- (a) Show that $f'(0) = p$. [2]

The function f and its derivatives satisfy

$$(1 - x^2)f^{(n+2)}(x) - (2n + 1)xf^{(n+1)}(x) + (p^2 - n^2)f^{(n)}(x) = 0, n \in \mathbb{N}$$

where $f^{(n)}(x)$ denotes the n th derivative of $f(x)$ and $f^{(0)}(x)$ is $f(x)$.

- (b) Show that $f^{(n+2)}(0) = (n^2 - p^2)f^{(n)}(0)$. [1]
- (c) For $p \in \mathbb{R} \setminus \{\pm 1, \pm 3\}$, show that the Maclaurin series for $f(x)$, up to and including the x^5 term, is

$$px + \frac{p(1 - p^2)}{3!}x^3 + \frac{p(9 - p^2)(1 - p^2)}{5!}x^5. [4]$$

- (d) Hence or otherwise, find $\lim_{x \rightarrow 0} \frac{\sin(p \arcsin x)}{x}$. [2]
- (e) If p is an odd integer, prove that the Maclaurin series for $f(x)$ is a polynomial of degree p . [4]