

# Markscheme

May 2018

Chemistry


Standard level


Paper 2

13 pages



This markscheme is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Global Centre, Cardiff.

Question		Answers	Notes	Total
1.	a	$n(\text{H}_2\text{SO}_4) \llcorner = 0.0500 \text{ dm}^3 \times 0.100 \text{ mol dm}^{-3} \llcorner = 0.00500/5.00 \times 10^{-3} \llcorner \llcorner \text{mol} \llcorner \llcorner \checkmark$		1
1.	b	$\text{H}_2\text{SO}_4(\text{aq}) + \text{Mg}(\text{OH})_2(\text{s}) \rightarrow \text{MgSO}_4(\text{aq}) + 2\text{H}_2\text{O}(\text{l}) \checkmark$	<i>Accept an ionic equation.</i>	1
1.	c	$\llcorner n(\text{H}_2\text{SO}_4) = \frac{1}{2} \times n(\text{NaOH}) = \frac{1}{2} (0.02080 \text{ dm}^3 \times 0.1133 \text{ mol dm}^{-3}) \llcorner$ $0.001178/1.178 \times 10^{-3} \llcorner \llcorner \text{mol} \llcorner \llcorner \checkmark$		1
1.	d	$n(\text{H}_2\text{SO}_4) \text{ reacted } \llcorner = 0.00500 - 0.001178 \llcorner = 0.00382/3.82 \times 10^{-3} \llcorner \llcorner \text{mol} \llcorner \llcorner \checkmark$		1
1.	e	$n(\text{Mg}(\text{OH})_2) \llcorner = n(\text{H}_2\text{SO}_4) = \llcorner = 0.00382/3.82 \times 10^{-3} \llcorner \llcorner \text{mol} \llcorner \llcorner \checkmark$ $m(\text{Mg}(\text{OH})_2) \llcorner = 0.00382 \text{ mol} \times 58.33 \text{ g mol}^{-1} \llcorner = 0.223 \llcorner \llcorner \text{g} \llcorner \llcorner \checkmark$	<i>Award [2] for correct final answer.</i>	2
1.	f	$\% \text{ Mg}(\text{OH})_2 \llcorner = \frac{0.223 \text{ g}}{1.24 \text{ g}} \times 100 \llcorner = 18.0 \llcorner \llcorner \% \llcorner \llcorner \checkmark$	<i>Answer must show three significant figures.</i>	1

Question		Answers	Notes	Total
2.	a	 <p>both axes correctly labelled ✓                      correct shape of curve starting at origin ✓  <math>E_{a(catalyst)} &lt; E_{a(without catalyst)}</math> on x-axis ✓</p>	<p><b>M1:</b>                      Accept "speed" for x-axis label.                      Accept "number of particles", "N", "frequency" or "probability «density»" for y-axis label.                      Do <b>not</b> accept "potential energy" for x-axis label.</p> <p><b>M2:</b>                      Do <b>not</b> accept a curve that touches the x-axis at high energy.                      Do <b>not</b> award M2 if two curves are drawn.</p> <p><b>M3:</b>                      Ignore any shading under the curve.</p>	<b>3</b>

Question			Answers	Notes	Total
2.	b	i	 <p>curve starting from origin with steeper gradient <b>AND</b> reaching same maximum volume ✓</p>		1
2.	b	ii	<p>rate decreases <b>OR</b> slower reaction ✓</p> <p>«ethanoic acid» partially dissociated/ionized «in solution/water» <b>OR</b> lower <math>[H^+]</math> ✓</p>	Accept "weak acid" or "higher pH".	2

Question		Answers	Notes	Total
2.	c	<p>«pH» converts «wide range of <math>[H^+]</math>» into simple «log» scale/numbers  <b>OR</b>            «pH» avoids need for exponential/scientific notation  <b>OR</b>            «pH» converts small numbers into values «typically» between 0/1 and 14  <b>OR</b>            «pH» allows easy comparison of values of <math>[H^+]</math> ✓</p>	<p>Accept “uses values between 0/1 and 14”.</p> <p>Do <b>not</b> accept “easier to use”.</p> <p>Do <b>not</b> accept “easier for calculations”.</p>	1
2.	d	<p>«species» do not differ by a «single» proton/<math>H^+</math>  <b>OR</b>            conjugate base of <math>H_3PO_4</math> is <math>H_2PO_4^-</math> «not <math>HPO_4^{2-}</math>»  <b>OR</b>            conjugate acid of <math>HPO_4^{2-}</math> is <math>H_2PO_4^-</math> «not <math>H_3PO_4</math>» ✓</p>	<p>Do <b>not</b> accept “hydrogen/H” for “<math>H^+</math>/proton”.</p>	1

Question			Answers	Notes	Total
3.	a	i	 <p>4 levels showing convergence at higher energy ✓</p>		1
3.	a	ii	 <p>arrows (pointing down) from <math>n = 3</math> to <math>n = 2</math> <b>AND</b> <math>n = 4</math> to <math>n = 2</math> ✓</p>		1

Question			Answers	Notes	Total						
3.	b	i	same number of shells/«outer» energy level/shielding <b>AND</b> nuclear charge/number of protons/ $Z_{\text{eff}}$ increases «causing a stronger pull on the outer electrons» ✓		1						
3.	b	ii	$\text{K}^+$ 19 protons <b>AND</b> $\text{Cl}^-$ 17 protons <b>OR</b> $\text{K}^+$ has «two» more protons ✓ same number of electrons/isoelectronic «thus pulled closer together» ✓		2						
3.	c	i	<table border="1" style="display: inline-table; vertical-align: middle;"><tr><td>1</td></tr></table> <table border="1" style="display: inline-table; vertical-align: middle;"><tr><td>↑↓</td><td>↑↓</td><td>↑↓</td><td>↑↓</td><td>↑↓</td></tr></table>	1	↑↓	↑↓	↑↓	↑↓	↑↓		1
1											
↑↓	↑↓	↑↓	↑↓	↑↓							
3.	c	ii	<i>Anode (positive electrode):</i> $\text{Cu(s)} \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{e}^-$ ✓  <i>Cathode (negative electrode):</i> $\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu(s)}$ ✓	<i>Accept <math>\text{Cu(s)} - 2\text{e}^- \rightarrow \text{Cu}^{2+}(\text{aq})</math>.</i> <i>Accept <math>\rightleftharpoons</math> for <math>\rightarrow</math>.</i> <i>Award [1 max] if the equations are at the wrong electrodes.</i>	2						
3.	c	iii	«external» circuit/wire <b>AND</b> from positive/anode to negative/cathode electrode ✓	<i>Accept “through power supply/battery” instead of “circuit”.</i>	1						









Question			Answers	Notes	Total
4.	a		<p>bonds broken: <math>4(\text{C-H}) + 2(\text{H-O}) / 4(414) + 2(463) / 2582</math> «kJ» ✓  bonds made: <math>3(\text{H-H}) + \text{C}\equiv\text{O} / 3(436) + 1077 / 2385</math> «kJ» ✓</p> $\Delta H \llcorner = \sum \text{BE}_{(\text{bonds broken})} - \sum \text{BE}_{(\text{bonds made})} = 2582 - 2385 \llcorner = \llcorner + \llcorner 197 \llcorner \llcorner \text{kJ} \llcorner \llcorner \llcorner$	<p>Award <b>[3]</b> for correct final answer.  Award <b>[2 max]</b> for <math>-197</math> «kJ».</p>	3
4.	b	i	$\Delta H_f^\ominus$ for any element = 0 «by definition» <b>OR</b> no energy required to form an element «in its stable form» from itself ✓		1
4.	b	ii	$\Delta H^\ominus \llcorner = \sum \Delta H_f^\ominus (\text{products}) - \sum \Delta H_f^\ominus (\text{reactants}) = -111 + 0 - [-74.0 + (-242)] \llcorner$ $= \llcorner + \llcorner 205 \llcorner \llcorner \text{kJ} \llcorner \llcorner \llcorner$		1
4.	b	iii	«bond enthalpies» averaged values «over similar compounds» <b>OR</b> «bond enthalpies» are not specific to these compounds ✓		1

5.	a		<p>Q: non-equilibrium concentrations <b>AND</b> <math>K_c</math>: equilibrium concentrations  <b>OR</b>  Q: «measured» at any time <b>AND</b> <math>K_c</math>: «measured» at equilibrium ✓</p>		1
5.	b		$\llcorner Q = \frac{[\text{SO}_3]^2}{[\text{SO}_2]^2 [\text{O}_2]} = \frac{1.00^2}{1.00^2 \times 2.00} \llcorner = 0.500 \llcorner$ <p>reverse reaction favoured/reaction proceeds to the left <b>AND</b>  <math>Q &gt; K_c / 0.500 &gt; 0.282</math> ✓</p>	Do <b>not</b> award M2 without M1.	2

Question			Answers	Notes	Total
6.	a	i	polar bonds «between H and group 16 element» <b>OR</b> difference in electronegativities «between H and group 16 element» ✓  uneven distribution of charge/electron cloud <b>OR</b> non-linear/bent/V-shaped/angular shape «due to lone pairs» <b>OR</b> polar bonds/dipoles do not cancel out ✓	M2: Do <b>not</b> accept “net/overall dipole moment” without further explanation. Accept “non-symmetrical «shape/distribution of charge»”.	2
6.	a	ii	number of electrons increases ✓  London/dispersion/instantaneous induced dipole-induced dipole forces increase ✓	M1: Accept “ $M_r/A_r$ increases” or “molecules become larger in size/mass/surface area”.	2
6.	b		Electron domain geometry: tetrahedral ✓  Molecular geometry: bent/V-shaped/angular ✓	Both marks can be awarded for clear diagrams. Electron domain geometry requires a 3-D diagram showing the tetrahedral arrangement.	2

Question		Answers	Notes	Total
7.	a	<p><i>Physical evidence:</i>                      equal C–C bond «lengths/strengths»  <b>OR</b>                      regular hexagon  <b>OR</b>                      «all» C–C have bond order of 1.5  <b>OR</b>                      «all» C–C intermediate between single and double bonds ✓</p> <p><i>Chemical evidence:</i>                      undergoes substitution reaction «more readily than addition»  <b>OR</b>                      does not discolour/react with bromine water  <b>OR</b>                      substitution forms only one isomer for 1,2-disubstitution «presence of alternate double bonds would form two isomers»  <b>OR</b>                      more stable than expected «compared to hypothetical molecule cyclohexa-1,3,5-triene»  <b>OR</b>                      enthalpy change of hydrogenation/combustion is less exothermic than predicted «for cyclohexa-1,3,5-triene» ✓</p>	<p><i>M1:</i>                      Accept “all C–C–C bond angles are equal”.</p>	2

Question			Answers	Notes	Total
7.	b	i	$3\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}(\text{l}) + \text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 8\text{H}^+(\text{aq}) \rightarrow 3\text{CH}_3\text{CH}_2\text{CHO}(\text{aq}) + 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l})$ correct reactants and products ✓ balanced equation ✓		2
7.	b	ii	Aldehyde: by distillation «removed from reaction mixture as soon as formed» ✓  Carboxylic acid: «heat mixture under» reflux «to achieve complete oxidation to –COOH» ✓	Accept clear diagrams or descriptions of the processes.	2
7.	c	i	$\left\langle \frac{136}{48 + 4 + 16} = 2 \right\rangle$ C <sub>8</sub> H <sub>8</sub> O <sub>2</sub> ✓		1
7.	c	ii	A: C–H «in alkanes, alkenes, arenes» <b>AND</b> B: C=O «in aldehydes, ketones, carboxylic acids and esters» ✓		1

Question			Answers	Notes	Total
7.	c	iii	<p>Any two of:</p>  <p><b>OR</b> <math>C_6H_5COOCH_3</math> ✓</p>  <p><b>OR</b> <math>CH_3COOC_6H_5</math> ✓</p>  <p><b>OR</b> <math>HCOOCH_2C_6H_5</math> ✓</p>	<p>Do <b>not</b> penalize use of Kekule structures for the phenyl group.</p> <p>Accept the following structures:</p>    <p>Award <b>[1 max]</b> for two correct aliphatic/linear esters with the molecular formula <math>C_8H_8O_2</math>.</p>	2
7.	c	iv	<p><math>C_6H_5COOCH_3</math> «signal at 4 ppm (3.7 – 4.8 range in data table) due to alkyl group on ester» ✓</p>		1