

Markscheme

November 2018

Sports, exercise and health science

Higher level

Paper 3

This markscheme is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Global Centre, Cardiff.

Option A — Optimizing physiological performance

Q	uestior	Answers	Notes	Total
1.	а	3200 « g » √		1
1.	b	3200–2800 ✓ = 400 «g» ✓	Accept the subtraction in a different order.	2
1.	С	CWI did not affect muscle mass ACT helped in the development of muscle mass ✓	Accept in the converse.	2
		CWI is «significantly» less effective than ACT in developing muscle mass ✓		

-3-

2.	а	overreaching is transient overtraining ✓ increasing frequency/intensity/duration of an exercise for improvement ✓		1
2.	b	overtraining is when an athlete attempts to do more training than he or she is able to physically and/or mentally tolerate ✓		1
2.	С	decreased appetite. Noticeable behavioural change in food intake leading to body weight loss/fat and muscle loss ✓	Award [1 max] for listing three indicators.	
		chronic soreness such as muscle or bone tenderness/soreness «which is a sign the muscles are not recovering» ✓		
		fatigue indicators including sleep disturbance «combination of nervous system and or hormonal system overload» / nausea ✓		3 max
		elevated resting HR / BP ✓		
		unexplained decline in performance ✓		

_	4	-
---	---	---

	increased susceptibility to infections/reduced immune function / continual catabolic	
	state ✓	

C	Questio	n Answers	Notes	Total
2.	d	periodization / mesocycle would be used for training for the world championships ✓		
		there may be multiple preparation phases within a macrocycle ✓		
		the preparation phase can be broken into mesocycles with a specific focus (1–3 months) \checkmark		
		used for optimizing performance where training will start broad and become more specific <i>eg</i> fitness ✓		4 max
		multiple mesocycles can be broken down into microcycles (7–14 days) ✓		
		elicit specific adaptations for the sprinter by balancing high intensity work with rest and recovery \checkmark		

C	uesti	on	Answers	Notes	Total	
3.	а	i	low-intensity exercise to promote recovery either immediately after / or in the days following, an intense training session or competition ✓	OWTTE	1	
3.	а	ii	raised circulation rate ✓ enhanced blood lactate removal ✓ accelerated raising of blood pH ✓		2 max	
3.	b	i	shivering ✓ non-shivering thermogenesis ✓ «peripheral» vasoconstriction ✓		1 max	
3.	b	ii	Strengths: acts as an analgesic and anti-inflammatory for soft tissue ✓ some methods are easily affordable ✓ perception of enhanced recovery rates lead to improved performance ✓ Limitations: current recommendations are largely based on anecdotal rather than scientific research ✓ can be costly ✓ Risks associated with exposure to prolonged or extreme cold ✓	Award [2 max] for strengths. Award [2 max] for limitations.	3 max	

C	uestion		Answers			Notes	Total
4.	a	Hemoglobin saturation Respiratory rates at exercise Submaximal heart rate Effectiveness of glycolysis Lactic acid production Maximal oxygen consumption (VO ₂ max) Body mass Cardiac output	Sea level High (98.5 %) Lower Normal Normal Normal Normal Normal Normal	Altitude Lower (~60 %) Higher Elevated Decreased Elevated Reduced Decrease Increase	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \		2 max
4.	b	less oxygen at low oxygen less smaller reduction in hemoglorestore acid-base balance Muscle adaptations: reduced lean body mass increased capillary density, princreased mitochondrial denerythrpoietic response increased	Blood adaptations: increased number of red blood cells / viscosity to compensate for each carrying less oxygen at low oxygen levels ✓ smaller reduction in hemoglobin saturation ✓ excretion of base via the kidneys to restore acid-base balance ✓ Muscle adaptations: reduced lean body mass ✓ increased capillary density, particularly in skeletal muscle to aid saturation ✓ increased mitochondrial density and enzyme concentration ✓ erythrpoietic response increases ✓ Cardiorespiratory adaptations:				2 max

Option B — Psychology of sport

C	uestion	Answers	Notes	Total
5.	а	control ✓		1
5.	b	38.33–32.21 ✓ = 6.12 «kg» ✓	Accept the subtraction in a different order.	2
5.	С	Data: imagery improved 1RM «5.88 kg» score more than the control group «0.91 kg» ✓ imagery improvement «5.88 kg» was «almost» as effective as physical practice improvement «6.12 kg» ✓ Theory: cognitive-based imagery aids task performance by improving focus / concentration ✓ cognitive-based imagery aids skill learning ✓ motivational-based imagery improves confidence ✓ imagery can be used to improve motivation ✓	Mere presentation of figures from table without stating improvement is not sufficient for mark. Reference to numbers must be the difference in values. Award [2 max] for theoretical points.	3 max

-7 -

Q	uestion	Answers	Notes	Total
6.	а	the internal mechanisms and external stimuli which arouse and direct our behaviour ✓	Accept other appropriate definitions.	1
6.	b	extrinsic rewards can be a controlling influence on behaviour \checkmark extrinsic/controlling rewards reduce intrinsic motivation «while possibly increasing extrinsic motivation» \checkmark extrinsic rewards seen as information providing feedback on performance \checkmark information rewards can increase intrinsic motivation \checkmark intrinsic motivation leads to greater satisfaction with performance therefore satisfaction may be decreased with extrinsic rewards \checkmark		3 max

-8-

7.	а				Accept any appropriate example containing reference to judgement against a self-referenced numeric value.	1	
7.	b	characterised by thoughts characterised by physiological response involves worry and self-doubt involves butterflies, dry mouth, shaking, etc. typically increases immediately before	Cognitive Y N Y N Y N	Somatic N Y N N Y	✓ ✓ ✓ ✓		2 max
7.	С	progressive muscular relaxation (PMR) reduce breathing techniques can control heart rate and tension ✓	preathing techniques can control heart rate and respiratory rate, reducing			Award [1 max] for list. Accept other techniques as appropriate.	2 max

	self-talk ✓	
	mental imagery ✓	
	thought stopping 🗸	
	music √	

-9-

C	uestion	Aı	nswers			Notes	Total
8.	а		Talent identification (TI)	Multidimensional talent identification and development (TID)			
		predominantly physiological assessments	Y	Ň	✓		3 max
		incorporates psychological skill	N	Υ	✓		
		recognises the evolution of talent	N	Υ	✓		
		monitoring takes place as a snapshot	Υ	N	✓		
		monitoring takes place over time	N	Υ	✓		
8.	b	stages were initiation, development, maduring early stages, practice is coach- of as athletes master skill, practice become development is dependent on opportunity	oloom (1985) / Cote (1999) suggested stages of athlete development ✓ stages were initiation, development, mastery, and maintenance / perfection ✓ during early stages, practice is coach- or parent-led ✓ as athletes master skill, practice becomes more self-determined ✓ development is dependent on opportunities, obstacles, and progressions ✓ psychological skill is key to coping with unstable periods and determines				3 max

C	uestion	Answers	Notes	Total
8.	c	relationship is bi-directional, in that motivated athletes engage in more self-regulated learning, which in turn serves to enhance motivation ✓ stages forethought, monitoring, reflection Forethought (planning) phase: athletes must see value in task to spend time planning for it ✓	Notes	Total
		higher self-efficacy beliefs increase the use of self-regulation strategies ✓ Monitoring phase: intrinsic motivation affects level of effort in completing tasks and use of self-regulation strategies ✓ Reflection phase: athlete attributions affect future engagement in self-regulation strategies ✓		4 max

Option C — Physical activity and health

C	Questi	ion	Answers	Notes	Total
9.	а	i	Southeast Asia ✓		1
9.	а	ii	60-30 ✓ = 30 «%» ✓	Accept the subtraction in a different order.	2
9.	a	iii	Compare: adults aged ≥ 60 highest proportion of physical inactivity in both regions ✓ from 30+, there is a similar trend in increasing levels of inactivity ✓ Contrast: when comparing each age group, Americas have more inactivity than Western Pacific ✓ inactivity increases with age in Americas but does not in Western Pacific ✓ OR there is a greater increase in inactivity from 30–44 to 45–59 and to >60 in Americas compared to Western Pacific ✓ 15–29 year olds are the most active / least inactive in Americas but they are more inactive than 30–44 years and 45–59 years in Western Pacific ✓	Award [2 max] for contrast. Must be clear that comparison is age group to same age group between regions. Accept other appropriate interpretations.	3 max
9.	b		a condition that involves narrowing or blockage of blood vessels that supply the heart «leading to heart attack / angina» ✓		1

– 11 –

	1	2	
_	- 1	2	_

Question		Answers	Notes	Total
9.	С	Inactive individuals are more likely to have:	Award [2 max] for list.	
		high blood pressure √		
		atherosclerosis ✓		
		obesity ✓		3 max
		type 2 diabetes ✓		
		low HDL-cholesterol ✓		

10.	а	a state of emotional or affective arousal of varying, and not permanent duration 🗸	Accept other appropriate definitions.	1	
10.	b	exercise seems to have a positive correlation with alleviating depression \checkmark being sedentary has been shown to be related to higher levels of depression \checkmark no causal link has been established between exercise and depression \checkmark			
		exercise involving rhythmical abdominal breathing is likely to have a positive effect \checkmark		4 may	
		exercise at least 20 minutes in duration is likely have a positive effect \checkmark exercise may increase the release of endorphins / increase serotonin / norepinephrine synthesis \checkmark exercise groups convey a sense of mastery and increased self-esteem \checkmark may also provide social interaction and promote social support \checkmark		4 max	

C	uestion	Answers	Notes	Total
11.	a	Acute injuries: occur suddenly as a result of a specific injury mechanism ✓ Chronic injuries: develop over a period of time and are often caused by repetitive activity ✓		2
11.	b	rapid increase in training distance or intensity \(\square \) warm-up \(\square \) ignoring warning signs of discomfort can lead to overuse injuries \(\square \) running technique \(\square \) twists and turns \(\square \) running surface \(\section \) footwear \(\section \) footwear \(\section \) previous injuries \(\section \) gight muscles straining tendons \(\square \) running experience \(\section \) provious injuries \(\section \) running experience \(\section \) provious provious injuries \(\section \) running experience \(\section \) provious provious injuries \(\section \) running experience \(\section \) provious injuries \(\section \) running experience \(\section \) provious provious injuries \(\section \) running experience \(\section \) provious provious injuries \(\section \) running experience \(\section \) running experience \(\section \) provious provious injuries \(\section \) running experience \(\section \) running experience \(\section \) provious provious injuries \(\section \) running experience \(\section \		3 max

– 13 –

C	uestion	Answers	Notes	Total
12.	а	underlying medical history / genetic disorders ✓ underlying problem when accompanied with high intensity exercise ✓ underlying problem when accompanied with habitual weekly exercise ✓		2 max
12.	b	moderate exercise (eg walking) is associated with a lower risk of mortality ✓ improved metabolic rates and VO2max improves aerobic capacity ✓ increased energy expenditure reduces risk of obesity ✓ improved plasma lipid profiles reduce risk of atherosclerosis ✓ decreased adiposity reduces risk of atherosclerosis ✓ decreased blood pressure reduces risk of cardiovascular disease ✓ reduced risk of skeletal injuries and potential periods of physical inactivity ✓ Social well-being eg, walking with groups/friends ✓ Psychological benefits eg, increased self-esteem from losing weight ✓		3 max

– 14 –

Option D — Nutrition for sport, exercise and health

Q	uestic	on	Answers	Notes	Total
13.	а		1.55 «minutes» ✓		1
13.	b		29.49–26.46 «minutes» ✓ = 3.03 «minutes» ✓	Accept subtraction in a different order.	2
13.	С		there was no significant change in running or cycling performance from start to end of study for the control group ✓	Award [2 max] if no conclusion.	
			there was no significant change in running performance from start to end of study for the early consumption group \checkmark		
			there was a significant change / improvement in cycling performance between start and end of study for the early consumption group ✓		3 max
			Conclusion:		
			early carbohydrate consumption may be beneficial in some activities/sport ✓		

14.	а	pepsin ✓ trypsin ✓	Two required in list to award [1] mark.	1 max
14.	b	a catalyst for the breakdown of large food molecules into smaller molecules ✓ smaller molecules are more soluble ✓ substances, which can be absorbed from the gut into the bloodstream ✓ speed up the rate of digestion ✓ carbohydrates are acted on by amylase ✓ OR proteins are acted on by pepsin ✓ OR fats are acted on by lipase ✓	Award 1 [max] for specific example. Accept other relevant examples.	3 max

– 16 **–**

Q	Question		Answers	Notes	Total
15.	а		low: apples/fish sticks/butter beans/kidney beans/lentils/sausage/fructose/peanuts/tuna ✓	N.B. The guide has brown rice as medium GI. Accept any suitable example (<15).	2
15.	b		Before race: simple sugars/high-GI foods immediately pre-competition causes blood sugar to rise rapidly/ excessive insulin release/ negatively impacts endurance ✓		
			low-GI immediately pre-competition has a slower rate of glucose absorption into the blood/ eliminates the insulin surge/ is beneficial for endurance ✓		3
			During race:		
			Triathlete should consume high-Gl «in optimal amounts» during a race ✓		

16.	а	Several causes but most commonly:	
		high insulin levels among diabetics ✓	
		Can also exist in non-diabetics through:	
		insufficient food intake ✓	
		excessive exercise ✓	3 max
		illness ✓	
		postponing or skipping a meal or snack	
		drinking alcohol	

– 17 –

Question		on	Answers	Notes	Total	
16.	b		glucose uptake into a cell is facilitated by the glucose transport proteins GLUT4 and GLUT1 ✓			
			GLUT4 transporters are stored inside intracellular vesicles that are translocated to cell membrane to allow greater glucose movement into cell ✓			
			GLUT4 transporters can be stimulated without insulin during physical exercise from other stimuli such as calcium ions ✓		3 max	
			glucose taken into muscle cells is quickly converted to glucose-6-phosphate to maintain glucose movement ✓			

17.		antioxidants are found in many foods and so are consumed as part of a healthy, well-balanced diet ✓		4 max
		free radicals are unstable and destructive to nearby molecules ✓		

OR
free radicals can: affect cell / mitochondrial membrane integrity / permeability / impair the function of molecules (<i>eg</i> enzymes) / impair DNA structure ✓
free radicals are a by-product of normal cell function that can lead to oxidative stress without sufficient antioxidants ✓
many athletes consume antioxidants in dietary supplements as extra defence against free radical damage ✓
no consistent evidence that these supplements reduce oxidative stress or have positive training or performance impact ✓
excess intake above RDA may have detrimental effects on the body 🗸
a lack of adequate regulation means that some products are poorly formulated and may even contain banned substances ✓