

**Mathematics**  
**Higher level**  
**Paper 3 – sets, relations and groups**

Thursday 15 November 2018 (afternoon)

1 hour

---

**Instructions to candidates**

- Do not open this examination paper until instructed to do so.
- Answer all the questions.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A graphic display calculator is required for this paper.
- A clean copy of the **mathematics HL and further mathematics HL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[50 marks]**.

Please start each question on a new page. Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working. For example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

1. [Maximum mark: 13]

Consider the binary operation  $*$  defined on the set of rational numbers,  $\mathbb{Q}$  by  $a*b = a + b + ab$ .

- (a) Show that the operation  $*$  is
  - (i) commutative;
  - (ii) associative. [5]
- (b) Solve the equation  $a*b = -1$ . [3]

Let  $S = \{x \in \mathbb{Q} \mid x \neq -1\}$ .

- (c) Show that  $\{S, *\}$  is an Abelian group. [5]

2. [Maximum mark: 8]

Consider two subsets  $X$  and  $Y$  of a universal set  $U$ .

- (a) Use De Morgan's laws to prove that  $\left[ (X' \cup Y')' \cap Y' \right]' = U$ . [4]

Let  $X = \{(m, n) \in \mathbb{Z} \times \mathbb{Z} \mid m = n^2\}$  and  $Y = \{(m, n) \in \mathbb{Z} \times \mathbb{Z} \mid 0 \leq m + n \leq 6\}$ .

- (b) List all elements of  $X \cap Y$ . [4]

3. [Maximum mark: 10]

A relation  $R$  is defined on  $\mathbb{R} \times \mathbb{R}$  by  $(a, b)R(c, d) \Leftrightarrow 3(a - c) = 2(b - d)$ .

- (a) Show that  $R$  is an equivalence relation. [8]
- (b) Find the equivalence class containing  $(1, 2)$  and describe it geometrically. [2]

4. [Maximum mark: 9]

Consider the functions  $f, g : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$  defined by

$$f((x, y)) = (x + y, x - y) \text{ and } g((x, y)) = (xy, x + y).$$

(a) Find

(i)  $(f \circ g)((x, y))$ ;

(ii)  $(g \circ f)((x, y))$ . [5]

(b) State with a reason whether or not  $f$  and  $g$  commute. [1]

(c) Find the inverse of  $f$ . [3]

5. [Maximum mark: 10]

Consider a finite group  $\{G, *\}$ . Let  $H$  be a subgroup of  $G$  of order  $m$  such that  $G \setminus H \neq \emptyset$ . Let  $a$  be a fixed element of  $G \setminus H$ . Consider the set  $A = \{a*h \mid h \in H\}$ .

(a) Show that  $A \cap H = \emptyset$ . [3]

Consider the function  $f: H \rightarrow A$  defined by  $f(h) = a*h$ .

(b) Show that  $f$  is a bijection. [4]

(c) Find the number of elements in the set  $A \cup H$ . [3]