

Markscheme

November 2018

Statistics and probability

Higher level

Paper 3

10 pages

This markscheme is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Global Centre, Cardiff.

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a valid **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Mark according to RM™ Assessor instructions and the document “**Mathematics HL: Guidance for e-marking November 2018**”. It is essential that you read this document before you start marking. In particular, please note the following:

- Marks must be recorded using the annotation stamps. Please check that you are entering marks for the right question.
- If a part is **completely correct**, (and gains all the “must be seen” marks), use the ticks with numbers to stamp full marks.
- If a part is completely wrong, stamp **A0** by the final answer.
- If a part gains anything else, it **must** be recorded using **all** the annotations.
- All the marks will be added and recorded by RM™ Assessor.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, eg **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (eg substitution into a formula) and **A1** for using the **correct** values.
- Where the markscheme specifies (**M2**), **N3**, etc., do **not** split the marks.

- Once a correct answer to a question or part-question is seen, ignore further correct working. However, if further working indicates a lack of mathematical understanding do not award the final **A1**. An exception to this may be in numerical answers, where a correct exact value is followed by an incorrect decimal. However, if the incorrect decimal is carried through to a subsequent part, and correct **FT** working shown, award **FT** marks as appropriate but do not award the final **A1** in that part.

Examples

	Correct answer seen	Further working seen	Action
1.	$8\sqrt{2}$	5.65685... (incorrect decimal value)	Award the final A1 (ignore the further working)
2.	$\frac{1}{4}\sin 4x$	$\sin x$	Do not award the final A1
3.	$\log a - \log b$	$\log(a - b)$	Do not award the final A1

3 N marks

Award **N** marks for **correct** answers where there is **no** working.

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

4 Implied marks

Implied marks appear in **brackets** eg (**M1**), and can only be awarded if **correct** work is seen or if implied in subsequent working.

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 Follow through marks

Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (eg $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 Misread

If a candidate incorrectly copies information from the question, this is a misread (**MR**). A candidate should be penalized only once for a particular misread. Use the **MR** stamp to indicate that this has been a misread. Then deduct the first of the marks to be awarded, even if this is an **M** mark, but award all others so that the candidate only loses [**1 mark**].

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (eg $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).

7 Discretionary marks (**d**)

An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. In such cases the annotation **DM** should be used and a brief **note** written next to the mark explaining this decision.

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2 \sin(5x - 3)$, the markscheme gives:

$$f'(x) = (2 \cos(5x - 3))5 \quad (= 10 \cos(5x - 3)) \quad \mathbf{A1}$$

Award **A1** for $(2 \cos(5x - 3))5$, even if $10 \cos(5x - 3)$ is not seen.

10 Accuracy of Answers

Candidates should **NO LONGER** be penalized for an accuracy error (**AP**).

*If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy. When this is not specified in the question, all numerical answers should be given exactly or correct to three significant figures. Please check work carefully for **FT**.*

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

12 Calculators

A GDC is required for paper 3, but calculators with symbolic manipulation features (eg TI-89) are not allowed.

Calculator notation The mathematics HL guide says:

Students must always use correct mathematical notation, not calculator notation.

Do **not** accept final answers written using calculator notation. However, do not penalize the use of calculator notation in the working.

13 More than one solution

Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise.

1. (a) $E(2X + 7Y) = 2E(X) + 7E(Y) = 6 + 28 = 34$ (M1)A1
[2 marks]
- (b) $\text{Var}(X) = E(X) = 3$ and $\text{Var}(Y) = E(Y) = 4$ (R1)
 $\text{Var}(4X - 3Y) = 16\text{Var}(X) + 9\text{Var}(Y) = 48 + 36$ (M1)
 $= 84$ A1
[3 marks]
- (c) use of $E(U^2) = \text{Var}(U) + (E(U))^2$ (M1)
 $E(X^2) = 3 + 3^2; E(Y^2) = 4 + 4^2$ A1
 $E(X^2 - Y^2) = E(X^2) - E(Y^2)$ (M1)
 $= -8$ A1
[4 marks]
- Total [9 marks]**

2. (a) (i) $\bar{t} = \frac{\sum_{i=1}^{75} t_i}{75} = 28.866\dots = 28.9$ A1
- (ii) $s_{n-1}^2 = \frac{75}{74} \left(\frac{\sum_{i=1}^{75} t_i^2}{75} - \bar{t}^2 \right) = 188.9009\dots = 189$ (M1)A1

Note: Accept all answers that round to 28.9 and 189.

Note: Award **M0** if division by 75.

- (b) attempting to find a confidence interval. (M1)
- (i) 90% interval: (26.2, 31.5) A1
- (ii) 95% interval: (25.7, 32.0) A1

Note: Accept any values which round to within 0.1 of the correct value.

Note: Award **M1A1A0** if only confidence limits are given in the form 28.9 ± 2.6 .

- (c) 26 lies within the 95% interval but not within the 90% interval R1

Note: Award **R1** for considering whether or not one or two of the intervals contain 26.

the belief is supported at the 5% level (accept 95%) A1

the belief is not supported at the 10% level (accept 90%) A1

Note: **FT** their intervals but award **R1A1A0** if both intervals give the same conclusion.

[3 marks]

Total [9 marks]

3.

Note: Accept all answers that round to the correct 2sf answer in (a), (b) and (c) but not in (d).

(a) (i) $X \sim N(550, 8^2)$ (M1)
 $P(X > 560) = 0.10564\dots = 0.106$ A1

(ii) $X_i \sim N(550, 8^2), i = 1, \dots, 11$
let $Y = \sum_{i=1}^{11} X_i$
 $E(Y) = 11 \times 550$ (6050) A1
 $\text{Var}(Y) = 11 \times 8^2$ (704) (M1)A1
 $P(Y \leq 6000) = 0.02975\dots = 0.0298$ A1

[6 marks]

(b) (i) t distribution with 7 degrees of freedom A1A1
(ii) $p = 0.25779\dots = 0.258$ A2
(iii) $p > 0.05$ R1
therefore we conclude that there is no evidence to reject H_0 A1

Note: FT their p -value.

Note: Only award A1 if R1 awarded.

[6 marks]

(c) (i) $H_0 : \rho = 0, H_1 : \rho > 0$ A1

Note: Do not accept r in place of ρ .

(ii) $r = 0.782$ A2
(iii) $0.01095\dots = 0.0110$ A1
since $0.0110 < 0.05$ R1
there is positive association between weight and length A1

Note: FT their p -value.

Note: Only award A1 if R1 awarded.

Note: Conclusion must be in context.

[6 marks]

continued...

Question 3 continued

- (d) regression line of y (weight) on x (length) is **(M1)**
- $y = 0.8267\dots x + 255.96\dots$ **(A1)**
- $x = 360$ gives $y = 554$ **A1**

Note: Award **M1A0A0** for the wrong regression line, that is $y = 0.7393\dots x - 51.62\dots$

[3 marks]

Total [21 marks]

- 4. (a) $G_x(t) = \sum_{x=1}^{\infty} pq^{x-1}t^x$ or equivalent **M1A1**

Note: Condone omission of limits on summation.

$$= \sum_{x=1}^{\infty} pt(qt)^{x-1}$$

recognition of a geometric series **M1**

$$= \frac{pt}{1-qt}$$

AG

[3 marks]

- (b) $G'_x(t) = \frac{p(1-qt) + pqt}{(1-qt)^2}$ **M1A1**

$$= \frac{p}{(1-qt)^2}$$

$$E(X) = G'_x(1) = \frac{p}{(1-q)^2} = \frac{p}{p^2}$$

M1A1

$$= \frac{1}{p}$$

AG

[4 marks]

- (c) **METHOD 1**

$$G_y(t) = pt^5 + pqt^7 + pq^2t^9 + \dots$$

recognition of geometric series **M1A1**

(M1)

$$= \frac{pt^5}{1-qt^2}$$

A1

continued...

Question 4 continued

METHOD 2

$$G_Y(t) = E(t^Y) = E(t^{2X+3})$$

M1

$$= t^3 E\left((t^2)^X\right)$$

A1

$$= t^3 G_X(t^2)$$

(M1)

$$= \frac{pt^5}{1-qt^2}$$

A1

[4 marks]

Total [11 marks]
