

# Markscheme

November 2020

**Physics**

**Higher level**

**Paper 2**

19 pages

No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without written permission from the IB.

Additionally, the license tied with this product prohibits commercial use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, is not permitted and is subject to the IB's prior written consent via a license. More information on how to request a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite de l'IB.

De plus, la licence associée à ce produit interdit toute utilisation commerciale de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, n'est pas autorisée et est soumise au consentement écrit préalable de l'IB par l'intermédiaire d'une licence. Pour plus d'informations sur la procédure à suivre pour demander une licence, rendez-vous à l'adresse suivante : <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin que medie la autorización escrita del IB.

Además, la licencia vinculada a este producto prohíbe el uso con fines comerciales de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales— no está permitido y estará sujeto al otorgamiento previo de una licencia escrita por parte del IB. En este enlace encontrará más información sobre cómo solicitar una licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

Question			Answers	Notes	Total
1.	a	i	zero ✓		1
1	a	ii	Blades exert a downward force on the air ✓  air exerts an equal and opposite force on the blades «by Newton's third law» <b>OR</b> air exerts a reaction force on the blades «by Newton's third law» ✓	<i>Downward direction required for MP1.</i>	2
1	a	iii	«lift force/change of momentum in one second» = $1.7v$ ✓ $1.7v = (0.95 + 0.45) \times 9.81$ ✓ $v = 8.1 \text{ ms}^{-1}$ AND answer expressed to 2 sf only ✓	<i>Allow 8.2 from g = 10 ms<sup>-2</sup>.</i>	3
1	a	iv	<b>ALTERNATIVE 1</b>  power «=rate of energy transfer to the air = $\frac{1}{2} \frac{\Delta m}{\Delta t} v^2$ » = $\frac{1}{2} \times 1.7 \times 8.1^2$ ✓ $= 56 \text{ W}$ ✓  <b>ALTERNATIVE 2</b>  Power «= Force x v ave» = $(0.95 + 0.45) \times 9.81 \times \frac{8.1}{2}$ ✓ $= 56 \text{ W}$ ✓		2

Question		Answers	Notes	Total
1	b	<p>vertical force= lift force – weight <b>OR</b> = <math>0.45 \times 9.81</math> <b>OR</b> = 4.4 «N» ✓</p> <p>acceleration = <math>\frac{0.45 \times 9.81}{0.95} = 4.6</math> «ms<sup>-2</sup>» ✓</p>		2

Question		Answers	Notes	Total
2.	a	<p>arrow downwards labelled weight/W/mg and arrow upwards labelled friction/F ✓</p> <p>arrow horizontally to the left labelled «normal» reaction/N ✓</p>	<p><i>Ignore point of application of the forces but do not allow arrows that do not touch the object.</i></p> <p><i>Do not allow horizontal force to be labelled 'centripetal' or R.</i></p>	2
2	b	<p>See <math>F = \mu N</math> AND <math>N = mR\omega^2</math> ✓</p> <p>«substituting for N» <math>\mu m\omega^2 R = mg</math> ✓</p>		2

Question		Answers	Notes	Total
2	c	<p><b>ALTERNATIVE 1</b></p> <p>minimum required angular velocity <math>\ll= \sqrt{\frac{9.81}{0.40 \times 3.5}} \gg = 2.6 \text{ rad s}^{-1}</math> ✓</p> <p>actual angular velocity <math>\ll= \frac{2\pi}{\left(\frac{60}{28}\right)} \gg = 2.9 \text{ rad s}^{-1}</math> ✓</p> <p>actual angular velocity is greater than the minimum, so the person does not slide ✓</p> <p><b>ALTERNATIVE 2</b></p> <p>minimum friction force <math>= mg = \ll 9.81m \gg</math> ✓</p> <p>actual friction force <math>\ll= \mu m R \omega^2 = 0.40 \text{ m} \times 3.5 \left(2\pi \frac{28}{60}\right)^2 \gg = 12.0 \text{ m}</math> ✓</p> <p>actual friction force is greater than the minimum frictional force so the person does not slide ✓</p>	Allow 2.7 from $g = 10 \text{ ms}^{-2}$ .	3

Question			Answers	Notes	Total
3.	a	i	« $15 \times 30 \times 60$ » = 27000 «J» ✓		1
3	a	ii	$27 \times 10^3 = 0.32 \times c \times (290 - 250)$ OR 2100 ✓ $J \text{ kg}^{-1} \text{ K}^{-1}$ OR $J \text{ kg}^{-1} {}^\circ\text{C}^{-1}$ ✓	Allow any appropriate unit that is $\frac{\text{energy}}{\text{mass} \times \text{temperature}}$	2
3	b		«intermolecular» bonds are formed during freezing ✓  bond-forming process releases energy <b>OR</b> «intermolecular» PE decreases «and the difference is transferred as heat» ✓  «average random» KE of the molecules does not decrease/change ✓ temperature is related to «average» KE of the molecules «hence unchanged» ✓	To award MP3 or MP4 molecules/particles/atoms must be mentioned.	3 max
3	c		mass of frozen oil «= $\frac{27 \times 10^3}{130 \times 10^3}$ » = 0.21 «kg» ✓  unfrozen mass «= 0.32 - 0.21» = 0.11 «kg» ✓		2

Question		Answers	Notes	Total
4.	a	wavelength = $\frac{340}{850} = 0.40 \text{ «m»}$ ✓ path difference = 1.8 «m» ✓ $1.8 \text{ «m»} = 4.5\lambda$ <b>OR</b> $\frac{1.8}{0.20} = 9 \text{ «half-wavelengths»}$ ✓  waves meet in antiphase «at P» <b>OR</b> destructive interference/superposition «at P» ✓	Allow approach where path length is calculated in terms of number of wavelengths; along path A (56.25) and path B (60.75) for MP2, hence path difference 4.5 wavelengths for MP3	4
4	b	«equally spaced» maxima and minima ✓  a maximum at Q ✓  four «additional» maxima «between P and Q» ✓		2 max
4	c	the amplitude of sound at Q is halved ✓ «intensity is proportional to amplitude squared hence» $\frac{I_A}{I_0} = \frac{1}{4}$ ✓		2

Question			Answers	Notes	Total
4.	d	i	<p>speed of sound relative to the microphone is less ✓</p> <p>wavelength unchanged «so frequency is lower»</p> <p><b>OR</b></p> <p>fewer waves recorded in unit time/per second «so frequency is lower» ✓</p>		2
4	d	ii	$845 = 850 \times \frac{340 - v}{340} \quad \checkmark$ $v = 2.00 \text{ «m s}^{-1}\text{»} \quad \checkmark$		2


Question			Answers	Notes	Total
5.	a		<p>current is not «directly» proportional to the potential difference  <b>OR</b>          resistance of X is not constant  <b>OR</b>          resistance of X changes «with current/voltage» ✓</p>		1
5	b	i	<p><b>ALTERNATIVE 1</b></p> <p>voltage across X = 2.3 «V» ✓</p> <p>voltage across R «= <math>4.0 - 2.3</math>» = 1.7 «V» ✓</p> <p>resistance of variable resistor «= <math>\frac{1.7}{0.020}</math>» = 85 «Ω» ✓</p> <p><b>ALTERNATIVE 2</b></p> <p>overall resistance «= <math>\frac{4.0}{0.020}</math>» = 200 «Ω» ✓</p> <p>resistance of X «= <math>\frac{2.3}{0.020}</math>» = 115 «Ω» ✓</p> <p>resistance of variable resistor «= <math>200 - 115</math>» = 85 «Ω» ✓</p>		3
5	b	ii	power «= $4.0 \times 0.020$ » = 0.080 «W» ✓		1

Question			Answers	Notes	Total
5.	c	i	from 0 to 60 mA ✓		1
5	c	ii	<p><b>ALTERNATIVE 1</b></p> <p>current from the cell is greater «than 20 mA» ✓</p> <p>because some of the current must flow through section SQ of the potentiometer ✓</p> <p>overall power greater «than in part (b)» ✓</p> <p><b>ALTERNATIVE 2</b></p> <p>total/overall resistance decreases ✓</p> <p>because SQ and X are in parallel ✓</p> <p>overall power greater «than in part (b)» ✓</p>	<i>Allow the reverse argument.</i>	3


Question			Answers	Notes	Total
6.	a	i	energy required to «completely» separate the nucleons <b>OR</b> energy released when a nucleus is formed from its constituent nucleons ✓	Allow protons <b>AND</b> neutrons.	1
6	a	ii	the values «in SI units» would be very small ✓		1
6	a	iii	$140 \times 8.29 + 94 \times 8.59 - 235 \times 7.59$ <b>OR</b> 184 «MeV» ✓		1
6	b	i	see «energy => $180 \times 10^6 \times 1.60 \times 10^{-19}$ <b>AND</b> «mass => $235 \times 1.66 \times 10^{-27}$ ✓ $7.4 \times 10^{13}$ «J kg <sup>-1</sup> » ✓		2
6	b	ii	energy produced in one day = $\frac{1.2 \times 10^9 \times 24 \times 3600}{0.36} = 2.9 \times 10^{14}$ «J» ✓ mass = $\frac{2.9 \times 10^{14}}{7.4 \times 10^{13}} = 3.9$ «kg» ✓		2
6	b	iii	«specific energy of uranium is much greater than that of coal, hence» more energy can be produced from the same mass of fuel / per kg <b>OR</b> less fuel can be used to create the same amount of energy ✓		1

Question			Answers	Notes	Total
6	c	i	39 ✓	<i>Do not allow <math>^{94}_{39}X</math> unless the proton number is indicated.</i>	1
6	c	ii	75 «s» ✓		1
6.	c	iii	<p><b>ALTERNATIVE 1</b></p> <p><math>10 \text{ min} = 8 t_{1/2}</math> ✓</p> <p>mass remaining = <math>1.0 \times \left(\frac{1}{2}\right)^8 = 3.9 \times 10^{-3} \text{ «kg»}</math> ✓</p> <p><b>ALTERNATIVE 2</b></p> <p>decay constant = «<math>\frac{\ln 2}{75}</math>» = » <math>9.24 \times 10^{-3} \text{ «s}^{-1}</math> » ✓</p> <p>mass remaining = <math>1.0 \times e^{-9.24 \times 10^{-3} \times 600} = 3.9 \times 10^{-3} \text{ «kg»}</math> ✓</p>		2

Question			Answers	Notes	Total
7.	a		the «restoring» force/acceleration is proportional to displacement ✓	Allow use of symbols i.e. $F \propto -x$ or $a \propto -x$	1
7	b		Evidence of equating $m\omega^2x = \rho Agx$ «to obtain $\frac{\rho Ag}{m} = \omega^2$ » ✓ $\omega = \sqrt{\frac{1.03 \times 10^3 \times 2.29 \times 10^{-1} \times 9.81}{118}}$ OR 4.43 «rad s <sup>-1</sup> » ✓	Answer to at least 3 s.f.	2
7	c	i	« $E_k$ is a maximum when $x = 0$ hence» $E_{k,\max} = \frac{1}{2} \times 118 \times 4.4^2 (0.250^2 - 0^2)$ ✓ 71.4 «J» ✓		2

Question			Answers	Notes	Total
7.	c	ii	<p>energy never negative ✓</p> <p>correct shape with two maxima ✓</p> 		2

Question		Answers	Notes	Total
8.	a	<p><b>ALTERNATIVE 1</b>  work done on moving a positive test charge in any outward direction is negative ✓  potential difference is proportional to this work «so <math>V</math> decreases from A to B» ✓</p> <p><b>ALTERNATIVE 2</b>  potential gradient is directed opposite to the field so inwards ✓  the gradient indicates the direction of increase of <math>V</math> «hence <math>V</math> increases towards the centre/decreases from A to B» ✓</p> <p><b>ALTERNATIVE 3</b>  <math>V = \frac{kQ}{R}</math> so as <math>r</math> increases <math>V</math> decreases ✓  <math>V</math> is positive as <math>Q</math> is positive ✓</p> <p><b>ALTERNATIVE 4</b>  the work done per unit charge in bringing a positive charge from infinity ✓  to point B is less than point A ✓</p>		2

Question			Answers	Notes	Total
8.	b		<p>curve decreasing asymptotically for <math>r &gt; R</math> ✓</p> <p>non – zero constant between 0 and <math>R</math> ✓</p> 		2
8	c	i	<p><math>\ll W = \frac{1.7 \times 10^{-16}}{q} = \gg 1.1 \times 10^3 \ll V \gg</math> ✓</p>		1
8	c	ii	<p><math>8.99 \times 10^9 \times Q \times \left( \frac{1}{5.0 \times 10^{-2}} - \frac{1}{1.0 \times 10^{-1}} \right) = 1.1 \times 10^3</math> ✓</p> <p><math>Q = 1.2 \times 10^{-8} \ll C \gg</math> ✓</p>		2
8	d		to highlight similarities between «different» fields ✓		1

Question			Answers	Notes	Total
9.	a		there is a magnetic flux «linkage» in the coil / coil cuts magnetic field ✓ this flux «linkage» changes as the angle varies/coil rotates ✓ «Faraday's law» connects induced emf with rate of change of flux «linkage» with time ✓	<i>Do not award MP2 or 3 for answers that don't discuss flux.</i>	3
9	b	i	$V_{\text{rms}} = \frac{25 \times 10^3}{\sqrt{2}} \text{ «=} 17.7 \times 10^3 \text{ V »} \checkmark$ $I_{\text{rms}} = \frac{8.5 \times 10^5}{17.7 \times 10^3} = 48 \text{ «A »} \checkmark$		2
9	b	ii	«power loss proportional to $I^2$ hence the step-up factor is $\sqrt{2.5 \times 10^2} \text{ »} 16 \checkmark$		1
9	b	iii	peak emf doubles ✓ $T$ halves ✓	<i>Must show at least 1 cycle.</i>	2

Question			Answers	Notes	Total
10.	a		$\lambda = \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{1.60 \times 10^{-19} \times 4.2 \times 10^8}$ OR $= 2.96 \times 10^{-15}$ «m» ✓	Answer to at least 2 s.f. (i.e. 3.0)	1
10	b	i	«the shape of the graph suggests that» electrons undergo diffraction «with carbon nuclei» ✓ only waves diffract ✓		2
10	b	ii	$\sin \theta_0 = \frac{2.96 \times 10^{-15}}{4.94 \times 10^{-15}}$ «= 0.599» ✓ 37 «degrees» OR 0.64/0.65 «rad» ✓		2
10	b	iii	the de Broglie wavelength of electrons is «much» longer than the size of a nucleus ✓  hence electrons would not undergo diffraction <b>OR</b> no diffraction pattern would be observed ✓		2
10	c		volume of a nucleus proportional to $\left(A^{\frac{1}{3}}\right)^3 = A$ AND mass proportional to $A$ ✓  the ratio $\frac{\text{mass}}{\text{volume}}$ independent of $A$ «hence density the same for all nuclei» ✓	Both needed for MP1	2