

# Markscheme

November 2021

Chemistry

Standard level

Paper 2

15 pages

© International Baccalaureate Organization 2021

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2021

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2021

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

## Subject Details: Chemistry Standard Level Paper 2 Markscheme


Candidates are required to answer **ALL** questions. Maximum total = **[50 marks]**.

1. Each row in the “Question” column relates to the smallest subpart of the question.
2. The maximum mark for each question subpart is indicated in the “Total” column.
3. Each marking point in the “Answers” column is shown by means of a tick (✓) at the end of the marking point.
4. A question subpart may have more marking points than the total allows. This will be indicated by “**max**” written after the mark in the “Total” column. The related rubric, if necessary, will be outlined in the “Notes” column.
5. An alternative word is indicated in the “Answers” column by a slash (/). Either word can be accepted.
6. An alternative answer is indicated in the “Answers” column by “**OR**”. Either answer can be accepted.
7. An alternative markscheme is indicated in the “Answers” column under heading **ALTERNATIVE 1** etc. Either alternative can be accepted.
8. Words inside chevrons « » in the “Answers” column are not necessary to gain the mark.
9. Words that are underlined are essential for the mark.
10. The order of marking points does not have to be as in the “Answers” column, unless stated otherwise in the “Notes” column.
11. If the candidate’s answer has the same “meaning” or can be clearly interpreted as being of equivalent significance, detail and validity as that in the “Answers” column then award the mark. Where this point is considered to be particularly relevant in a question it is emphasized by **OWTTE** (or words to that effect) in the “Notes” column.
12. Remember that many candidates are writing in a second language. Effective communication is more important than grammatical accuracy.
13. Occasionally, a part of a question may require an answer that is required for subsequent marking points. If an error is made in the first marking point then it should be penalized. However, if the incorrect answer is used correctly in subsequent marking points then **follow through** marks should be awarded. When marking, indicate this by adding **ECF** (error carried forward) on the script.
14. Do **not** penalize candidates for errors in units or significant figures, **unless** it is specifically referred to in the “Notes” column.
15. If a question specifically asks for the name of a substance, do not award a mark for a correct formula unless directed otherwise in the “Notes” column. Similarly, if the formula is specifically asked for, do not award a mark for a correct name unless directed otherwise in the “Notes” column.
16. If a question asks for an equation for a reaction, a balanced symbol equation is usually expected, do not award a mark for a word equation or an unbalanced equation unless directed otherwise in the “Notes” column.
17. Ignore missing or incorrect state symbols in an equation unless directed otherwise in the “Notes” column.

Question		Answers	Notes	Total
1.	a	$\left\langle \frac{8.802 \text{ g}}{44.01 \text{ g mol}^{-1}} \Rightarrow 0.2000 \text{ «mol of C/CO}_2\text{»} \right\rangle$ <p><b>AND</b> <math>\left\langle \frac{3.604 \text{ g}}{18.02 \text{ g mol}^{-1}} \Rightarrow 0.2000 \text{ «mol of H}_2\text{O»} / 0.4000 \text{ «mol of H»} \right\rangle</math></p> <p><b>OR</b></p> $\left\langle \frac{8.802 \text{ g}}{44.01 \text{ g mol}^{-1}} \times 12.01 \text{ g mol}^{-1} \Rightarrow 2.402 \text{ «g of C»} \right\rangle$ <p><b>OR</b></p> $\left\langle \frac{3.604 \text{ g}}{18.02 \text{ g mol}^{-1}} \times 2 \times 1.01 \text{ g mol}^{-1} \Rightarrow 0.404 \text{ «g of H»} \checkmark \right\rangle$ <p><math>\left\langle 4.406 \text{ g} - 2.806 \text{ g} = 1.600 \text{ «g of O»} \checkmark \right\rangle</math></p> $\left\langle \frac{2.402 \text{ g}}{12.01 \text{ g mol}^{-1}} = 0.2000 \text{ mol C}; \frac{0.404 \text{ g}}{1.01 \text{ g mol}^{-1}} = 0.400 \text{ mol H}; \right\rangle$ $\frac{1.600 \text{ g}}{16.00 \text{ g mol}^{-1}} = 0.1000 \text{ mol O} \checkmark$ <p><b>C<sub>2</sub>H<sub>4</sub>O</b> ✓</p>	<p><i>Award [3] for correct final answer.</i></p>	<p><b>3</b></p>

Question		Answers	Notes	Total									
1.	b	$\llcorner \frac{88.12 \text{ g mol}^{-1}}{44.06 \text{ g mol}^{-1}} = 2 \llcorner \text{C}_4\text{H}_8\text{O}_2 \checkmark$	<i>C<sub>2</sub>S<sub>2</sub> if CS used.</i>	1									
1.	c	<table border="1"> <thead> <tr> <th>Spectrum</th> <th>Identity</th> <th>Reason</th> </tr> </thead> <tbody> <tr> <td>A</td> <td>Propan-1-ol</td> <td>                     absence of carbonyl/C=O «absorption»/ no peak in 1700 - 1750 «cm<sup>-1</sup>» range   <b>OR</b>                      presence of hydroxyl/O-H in <u>alcohols</u> «absorption»/peak in 3200 – 3600 «cm<sup>-1</sup>» range ✓                 </td> </tr> <tr> <td>B</td> <td>Propanoic acid</td> <td> <b>ALTERNATIVE 1:</b>                      carbonyl/C=O <b>AND</b> hydroxyl/O-H «in carboxylic acids absorptions»  <b>OR</b>                      «strong» peaks in 2500 – 3000 «cm<sup>-1</sup>» <b>AND</b> 1700 – 1750 «cm<sup>-1</sup>» ranges ✓   <b>ALTERNATIVE 2:</b>                      O-H in carboxylic acids «absorption» <b>AND</b> 2500 – 3000 «cm<sup>-1</sup>» range ✓   <b>ALTERNATIVE 3:</b>                      strong/broad «peak» <b>AND</b> 2500 – 3000 «cm<sup>-1</sup>» range ✓                 </td> </tr> </tbody> </table>	Spectrum	Identity	Reason	A	Propan-1-ol	absence of carbonyl/C=O «absorption»/ no peak in 1700 - 1750 «cm <sup>-1</sup> » range  <b>OR</b> presence of hydroxyl/O-H in <u>alcohols</u> «absorption»/peak in 3200 – 3600 «cm <sup>-1</sup> » range ✓	B	Propanoic acid	<b>ALTERNATIVE 1:</b> carbonyl/C=O <b>AND</b> hydroxyl/O-H «in carboxylic acids absorptions» <b>OR</b> «strong» peaks in 2500 – 3000 «cm <sup>-1</sup> » <b>AND</b> 1700 – 1750 «cm <sup>-1</sup> » ranges ✓  <b>ALTERNATIVE 2:</b> O-H in carboxylic acids «absorption» <b>AND</b> 2500 – 3000 «cm <sup>-1</sup> » range ✓  <b>ALTERNATIVE 3:</b> strong/broad «peak» <b>AND</b> 2500 – 3000 «cm <sup>-1</sup> » range ✓	<p><i>Award [1 max] for correctly identifying all 3 compounds without valid reasons given.</i></p> <p><i>Accept specific values of wavenumbers within each range.</i></p>	3
Spectrum	Identity	Reason											
A	Propan-1-ol	absence of carbonyl/C=O «absorption»/ no peak in 1700 - 1750 «cm <sup>-1</sup> » range  <b>OR</b> presence of hydroxyl/O-H in <u>alcohols</u> «absorption»/peak in 3200 – 3600 «cm <sup>-1</sup> » range ✓											
B	Propanoic acid	<b>ALTERNATIVE 1:</b> carbonyl/C=O <b>AND</b> hydroxyl/O-H «in carboxylic acids absorptions» <b>OR</b> «strong» peaks in 2500 – 3000 «cm <sup>-1</sup> » <b>AND</b> 1700 – 1750 «cm <sup>-1</sup> » ranges ✓  <b>ALTERNATIVE 2:</b> O-H in carboxylic acids «absorption» <b>AND</b> 2500 – 3000 «cm <sup>-1</sup> » range ✓  <b>ALTERNATIVE 3:</b> strong/broad «peak» <b>AND</b> 2500 – 3000 «cm <sup>-1</sup> » range ✓											


			<p>C</p> <p>Propanal</p> <p>presence of carbonyl/C=O «absorption»/ peak in 1700 – 1750 «cm<sup>-1</sup>» range</p> <p><b>AND</b></p> <p>absence of hydroxyl/O-H «in carboxylic acids absorption»/ no «broad» peak in 2500 – 3000 «cm<sup>-1</sup>» range ✓</p>		
--	--	--	--	--	--

Question			Answers	Notes	Total
2.			increasing number of protons <b>OR</b> increasing nuclear charge ✓  «atomic» radius/size decreases <b>OR</b> same number of shells/electrons occupy same shell <b>OR</b> similar shielding «by inner electrons» ✓		2
3.	a	i		Accept any diagram with each P joined to the other three.  Accept any combination of dots, crosses and lines.	1
3.	a	ii	$\text{P}_4(\text{s}) + 6\text{Cl}_2(\text{g}) \rightarrow 4\text{PCl}_3(\text{l}) \quad \checkmark$		1
3.	b	i	Electron domain geometry: tetrahedral ✓ Molecular geometry: trigonal pyramidal ✓ Bond angle: 100«°» ✓	Accept any value or range within the range 91–108«°» for M3.	3

Question			Answers	Notes	Total
3.	b	ii	polar <b>AND</b> unsymmetrical distribution of charge <b>OR</b> polar <b>AND</b> dipoles do not cancel <b>OR</b> «polar as» dipoles «add to» give a «partial» positive «charge» at P and a «partial» negative «charge» at the opposite/Cl side of the molecule ✓	Accept “polar <b>AND</b> unsymmetrical molecule”.	1
3.	c	i	«-398.9 kJ mol <sup>-1</sup> - (-306.4 kJ mol <sup>-1</sup> ) =» -92.5 «kJ mol <sup>-1</sup> » ✓		1
3.	c	ii	«K <sub>c</sub> =» $\frac{[\text{PCl}_5]}{[\text{PCl}_3][\text{Cl}_2]}$ ✓		1
3.	c	iii	«shifts» left/towards reactants <b>AND</b> «forward reaction is» exothermic/ΔH is negative ✓		1



Question			Answers	Notes	Total
4.	a	i	«nucleophilic» substitution/S <sub>N</sub> 2 ✓	<i>Do not accept if “electrophilic” or “free radical” substitution is stated.</i>	1
4.	a	ii	«acts as a» nucleophile/Lewis base <b>OR</b> donates/provides lone pair «of electrons» <b>OR</b> attacks the «partially» positive carbon ✓		1
4.	a	iii	bond enthalpy C–I lower than C–Cl <b>OR</b> C–I bond weaker than C–Cl ✓  «weaker bond» broken more easily/with less energy <b>OR</b> lower $E_a$ «for weaker bonds» ✓	<i>Accept the bond enthalpy values for C–I and C–Cl for M1.</i>	2










Question			Answers	Notes	Total
4.	b	i	 <p>peak at <math>T_1</math> to right of <b>AND</b> lower than <math>T_2</math> ✓                      lines begin at origin <b>AND</b> <math>T_1</math> must finish above <math>T_2</math> ✓</p>		2

Question			Answers	Notes	Total
4.	b	ii	<p>«rate is» lower <b>AND</b> «average» kinetic energy of molecules is lower</p> <p><b>OR</b></p> <p>«rate is» lower <b>AND</b> less frequent collisions</p> <p><b>OR</b></p> <p>«rate is» lower <b>AND</b> fewer collisions per unit time ✓</p> <p>«rate is» lower <b>AND</b> fewer/smaller fraction of molecules/collisions have the <math>E \geq E_a</math> ✓</p>	<p><i>Lower «rate» needs to be mentioned once only.</i></p> <p><i>Do not accept “fewer collisions” without reference to time/frequency/probability for M1.</i></p>	2
5.		a	$\text{H}_3\text{PO}_4(\text{aq}) + \text{NaOH}(\text{aq}) \rightarrow \text{NaH}_2\text{PO}_4(\text{aq}) + \text{H}_2\text{O}(\text{l}) \checkmark$	<p><i>Accept net ionic equation.</i></p>	1
5.		b	$\text{H}_2\text{PO}_4^-(\text{aq}) + \text{H}^+(\text{aq}) \rightarrow \text{H}_3\text{PO}_4(\text{aq}) \checkmark$ $\text{H}_2\text{PO}_4^-(\text{aq}) + \text{OH}^-(\text{aq}) \rightarrow \text{HPO}_4^{2-}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \checkmark$	<p><i>Accept reactions of <math>\text{H}_2\text{PO}_4^-</math> with any acidic, basic or amphoteric species, such as <math>\text{H}_3\text{O}^+</math>, <math>\text{NH}_3</math> or <math>\text{H}_2\text{O}</math>.</i></p> <p><i>Accept <math>\text{H}_2\text{PO}_4^-(\text{aq}) \rightarrow \text{HPO}_4^{2-}(\text{aq}) + \text{H}^+(\text{aq})</math> for M2.</i></p>	2

Question		Answers	Notes	Total
5.	c	$\llcorner \text{NaOH} \frac{28.40 \text{ cm}^3}{1000} \times 0.5000 \text{ mol dm}^{-3} = 0.01420 \text{ mol} \llcorner$ $\llcorner \frac{0.01420 \text{ mol}}{3} \Rightarrow 0.004733 \llcorner \text{mol} \llcorner \checkmark$ $\llcorner \frac{0.004733 \text{ mol}}{\frac{25.00 \text{ cm}^3}{1000}} \Rightarrow 0.1893 \llcorner \text{mol dm}^{-3} \llcorner \checkmark$	<i>Award [2] for correct final answer.</i>	2
5.	d	$\llcorner \text{OH}^- \text{ is a} \llcorner \text{ proton acceptor} \llcorner \checkmark$		1
6	a	$\llcorner \text{amount of} \llcorner \text{ oxygen used to decompose the organic matter in water} \llcorner \checkmark$		1
6.	b	$\llcorner \frac{0.0001 \text{ g}}{0.1240 \text{ g}} \times 100 \% \Rightarrow 0.08 \llcorner \% \llcorner$ <p><b>OR</b></p> $\llcorner \frac{0.4 \text{ cm}^3}{1000.0 \text{ cm}^3} \times 100 \% \Rightarrow 0.04 \llcorner \% \llcorner \checkmark$ $\llcorner 0.08 \% + 0.04 \% \Rightarrow 0.12/0.1 \llcorner \% \llcorner \checkmark$	<i>Award [2] for correct final answer.</i> <i>Accept fractional uncertainties for M1, i.e., 0.0008 OR 0.0004.</i>	2

Question			Answers	Notes	Total
6.	c	i	$\ll \frac{37.50 \text{ cm}^3}{1000} \times 5.000 \times 10^{-4} \text{ mol dm}^{-3} \Rightarrow 1.875 \times 10^{-5} \text{ «mol» } \checkmark$		1
6.	c	ii	1:4 $\checkmark$	Accept "4 mol S <sub>2</sub> O <sub>3</sub> <sup>2-</sup> : 1 mol O <sub>2</sub> ", but not just 4:1.	1
6.	c	iii	$\ll 1.875 \times 10^{-5} \text{ mol} \times \frac{1}{4} \Rightarrow 4.688 \times 10^{-6} \text{ «mol» } \checkmark$ $\ll \frac{4.688 \times 10^{-6} \text{ mol}}{\frac{25.00 \text{ cm}^3}{1000}} \Rightarrow 1.875 \times 10^{-4} \text{ «mol dm}^{-3}\text{» } \checkmark$	Award [2] for correct final answer.	2
6.	c	iv	$\text{MnO}_2(\text{s}) + 2\text{e}^- + 4\text{H}^+(\text{aq}) \rightarrow \text{Mn}^{2+}(\text{aq}) + 2\text{H}_2\text{O}(\text{l}) \checkmark$		1
7.		a	$\ll q = mc\Delta T = 20.0 \text{ g} \times 4.18 \text{ J g}^{-1} \text{ }^\circ\text{C}^{-1} \times 57.3 \text{ }^\circ\text{C} \Rightarrow 4790 \text{ «J» } \checkmark$ $\ll \Delta H_c = -\frac{4790 \text{ J}}{8.75 \times 10^{-4} \text{ mol}} \Rightarrow -5470 \text{ «kJ mol}^{-1}\text{» } \checkmark$	Award [2] for correct final answer. Accept answers in the range -5470 to -5480 «kJ mol <sup>-1</sup> ». Accept correct answer in any units, e.g. - 5.47 «MJ mol <sup>-1</sup> » or 5.47 x 10 <sup>6</sup> «J mol <sup>-1</sup> ».	2

Question		Answers	Notes	Total
7.	b	$\text{Cl}\cdot + \text{C}_2\text{H}_6 \rightarrow \cdot\text{C}_2\text{H}_5 + \text{HCl} \checkmark$ $\cdot\text{C}_2\text{H}_5 + \text{Cl}_2 \rightarrow \text{Cl}\cdot + \text{C}_2\text{H}_5\text{Cl} \checkmark$  $\cdot\text{C}_2\text{H}_5 + \text{Cl}\cdot \rightarrow \text{C}_2\text{H}_5\text{Cl}$ <b>OR</b> $\text{Cl}\cdot + \text{Cl}\cdot \rightarrow \text{Cl}_2$ <b>OR</b> $\cdot\text{C}_2\text{H}_5 + \cdot\text{C}_2\text{H}_5 \rightarrow \text{C}_4\text{H}_{10} \checkmark$	<p><i>Do not penalize incorrectly placed radical sign, eg <math>\text{C}_2\text{H}_5\cdot</math>.</i></p>	3
8.	a	<p><i>Most <math>{}^4\text{He}^{2+}</math> passing straight through:</i>                      most of the atom is empty space  <b>OR</b>                      the space between nuclei is much larger than <math>{}^4\text{He}^{2+}</math> particles  <b>OR</b>                      nucleus/centre is «very» small «compared to the size of the atom» <math>\checkmark</math></p> <p><i>Very few <math>{}^4\text{He}^{2+}</math> deviating largely from their path:</i>                      nucleus/centre is positive «and repels <math>{}^4\text{He}^{2+}</math> particles»  <b>OR</b>                      nucleus/centre is «more» dense/heavy «than <math>{}^4\text{He}^{2+}</math> particles and deflects them»  <b>OR</b>                      nucleus/centre is «very» small «compared to the size of the atom» <math>\checkmark</math></p>	<p><i>Do not accept the same reason for both M1 and M2.</i>                      Accept “most of the atom is an electron cloud” for M1.</p> <p><i>Do not accept only “nucleus repels <math>{}^4\text{He}^{2+}</math> particles” for M2.</i></p>	2

Question			Answers	Notes	Total						
8.	b	i	<table border="1" style="width: 100%; text-align: center;"> <tr> <td></td> <td></td> <td></td> </tr> <tr> <td>1s</td> <td>2s</td> <td>2p</td> </tr> </table> <p>1s <b>AND</b> 2s as spheres ✓                      one or more 2p orbital(s) as figure(s) of 8 shape(s) of any orientation (p<sub>x</sub>, p<sub>y</sub> p<sub>z</sub>) ✓</p>				1s	2s	2p		<b>2</b>
											
1s	2s	2p									
8.	b	ii	<p>1s<sup>2</sup>2s<sup>2</sup>2p<sup>6</sup>3s<sup>2</sup>3p<sup>6</sup>4s<sup>1</sup>3d<sup>10</sup>  <b>OR</b>                      [Ar] 4s<sup>1</sup>3d<sup>10</sup> ✓</p>	Accept configuration with 3d before 4s.	<b>1</b>						