

Markscheme

November 2021

Mathematics: applications and interpretation

Higher level

Paper 1



© International Baccalaureate Organization 2021

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2021

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2021

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

Instructions to Examiners

Abbreviations

- **M** Marks awarded for attempting to use a correct **Method**.
- A Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- **R** Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- **FT** Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do not automatically award full marks for a correct answer; all working must be checked, and marks awarded according to the markscheme.
- It is generally not possible to award MO followed by A1, as A mark(s) depend on the preceding M mark(s), if any.
- Where **M** and **A** marks are noted on the same line, e.g. **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **A1** for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the **AG** line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this
 working is incorrect and/or suggests a misunderstanding of the question. This will encourage a
 uniform approach to marking, with less examiner discretion. Although some candidates may be
 advantaged for that specific question item, it is likely that these candidates will lose marks
 elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used in a subsequent part. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part. Examples:

	Correct	Further	Any FT issues?	Action
	answer seen	working seen		Action
1.		5.65685	No.	Award A1 for the final mark
	$8\sqrt{2}$	(incorrect	Last part in question.	(condone the incorrect further
	·	decimal value)		working)
2.	35	0.468111	Yes.	Award A0 for the final mark
	$\frac{33}{72}$	(incorrect	Value is used in	(and full FT is available in
	1/2	decimal value)	subsequent parts.	subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (M1), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is **(M1)A1**, it is possible to award full marks for *their* correct answer, **without working being seen**. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (e.g. probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (*MR*). A candidate should be penalized only once for a particular misread. Use the *MR* stamp to indicate that this has been a misread and do not award the first mark, even if this is an *M* mark, but award all others as appropriate.

- If the question becomes much simpler because of the MR, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- **MR** can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by METHOD 1, METHOD 2, etc.
- Alternative solutions for parts of questions are indicated by **EITHER** . . . **OR**.

7 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1,000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "from the use of 3 sf values".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an **A** mark to be awarded, arithmetic should be completed, and any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$.

An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

A GDC is required for this paper, but If you see work that suggests a candidate has used any calculator not approved for IB DP examinations (eg CAS enabled devices), please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

1. (a)
$$m = \frac{6-0}{4-2} = 3$$

(M1)A1

[2 marks]

(b)
$$(m=)$$
 $-\frac{1}{3}$ $(-0.333, -0.333333...)$

A1

[1 mark]

(c) an equation of line with a correct intercept and either of their gradients from (a) or (b)

(M1)

e.g.
$$y = -\frac{1}{3}x + 4$$
 OR $y - 4 = -\frac{1}{3}(x - 0)$

Note: Award *(M1)* for substituting either of their gradients from parts (a) or (b) and point B or (3, 3) into equation of a line.

$$x+3y-12=0$$
 or any integer multiple

A1

[2 marks]

(d)
$$(x =) 12$$

A1

[1mark] Total: [6 marks]

2. (a) attempt at using trapezoidal rule formula

(M1)

$$\frac{1}{2} \left(\frac{2-0}{5} \right) \left(30 + 50 + 2(50 + 60 + 40 + 20) \right)$$

A1

(total carbon =) 84 tonnes

A1 [3 marks]

(b)
$$\left| \frac{84-72}{72} \right| \times 100\%$$

(M1)

Note: Award *(M1)* for correct substitution of final answer in part (a) into percentage error formula.

A1

[2 marks] Total: [5 marks]

3. (a)
$$h(4) = \frac{640}{4^2} + 0.5$$
 OR $h(14) = \frac{640}{14^2} + 0.5$ (M1)

Note: Award *(M1)* for substituting 4 or 14 into h. This can be implicit from seeing 3.77 (3.76530...) or 40.5.

$$3.77 \le h(x) \le 40.5 \quad (3.76530... \le h(x) \le 40.5)$$

A1A1

Note: Award A1 for both correct endpoints seen, A1 for the endpoints in a correct interval.

[3 marks]

(b) (i)
$$h(x) = 10$$
 OR $h^{-1}(x) = \sqrt{\frac{640}{x - 0.5}}$ OR $h^{-1}(10) = \sqrt{\frac{640}{10 - 0.5}}$ (M1)
 $(x =) 8.21 \text{ cm } (8.20782...)$

(ii) a tin that is 10 cm high will have a diameter of 8.21 cm (8.20782...)

Note: Condone a correct answer expressed as the converse.

(iii)
$$4 \le h^{-1} \le 14$$

Note: Accept $4 \le y \le 14$. Do not *FT* in this part.

[4 marks] Total: [7 marks]

$$AP = \sqrt{(10-6)^2 + (6-2)^2} \quad \text{OR}$$

$$BP = \sqrt{(10-14)^2 + (6-2)^2} \quad \text{OR}$$

$$DP = \sqrt{(10-10.8)^2 + (6-11.6)^2} \quad OR$$

$$BQ = \sqrt{(13-14)^2 + (7-2)^2} \quad OR$$

$$CQ = \sqrt{(13-18)^2 + (7-6)^2}$$
 OR

$$DQ = \sqrt{(13-10.8)^2 + (7-11.6)^2}$$

(AP or BP or DP =)
$$\sqrt{32} = 5.66$$
 (5.65685...) **AND**

(BQ or CQ or DQ =)
$$\sqrt{26} = 5.10 (5.09901...)$$

A1

$$\sqrt{32} > \sqrt{26}$$
 OR AP (or BP or DP) is greater than BQ (or CQ or DQ)

A1

AG

Note: Follow through from their values provided their AP (or BP or DP) is greater than their BQ (or CQ or DQ).

[3 marks]


continued...

Question 4 continued

(b) (i) x = 14

A1

(ii)



A1A1

Note: Award A1 for each correct straight line. Do not FT from their part b(i).

[3 marks] Total: [6 marks]

5. (a)
$$N = 360$$

 $I\% = 3.8$
 $PV = (\pm)170\,000$
 $FV = 0$
 $P/Y = 12$
 $C/Y = 12$

(M1)(A1)

Note: Award *(M1)* for an attempt to use a financial app in their technology with at least two entries seen, award A1 for all entries correct. Accept a positive or negative value for PV.

$$(PMT =) 792.13 \text{ AUD}$$

A1

Note: Accept an answer of -792.13. Do not award final **A1** if answer is not given correct to 2 dp

[3 marks]

(b) (i)
$$N = 120$$

 $I\% = 3.8$
 $PV = (\pm)170000$
 $PMT = (\mp)792.13$
 $P/Y = 12$
 $C/Y = 12$

(M1)(A1)

Note: Award *(M1)* for an attempt to use a financial app in their technology with a least two entries seen, award *A1* for all entries correct. PV and PMT must have opposite signs.

$$(FV =) 133019.94 \text{ AUD}$$

A1

A1

Note: Do not award final *A1* if answer is not given correct to 2 dp, unless already penalized in part (a). Accept 133020.30 from use of exact value for PMT.

interest paid: (95055.60 – 36980.06 =) 58075.54 AUD

Note: Allow 58075.60 or 58075.90 from use of some exact values from parts (a) and (b)(i). If their answer to part (b)(i) is greater than 170 000 then award at most *(M1)(M1)(A0)* for follow through in part (b)(ii).

[6 marks] Total: [9 marks]

https://xtremepape.rs/

6. (a)
$$10 = \frac{2}{1-r}$$
 (M1) $r = 0.8$

[2 marks]

(b)
$$2 \times (0.8)^{n-1} < 0.5$$
 OR $2 \times (0.8)^{n-1} = 0.5$ (M1) $(n >) 7.212....$ (A1) $n = 8$

Note: If n = 7 is seen, with or without seeing the value 7.212... then award **M1A1A0**.

[3 marks]

Total: [5 marks]

(b) recognition that all entries add up to 120 (M1) a = 120 - 6 - 13 - 26 - b OR a = 75 - b A1 [2 marks]

(c) (i)
$$\frac{6 \times 1 + 13 \times 2 + 26 \times 3 + (75 - b) \times 4 + b \times 5}{120} = 3.65$$
 (M1)(A1)

Note: Award *(M1)* for attempt to substitute into mean formula, LHS expression is sufficient for the *M* mark. Award *(A1)* for correct substitutions in one variable **OR** in two variables, followed by evidence of solving simultaneously with a + b = 75.

$$(b=) 28$$


(ii) 120 – their part (c)(i) seen (e.g. 92 indicated on graph) (M1) 84

[5 marks] Total: [8 marks]

- **8**. (a) (i) *a*
 - (ii) the hill is at its steepest / largest slope of hill


 A1

 [2 marks]



Note: Award **(A1)** for decreasing function from 0 to b and d to f and increasing from b to d; **(A1)** for minimum at b and max at d; **(A1)** for starting at height of 60 and finishing at a height of 0 at f. If reasonable curvature not evident on graph (i.e. only straight lines used) award **A1A0A1**.

[3 marks] Total: [5 marks] **9.** (a)



A1A1

[2 marks]

(b)
$$A = \begin{pmatrix} 0.8 & 0.6 \\ 0.2 & 0.4 \end{pmatrix}$$
 (A1)


$$A^{180} = \begin{pmatrix} 0.75 & 0.75 \\ 0.25 & 0.25 \end{pmatrix} \tag{M1}$$

0.75

[3 marks] Total: [5 marks] 10. (a) g(0) = 16 M1A1

[2 marks]

(b)



y-asymptote (y = 4)

concave up decreasing curve and passing through $(0,\,16)$

A1 A1

[2 marks] Total: [4 marks]

11. METHOD 1

attempt to find AC using cosine rule
$$7^2 = 10^2 + AC^2 - 2 \times 10 \times AC \times \cos 40^\circ$$
 (A1)

attempt to solve a quadratic equation (M1)

$$AC = 4.888...$$
 AND $10.432...$ (A1)

Note: At least AC = 4.888... must be seen, or implied by subsequent working.

minimum area =
$$\frac{1}{2} \times 10 \times 4.888... \times \sin(40^\circ)$$
 M1

Note: Do not award *M1* if incorrect value for minimizing the area has been chosen.

$$=15.7 \text{ m}^2$$

METHOD 2

attempt to find \hat{ACB} using the sine Rule

$$\frac{\sin C}{10} = \frac{\sin 40}{7} \tag{A1}$$

$$C = 66.674...^{\circ}$$
 OR $113.325....^{\circ}$ (A1)

EITHER

B = 180 - 40 - 113.325...

$$B = 26.675...^{\circ}$$
 (A1)

area =
$$\frac{1}{2} \times 10 \times 7 \times \sin(26.675...^{\circ})$$

OR

sine rule or cosine rule to find
$$AC = 4.888...$$
 (A1)

minimum area =
$$\frac{1}{2} \times 10 \times 4.888... \times \sin(40^\circ)$$
 M1

THEN

$$=15.7 \text{ m}^2$$

Note: Award **A0M1A0** if the wrong length AC or the wrong angle B selected but used correctly finding a value of 33.5 m^2 for the area.

Total: [6 marks]

12. (a) m = -0.695 (-0.695383...); b = 4.63 (4.62974...)

A1A1

[2 marks]

(b)
$$\ln x = -0.695 (\ln 25) + 4.63$$

 $\ln x = 2.39288...$

M1 (A1)

$$x = 10.9\%$$

A1


[3 marks] Total: [5 marks]

13. (a)
$$\left(\frac{dy}{dx} = e^0 - 1\right) = 0$$

A1

[1 mark]

(b)



gradient = 0 at (0, 1) correct shape

A1 A1


Note: Award second A1 for horizontal asymptote of y = 0, and general symmetry about the y-axis.

[2 marks]

continued...

Question 13 continued

(c)



positive gradient at origin **A1** (i) **A1** correct shape

Note: Award second A1 for a single maximum in 1st quadrant and tending toward an asymptote.

positive gradient at (0, 0.75)**A1** (ii) correct shape **A1**

Note: Award second *A1* for a single minimum in 2nd quadrant, single maximum in 1st quadrant and tending toward an asymptote.

> [4 marks] Total: [7 marks]

14. (a) let *X* be the random variable "the weight of a sack of potatoes" P(X < 50)

(M1)**A1**

= 0.588 kg (0.587929...)

[2 marks]

P(X < l) = 0.25(b) (M1)49.2 kg (49.1929...)

A1 [2 marks]

(c) attempt to sum 10 independent random variables

attempt to sum
$$10$$
 independent random variables (M1)
$$Y = \sum_{i=1}^{10} X_i \sim N(498, 10 \times 0.9^2)$$
 (A1)

$$P(Y > 500) = 0.241$$

[3 marks] Total: [7 marks]

15. (a)
$$15 = 3 + 4r + 2r\theta$$
 M1 $12 = 2r(2 + \theta)$

Note: Award A1 for any reasonable working leading to expected result e,g, factorizing r.

$$r = \frac{6}{2 + \theta}$$

[2 marks]

(b) (i) attempt to use sector area to find volume
$$volume = \frac{1}{2}r^2\theta \times 1$$

$$= \frac{1}{2} \times \frac{36}{(2+\theta)^2} \times \theta \quad \left(= \frac{18\theta}{(2+\theta)^2} \right)$$
 A1

(ii)
$$\frac{dV}{d\theta} = \frac{(2+\theta)^2 \times 18 - 36\theta (2+\theta)}{(2+\theta)^4}$$
 M1A1A1

$$\frac{\mathrm{d}V}{\mathrm{d}\theta} = \frac{36 - 18\theta}{\left(2 + \theta\right)^3}$$

(iii)
$$\frac{\mathrm{d}V}{\mathrm{d}\theta} = \frac{36 - 18\theta}{(2 + \theta)^3} = 0$$

Note: Award this *M1* for simplified version equated to zero. The simplified version may have been seen in part (b)(ii).

$$\theta = 2$$

[7 marks] Total: [9 marks]

16. (a)
$$\overrightarrow{OS} = \begin{pmatrix} 300 \\ 100 \end{pmatrix} + t \begin{pmatrix} -12 \\ 15 \end{pmatrix}$$

A1

[1 mark]

(b) attempt to find the vector from L to S

$$\overrightarrow{LS} = \begin{pmatrix} 171 \\ -183 \end{pmatrix} + t \begin{pmatrix} -12 \\ 15 \end{pmatrix}$$

(M1) A1

EITHER

$$\left| \overrightarrow{LS} \right| = \sqrt{(171 - 12t)^2 + (15t - 183)^2}$$

(M1)(A1)

minimize to find *t* on GDC

(M1)

OR

S closest when
$$\overrightarrow{LS} \cdot \begin{pmatrix} -12 \\ 15 \end{pmatrix} = 0$$

(M1)

$$\begin{pmatrix} 171 \\ -183 \end{pmatrix} + t \begin{pmatrix} -12 \\ 15 \end{pmatrix}) \bullet \begin{pmatrix} -12 \\ 15 \end{pmatrix} = 0$$
$$-2052 + 144t - 2745 + 225t = 0$$

(M1)(A1)

OR

S closest when
$$\overrightarrow{LS} \cdot \begin{pmatrix} -12 \\ 15 \end{pmatrix} = 0$$

(M1)

$$\overrightarrow{LS} = \begin{pmatrix} 5k \\ 4k \end{pmatrix}$$

$$\overrightarrow{OS} = \begin{pmatrix} 129 + 5k \\ 283 + 4k \end{pmatrix}$$

(A1)

Solving simultaneously

(M1)

THEN

$$t = 13$$

A1

[6 marks]

(c) the alarm will sound

 $|\overrightarrow{LS}| = 19.2.... < 20$

A1 R1

Note: Do not award A1R0.

[2 marks] Total: [9 marks]

17. (a) attempt to use
$$V = \pi \int_a^b x^2 dy$$
 (M1)

$$x = e^{\frac{y}{6}}$$
 or any reasonable attempt to find x in terms of y (M1)

$$V = \pi \int_0^h e^{\frac{y}{3}} dy$$

Note: Correct limits must be seen for the A1 to be awarded.

$$=\pi \left[3e^{\frac{y}{3}}\right]_0^h \tag{A1}$$

Note: Condone the absence of limits for this *A1* mark.

$$=3\pi \left[e^{\frac{h}{3}}-e^{0}\right]$$

$$=3\pi \left[e^{\frac{h}{3}}-1\right]$$

Note: If the variable used in the integral is x instead of y (i.e. $V = \pi \int_0^h e^{\frac{x}{3}} dx$) and the candidate has not stated that they are interchanging x and y then award at most M1M1A0A1A1AG.

[5 marks]

(b) maximum volume when h = 9 cm max volume = 180 cm³ A1

> [2 marks] Total: [7 marks]