

© International Baccalaureate Organization 2022

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2022

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2022

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

Mathematics: analysis and approaches
Higher level
Paper 3

Thursday 12 May 2022 (morning)

1 hour

Instructions to candidates

- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Answer all the questions in the answer booklet provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics: analysis and approaches formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[55 marks]**.

Answer **all** questions in the answer booklet provided. Please start each question on a new page. Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. Solutions found from a graphic display calculator should be supported by suitable working. For example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

1. [Maximum mark: 28]

This question asks you to explore properties of a family of curves of the type $y^2 = x^3 + ax + b$ for various values of a and b , where $a, b \in \mathbb{N}$.

(a) On the same set of axes, sketch the following curves for $-2 \leq x \leq 2$ and $-2 \leq y \leq 2$, clearly indicating any points of intersection with the coordinate axes.

(i) $y^2 = x^3, x \geq 0$ [2]

(ii) $y^2 = x^3 + 1, x \geq -1$ [2]

(b) (i) Write down the coordinates of the two points of inflexion on the curve $y^2 = x^3 + 1$. [1]

(ii) By considering each curve from part (a), identify two key features that would distinguish one curve from the other. [1]

Now, consider curves of the form $y^2 = x^3 + b$, for $x \geq -\sqrt[3]{b}$, where $b \in \mathbb{Z}^+$.

(c) By varying the value of b , suggest two key features common to these curves. [2]

Next, consider the curve $y^2 = x^3 + x, x \geq 0$.

(d) (i) Show that $\frac{dy}{dx} = \pm \frac{3x^2 + 1}{2\sqrt{x^3 + x}}$, for $x > 0$. [3]

(ii) Hence deduce that the curve $y^2 = x^3 + x$ has no local minimum or maximum points. [1]

The curve $y^2 = x^3 + x$ has two points of inflexion. Due to the symmetry of the curve these points have the same x -coordinate.

(e) Find the value of this x -coordinate, giving your answer in the form $x = \sqrt{\frac{p\sqrt{3} + q}{r}}$, where $p, q, r \in \mathbb{Z}$. [7]

(This question continues on the following page)

(Question 1 continued)

$P(x, y)$ is defined to be a rational point on a curve if x and y are rational numbers.

The tangent to the curve $y^2 = x^3 + ax + b$ at a rational point P intersects the curve at another rational point Q .

Let C be the curve $y^2 = x^3 + 2$, for $x \geq -\sqrt[3]{2}$. The rational point $P(-1, -1)$ lies on C .

- (f) (i) Find the equation of the tangent to C at P . [2]
- (ii) Hence, find the coordinates of the rational point Q where this tangent intersects C , expressing each coordinate as a fraction. [2]
- (g) The point $S(-1, 1)$ also lies on C . The line $[QS]$ intersects C at a further point. Determine the coordinates of this point. [5]

Turn over

2. [Maximum mark: 27]

This question asks you to investigate conditions for the existence of complex roots of polynomial equations of degree 3 and 4.

The cubic equation $x^3 + px^2 + qx + r = 0$, where $p, q, r \in \mathbb{R}$, has roots α, β and γ .

(a) By expanding $(x - \alpha)(x - \beta)(x - \gamma)$ show that:

$$p = -(\alpha + \beta + \gamma)$$

$$q = \alpha\beta + \beta\gamma + \gamma\alpha$$

$$r = -\alpha\beta\gamma. \quad [3]$$

(b) (i) Show that $p^2 - 2q = \alpha^2 + \beta^2 + \gamma^2$. [3]

(ii) Hence show that $(\alpha - \beta)^2 + (\beta - \gamma)^2 + (\gamma - \alpha)^2 = 2p^2 - 6q$. [3]

(c) Given that $p^2 < 3q$, deduce that α, β and γ cannot all be real. [2]

Consider the equation $x^3 - 7x^2 + qx + 1 = 0$, where $q \in \mathbb{R}$.

(d) Using the result from part (c), show that when $q = 17$, this equation has at least one complex root. [2]

Noah believes that if $p^2 \geq 3q$ then α, β and γ are all real.

(e) (i) By varying the value of q in the equation $x^3 - 7x^2 + qx + 1 = 0$, determine the smallest positive integer value of q required to show that Noah is incorrect. [2]

(ii) Explain why the equation will have at least one real root for all values of q . [1]

(This question continues on the following page)

(Question 2 continued)

Now consider polynomial equations of degree 4.

The equation $x^4 + px^3 + qx^2 + rx + s = 0$, where $p, q, r, s \in \mathbb{R}$, has roots α, β, γ and δ .

In a similar way to the cubic equation, it can be shown that:

$$p = -(\alpha + \beta + \gamma + \delta)$$

$$q = \alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta$$

$$r = -(\alpha\beta\gamma + \alpha\beta\delta + \alpha\gamma\delta + \beta\gamma\delta)$$

$$s = \alpha\beta\gamma\delta.$$

- (f) (i) Find an expression for $\alpha^2 + \beta^2 + \gamma^2 + \delta^2$ in terms of p and q . [3]
- (ii) Hence state a condition in terms of p and q that would imply $x^4 + px^3 + qx^2 + rx + s = 0$ has at least one complex root. [1]
- (g) Use your result from part (f)(ii) to show that the equation $x^4 - 2x^3 + 3x^2 - 4x + 5 = 0$ has at least one complex root. [1]

The equation $x^4 - 9x^3 + 24x^2 + 22x - 12 = 0$, has one integer root.

- (h) (i) State what the result in part (f)(ii) tells us when considering this equation $x^4 - 9x^3 + 24x^2 + 22x - 12 = 0$. [1]
- (ii) Write down the integer root of this equation. [1]
- (iii) By writing $x^4 - 9x^3 + 24x^2 + 22x - 12$ as a product of one linear and one cubic factor, prove that the equation has at least one complex root. [4]

References: